



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 27, 2024 – 01:29 AM EDT

PDB ID : 7KF4
Title : Crystal structure from SARS-CoV-2 NendoU NSP15
Authors : Godoy, A.S.; Nakamura, A.M.; Pereira, H.M.; Noske, G.D.; Gawriljuk, V.O.;
Fernandes, R.S.; Oliveira, K.I.Z.; Oliva, G.
Deposited on : 2020-10-13
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4623 (2.64-2.60)
Clashscore	180529	5071 (2.64-2.60)
Ramachandran outliers	177936	5006 (2.64-2.60)
Sidechain outliers	177891	5006 (2.64-2.60)
RSRZ outliers	164620	4622 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	92% 7% .
1	B	349	93% 6% .
1	C	349	92% 7% .
1	D	349	90% 9% .
1	E	349	92% 7% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	349	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into two segments: a green segment on the left labeled '84%' and a yellow segment on the right labeled '14%'. A small red square is at the beginning of the bar, and a small black dot is at the end. A '%' symbol is positioned above the bar on the left side.</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 16984 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridylate-specific endoribonuclease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	348	2751	1768	447	526	10	0	1	0
1	B	348	2754	1770	447	526	11	0	1	0
1	C	348	2742	1763	446	522	11	0	1	0
1	D	348	2738	1759	444	524	11	0	1	0
1	E	348	2750	1767	446	526	11	0	1	0
1	F	348	2742	1763	446	522	11	0	1	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P0DTD1
A	-1	ALA	-	expression tag	UNP P0DTD1
A	0	MET	-	expression tag	UNP P0DTD1
B	-2	GLY	-	expression tag	UNP P0DTD1
B	-1	ALA	-	expression tag	UNP P0DTD1
B	0	MET	-	expression tag	UNP P0DTD1
C	-2	GLY	-	expression tag	UNP P0DTD1
C	-1	ALA	-	expression tag	UNP P0DTD1
C	0	MET	-	expression tag	UNP P0DTD1
D	-2	GLY	-	expression tag	UNP P0DTD1
D	-1	ALA	-	expression tag	UNP P0DTD1
D	0	MET	-	expression tag	UNP P0DTD1
E	-2	GLY	-	expression tag	UNP P0DTD1
E	-1	ALA	-	expression tag	UNP P0DTD1
E	0	MET	-	expression tag	UNP P0DTD1
F	-2	GLY	-	expression tag	UNP P0DTD1
F	-1	ALA	-	expression tag	UNP P0DTD1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	MET	-	expression tag	UNP P0DTD1

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	13	6	7	0	0
2	B	1	13	6	7	0	0
2	C	1	13	6	7	0	0
2	D	1	13	6	7	0	0
2	E	1	13	6	7	0	0
2	F	1	13	6	7	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	90	90	90	0	0
3	B	77	77	77	0	0
3	C	59	59	59	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	71	Total O 71 71	0	0
3	E	61	Total O 61 61	0	0
3	F	71	Total O 71 71	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uridylate-specific endoribonuclease

Chain A:  92% 7%



- Molecule 1: Uridylate-specific endoribonuclease

Chain B:  93% 6%




- Molecule 1: Uridylate-specific endoribonuclease

Chain C:  92% 7%



- Molecule 1: Uridylate-specific endoribonuclease

Chain D:  90% 9%




- Molecule 1: Uridylate-specific endoribonuclease

Chain E:  92% 7%



● Molecule 1: Uridylate-specific endoribonuclease



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.12Å 151.16Å 199.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.16 – 2.61 49.16 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.16-2.61) 99.7 (49.16-2.61)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.61Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.219 , 0.246 0.229 , 0.254	Depositor DCC
R_{free} test set	4004 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	59.3	Xtrriage
Anisotropy	0.593	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 64.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16984	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3252e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2811	0.60	0/3814
1	B	0.49	0/2814	0.61	0/3817
1	C	0.50	0/2802	0.68	0/3803
1	D	0.51	0/2798	0.68	0/3800
1	E	0.51	0/2810	0.68	0/3813
1	F	0.52	0/2802	0.72	0/3803
All	All	0.50	0/16837	0.66	0/22850

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2751	0	2742	14	0
1	B	2754	0	2749	11	0
1	C	2742	0	2730	10	0
1	D	2738	0	2712	12	0
1	E	2750	0	2738	14	0
1	F	2742	0	2730	28	0
2	A	13	0	5	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	13	0	5	1	0
2	C	13	0	5	0	0
2	D	13	0	5	0	0
2	E	13	0	5	0	0
2	F	13	0	5	0	0
3	A	90	0	0	1	0
3	B	77	0	0	0	0
3	C	59	0	0	0	0
3	D	71	0	0	0	0
3	E	61	0	0	0	0
3	F	71	0	0	2	0
All	All	16984	0	16431	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322:ILE:HB	1:A:327:ILE:HD13	1.60	0.84
1:C:322:ILE:HB	1:C:327:ILE:HD13	1.60	0.83
1:D:322:ILE:HB	1:D:327:ILE:HD13	1.62	0.80
1:C:1:SER:HB2	1:D:1:SER:HB2	1.68	0.75
1:F:265:LEU:HD23	1:F:280:ILE:HG12	1.67	0.75
1:B:148:VAL:HG13	1:B:151:LEU:HB2	1.74	0.70
1:E:230:TYR:HA	1:E:338:VAL:HG13	1.74	0.69
1:E:316:LYS:HE3	1:E:331:LEU:HD23	1.74	0.69
1:F:117:ALA:HB1	1:F:138:ARG:HE	1.59	0.68
1:D:151:LEU:HD22	1:D:186:VAL:HG11	1.77	0.66
1:A:104:MET:HE3	1:F:34:LYS:HD2	1.78	0.65
1:F:77:VAL:HG22	1:F:141:VAL:HG11	1.79	0.64
1:B:1:SER:HB2	1:E:1:SER:HB2	1.81	0.63
1:F:148:VAL:HG13	1:F:151:LEU:HB2	1.81	0.63
1:D:244:GLN:HA	1:D:288:SER:O	2.01	0.61
1:D:240:PHE:CZ	1:D:257:ARG:HD2	2.38	0.58
1:F:50:PRO:HD2	1:F:53:VAL:CG1	2.34	0.57
1:A:1:SER:HB2	1:F:1:SER:HB2	1.85	0.57
1:F:248:LEU:HD11	1:F:254:LEU:HG	1.87	0.57
1:E:98:THR:CG2	1:E:101:VAL:HB	2.35	0.56
1:F:318:VAL:HG11	1:F:341:PHE:HE1	1.70	0.56
1:F:77:VAL:CG2	1:F:141:VAL:HG11	2.37	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:313:VAL:O	1:D:333:CYS:HB2	2.08	0.54
1:A:322:ILE:HB	1:A:327:ILE:CD1	2.36	0.54
1:E:98:THR:HG23	1:E:101:VAL:HB	1.91	0.53
1:B:127:VAL:HB	1:B:130:GLN:HG3	1.91	0.51
1:E:127:VAL:HB	1:E:130:GLN:HG3	1.92	0.51
1:F:127:VAL:HB	1:F:130:GLN:HG3	1.91	0.51
1:A:127:VAL:HB	1:A:130:GLN:HG3	1.91	0.51
1:C:127:VAL:HB	1:C:130:GLN:HG3	1.91	0.51
1:C:240:PHE:CZ	1:C:257:ARG:HD2	2.46	0.51
1:F:319:LYS:HB3	1:F:326:GLU:HG2	1.92	0.50
1:A:300:ASP:HA	1:A:303:VAL:HG22	1.94	0.50
1:D:127:VAL:HB	1:D:130:GLN:HG3	1.93	0.50
1:F:135:ARG:HH21	1:F:148:VAL:HG22	1.75	0.50
1:F:31:VAL:HG23	1:F:42:LEU:HB2	1.92	0.50
1:C:266:GLU:HB3	1:C:279:PHE:HB3	1.94	0.49
1:F:337:HIS:HB3	3:F:843:HOH:O	2.11	0.49
1:D:266:GLU:HB3	1:D:279:PHE:HB3	1.94	0.49
1:A:266:GLU:HB3	1:A:279:PHE:HB3	1.95	0.48
1:C:207:SER:O	1:C:211:ILE:HG12	2.13	0.48
1:F:303:VAL:O	1:F:307:LYS:HB2	2.13	0.48
1:F:50:PRO:HD2	1:F:53:VAL:HG11	1.95	0.48
1:B:266:GLU:HB3	1:B:279:PHE:HB3	1.97	0.47
1:C:79:ILE:HG12	1:C:99:ILE:HD12	1.96	0.46
1:B:148:VAL:HG12	1:B:179:TYR:CZ	2.50	0.46
1:E:71:ILE:HD13	1:E:323:ASP:HB3	1.97	0.46
1:E:135:ARG:HH21	1:E:148:VAL:HG12	1.81	0.46
1:F:200:LEU:HD11	1:F:254:LEU:HB3	1.98	0.46
1:F:265:LEU:HD22	1:F:278:TYR:CG	2.51	0.46
1:D:322:ILE:HB	1:D:327:ILE:CD1	2.40	0.45
1:B:226:LYS:O	1:B:226:LYS:HG3	2.17	0.45
1:E:266:GLU:HB3	1:E:279:PHE:HB3	1.99	0.45
1:F:71:ILE:HD13	1:F:323:ASP:HB3	1.99	0.45
1:A:300:ASP:O	1:A:303:VAL:HG22	2.17	0.44
1:B:246:GLY:HA2	2:B:700:CIT:O1	2.18	0.44
1:A:34:LYS:HD3	1:F:104:MET:HE3	2.00	0.44
1:B:71:ILE:HD13	1:B:323:ASP:HB3	2.00	0.44
1:D:12:LYS:HE3	1:D:18:GLN:HE21	1.83	0.44
1:B:226:LYS:O	1:B:226:LYS:CG	2.65	0.44
1:D:163:ASN:ND2	1:E:279:PHE:CZ	2.86	0.44
1:F:266:GLU:HB3	1:F:279:PHE:HB3	1.99	0.43
1:F:31:VAL:HG22	1:F:43:PHE:HB3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:318:VAL:CG1	1:F:341:PHE:HE1	2.31	0.43
1:A:246:GLY:HA2	2:A:700:CIT:O2	2.18	0.43
1:C:12:LYS:HE3	1:C:18:GLN:HG3	2.00	0.43
1:A:12:LYS:HE3	1:A:18:GLN:HG3	2.00	0.43
1:F:334:LYS:HD3	1:F:334:LYS:HA	1.83	0.43
1:F:0:MET:HA	3:F:844:HOH:O	2.17	0.43
1:C:322:ILE:HB	1:C:327:ILE:CD1	2.39	0.42
1:F:75:LEU:HD13	1:F:141:VAL:HG21	2.01	0.42
1:E:121:VAL:HG22	1:E:123:PHE:CZ	2.55	0.42
1:E:10:VAL:HG21	1:E:22:VAL:HG11	1.99	0.42
1:E:79:ILE:HG23	1:E:121:VAL:HB	2.01	0.42
1:B:84:VAL:CG2	1:B:101:VAL:HG11	2.50	0.42
1:F:248:LEU:CD1	1:F:254:LEU:HG	2.50	0.42
1:A:0:MET:HA	3:A:846:HOH:O	2.20	0.41
1:A:330:MET:O	1:A:341:PHE:HA	2.20	0.41
1:D:84:VAL:CG2	1:D:101:VAL:HG11	2.51	0.41
1:E:84:VAL:CG2	1:E:101:VAL:HG11	2.51	0.41
1:B:232:PHE:O	1:B:236:VAL:HG22	2.21	0.41
1:C:84:VAL:CG2	1:C:101:VAL:HG11	2.51	0.41
1:A:84:VAL:CG2	1:A:101:VAL:HG11	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/349 (99%)	342 (99%)	5 (1%)	0	100	100
1	B	347/349 (99%)	343 (99%)	4 (1%)	0	100	100
1	C	347/349 (99%)	334 (96%)	13 (4%)	0	100	100
1	D	347/349 (99%)	335 (96%)	11 (3%)	1 (0%)	37	57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	347/349 (99%)	338 (97%)	9 (3%)	0	100	100
1	F	347/349 (99%)	338 (97%)	8 (2%)	1 (0%)	37	57
All	All	2082/2094 (99%)	2030 (98%)	50 (2%)	2 (0%)	48	70

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	226	LYS
1	D	334	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/309 (100%)	299 (97%)	10 (3%)	34	59
1	B	310/309 (100%)	297 (96%)	13 (4%)	25	48
1	C	307/309 (99%)	297 (97%)	10 (3%)	33	57
1	D	306/309 (99%)	291 (95%)	15 (5%)	21	41
1	E	309/309 (100%)	299 (97%)	10 (3%)	34	59
1	F	307/309 (99%)	283 (92%)	24 (8%)	10	21
All	All	1848/1854 (100%)	1766 (96%)	82 (4%)	23	46

All (82) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	A	147	SER
1	A	148	VAL
1	A	151	LEU
1	A	173	LYS
1	A	200	LEU
1	A	201	GLN
1	A	261	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	279	PHE
1	A	345	LEU
1	B	1	SER
1	B	38	VAL
1	B	89	LYS
1	B	147	SER
1	B	151	LEU
1	B	188	GLN
1	B	200	LEU
1	B	201	GLN
1	B	202	GLU
1	B	226	LYS
1	B	244	GLN
1	B	279	PHE
1	B	300	ASP
1	C	34	LYS
1	C	96	ILE
1	C	147	SER
1	C	148	VAL
1	C	200	LEU
1	C	261	SER
1	C	279	PHE
1	C	313	VAL
1	C	315	SER
1	C	344	LYS
1	D	1	SER
1	D	19	GLN
1	D	173	LYS
1	D	181	LYS
1	D	200	LEU
1	D	208	GLN
1	D	219	ASP
1	D	224	ARG
1	D	227	LEU
1	D	245	LEU
1	D	273	SER
1	D	279	PHE
1	D	288	SER
1	D	311	LEU
1	D	346	GLN
1	E	98	THR
1	E	147	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	151	LEU
1	E	200	LEU
1	E	245	LEU
1	E	264	GLU
1	E	279	PHE
1	E	300	ASP
1	E	314	VAL
1	E	338	VAL
1	F	1	SER
1	F	33	THR
1	F	34	LYS
1	F	89	LYS
1	F	110	LYS
1	F	128	ASP
1	F	151	LEU
1	F	163	ASN
1	F	173	LYS
1	F	202	GLU
1	F	206	ARG
1	F	215	GLU
1	F	219	ASP
1	F	227	LEU
1	F	248	LEU
1	F	273	SER
1	F	279	PHE
1	F	309	GLN
1	F	311	LEU
1	F	325	THR
1	F	331	LEU
1	F	334	LYS
1	F	335	ASP
1	F	339	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	201	GLN
1	D	18	GLN
1	D	244	GLN
1	D	337	HIS
1	F	244	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIT	A	700	-	12,12,12	1.95	6 (50%)	17,17,17	2.21	6 (35%)
2	CIT	F	700	-	12,12,12	0.25	0	17,17,17	0.43	0
2	CIT	C	700	-	12,12,12	0.25	0	17,17,17	0.43	0
2	CIT	B	700	-	12,12,12	0.17	0	17,17,17	0.42	0
2	CIT	D	700	-	12,12,12	2.01	6 (50%)	17,17,17	2.17	6 (35%)
2	CIT	E	700	-	12,12,12	1.96	6 (50%)	17,17,17	2.15	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	A	700	-	-	2/16/16/16	-
2	CIT	F	700	-	-	0/16/16/16	-
2	CIT	C	700	-	-	0/16/16/16	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	700	-	-	0/16/16/16	-
2	CIT	D	700	-	-	2/16/16/16	-
2	CIT	E	700	-	-	2/16/16/16	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	700	CIT	O5-C6	3.41	1.32	1.22
2	A	700	CIT	O5-C6	3.14	1.31	1.22
2	E	700	CIT	O4-C5	-3.10	1.20	1.30
2	E	700	CIT	O2-C1	-2.97	1.21	1.30
2	A	700	CIT	O3-C5	2.96	1.31	1.22
2	E	700	CIT	O5-C6	2.92	1.31	1.22
2	D	700	CIT	O1-C1	2.86	1.31	1.22
2	A	700	CIT	O2-C1	-2.85	1.21	1.30
2	D	700	CIT	O4-C5	-2.83	1.21	1.30
2	D	700	CIT	O2-C1	-2.77	1.21	1.30
2	A	700	CIT	O1-C1	2.68	1.30	1.22
2	E	700	CIT	O3-C5	2.66	1.30	1.22
2	D	700	CIT	O3-C5	2.57	1.30	1.22
2	A	700	CIT	O4-C5	-2.56	1.22	1.30
2	E	700	CIT	O1-C1	2.51	1.30	1.22
2	D	700	CIT	O6-C6	-2.39	1.21	1.30
2	E	700	CIT	O6-C6	-2.33	1.22	1.30
2	A	700	CIT	O6-C6	-2.24	1.22	1.30

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	700	CIT	O6-C6-C3	4.88	122.50	113.14
2	A	700	CIT	O6-C6-C3	4.86	122.46	113.14
2	D	700	CIT	O6-C6-C3	4.86	122.46	113.14
2	A	700	CIT	O5-C6-C3	-4.36	113.64	122.09
2	E	700	CIT	O5-C6-C3	-4.28	113.80	122.09
2	D	700	CIT	O5-C6-C3	-4.14	114.07	122.09
2	A	700	CIT	O1-C1-C2	-3.44	113.21	122.95
2	D	700	CIT	O3-C5-C4	-3.34	113.49	122.95
2	E	700	CIT	O1-C1-C2	-3.30	113.61	122.95
2	A	700	CIT	O3-C5-C4	-3.30	113.61	122.95
2	D	700	CIT	O1-C1-C2	-3.28	113.66	122.95
2	E	700	CIT	O3-C5-C4	-2.96	114.56	122.95
2	A	700	CIT	O2-C1-C2	2.95	123.71	114.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	700	CIT	O2-C1-C2	2.88	123.48	114.35
2	D	700	CIT	O2-C1-C2	2.75	123.07	114.35
2	D	700	CIT	O4-C5-C4	2.72	122.96	114.35
2	A	700	CIT	O4-C5-C4	2.50	122.27	114.35
2	E	700	CIT	O4-C5-C4	2.50	122.26	114.35

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	700	CIT	O1-C1-C2-C3
2	D	700	CIT	O2-C1-C2-C3
2	E	700	CIT	O2-C1-C2-C3
2	A	700	CIT	O1-C1-C2-C3
2	A	700	CIT	O2-C1-C2-C3
2	E	700	CIT	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	700	CIT	1	0
2	B	700	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	348/349 (99%)	-0.29	0 100 100	43, 65, 93, 149	1 (0%)
1	B	348/349 (99%)	-0.29	0 100 100	43, 67, 105, 210	1 (0%)
1	C	348/349 (99%)	-0.05	1 (0%) 90 88	51, 74, 97, 160	1 (0%)
1	D	348/349 (99%)	-0.13	1 (0%) 90 88	51, 69, 96, 170	1 (0%)
1	E	348/349 (99%)	-0.18	0 100 100	41, 68, 98, 238	1 (0%)
1	F	348/349 (99%)	0.16	3 (0%) 81 77	50, 82, 133, 212	1 (0%)
All	All	2088/2094 (99%)	-0.13	5 (0%) 92 90	41, 70, 105, 238	6 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	345	LEU	3.8
1	D	163	ASN	2.2
1	C	0	MET	2.1
1	F	263	PHE	2.1
1	F	232	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CIT	E	700	13/13	0.74	0.12	106,111,117,119	0
2	CIT	F	700	13/13	0.78	0.11	110,115,124,124	0
2	CIT	D	700	13/13	0.84	0.09	100,104,107,107	0
2	CIT	B	700	13/13	0.86	0.13	123,125,126,127	0
2	CIT	C	700	13/13	0.87	0.09	99,102,104,105	0
2	CIT	A	700	13/13	0.87	0.10	80,87,91,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.