



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2021 – 01:18 PM BST

PDB ID : 7NNZ  
Title : Crystal structure of Mycobacterium tuberculosis ArgF in complex with 5-methyl-4-phenylthiazol-2-amine.  
Authors : Mendes, V.; Gupta, P.; Burgess, A.; Sebastian-Perez, V.; Cattermole, E.; Meghir, C.; Blundell, T.L.  
Deposited on : 2021-02-25  
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.20  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.20


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	 95%
1	B	309	 88% 7%
1	C	309	 93% 7%
1	D	309	 92% 5%
1	E	309	 91% 6%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	309	 2% 93%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ornithine carbamoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	308	Total 2329	C 1455	N 421	O 448	S 5	0	3	0
1	B	297	Total 2248	C 1409	N 407	O 427	S 5	0	2	0
1	C	307	Total 2326	C 1452	N 422	O 446	S 6	0	2	0
1	D	298	Total 2237	C 1401	N 401	O 430	S 5	0	1	0
1	E	300	Total 2258	C 1414	N 405	O 434	S 5	0	1	0
1	F	301	Total 2281	C 1426	N 415	O 435	S 5	0	2	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P9WIT9
A	0	SER	-	expression tag	UNP P9WIT9
A	1	VAL	-	expression tag	UNP P9WIT9
B	-1	GLY	-	expression tag	UNP P9WIT9
B	0	SER	-	expression tag	UNP P9WIT9
B	1	VAL	-	expression tag	UNP P9WIT9
C	-1	GLY	-	expression tag	UNP P9WIT9
C	0	SER	-	expression tag	UNP P9WIT9
C	1	VAL	-	expression tag	UNP P9WIT9
D	-1	GLY	-	expression tag	UNP P9WIT9
D	0	SER	-	expression tag	UNP P9WIT9
D	1	VAL	-	expression tag	UNP P9WIT9
E	-1	GLY	-	expression tag	UNP P9WIT9
E	0	SER	-	expression tag	UNP P9WIT9
E	1	VAL	-	expression tag	UNP P9WIT9
F	-1	GLY	-	expression tag	UNP P9WIT9
F	0	SER	-	expression tag	UNP P9WIT9

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	1	VAL	-	expression tag	UNP P9WIT9

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	E	1	5	4	1	0	0
2	E	1	5	4	1	0	0
2	F	1	5	4	1	0	0

- Molecule 3 is 5-methyl-4-phenyl-1,3-thiazol-2-amine (three-letter code: PTZ) (formula: C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	S		
3	A	1	Total 13	C 10	N 2	S 1	0	0
3	C	1	Total 13	C 10	N 2	S 1	0	0
3	D	1	Total 13	C 10	N 2	S 1	0	0
3	E	1	Total 13	C 10	N 2	S 1	0	0
3	F	1	Total 13	C 10	N 2	S 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	351	Total 351	O 351	0	0
4	B	251	Total 251	O 251	0	0
4	C	322	Total 322	O 322	0	0
4	D	326	Total 326	O 326	0	0
4	E	290	Total 290	O 290	0	0
4	F	312	Total 312	O 312	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ornithine carbamoyltransferase




- Molecule 1: Ornithine carbamoyltransferase




- Molecule 1: Ornithine carbamoyltransferase



- Molecule 1: Ornithine carbamoyltransferase



- Molecule 1: Ornithine carbamoyltransferase





- Molecule 1: Ornithine carbamoyltransferase

Chain F: 2% 93%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.62Å 143.00Å 97.59Å 90.00° 117.58° 90.00°	Depositor
Resolution (Å)	48.96 – 1.68 81.21 – 1.68	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.96-1.68) 99.9 (81.21-1.68)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 1.68Å)	Xtrriage
Refinement program	PHENIX 1.18.2	Depositor
R, $R_{free}$	0.167 , 0.183 0.168 , 0.182	Depositor DCC
$R_{free}$ test set	12797 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.2	Xtrriage
Anisotropy	0.065	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	15641	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, PTZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/2381	0.69	0/3243
1	B	0.45	0/2296	0.68	0/3126
1	C	0.42	0/2375	0.64	0/3232
1	D	0.47	0/2281	0.66	0/3109
1	E	0.45	0/2302	0.67	0/3135
1	F	0.45	0/2328	0.66	0/3168
All	All	0.45	0/13963	0.67	0/19013

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2329	0	2290	5	0
1	B	2248	0	2222	15	0
1	C	2326	0	2296	11	0
1	D	2237	0	2193	9	0
1	E	2258	0	2227	9	0
1	F	2281	0	2257	5	0
2	A	10	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	5	0	0	0	0
2	D	15	0	0	0	0
2	E	10	0	0	0	0
2	F	5	0	0	0	0
3	A	13	0	0	0	0
3	C	13	0	0	0	0
3	D	13	0	0	1	0
3	E	13	0	0	0	0
3	F	13	0	0	0	0
4	A	351	0	0	0	0
4	B	251	0	0	0	0
4	C	322	0	0	0	0
4	D	326	0	0	0	0
4	E	290	0	0	0	0
4	F	312	0	0	0	0
All	All	15641	0	13485	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:54:ARG:HD2	1:F:70:VAL:HG11	1.70	0.72
1:C:292:ARG:HE	1:C:296:GLN:NE2	1.93	0.67
1:C:85:GLN:O	1:C:89:LYS:HG3	1.96	0.66
1:F:54:ARG:CD	1:F:70:VAL:HG11	2.26	0.64
1:D:54:ARG:HD2	1:D:70:VAL:HG11	1.81	0.63
1:B:78:LEU:HD21	1:B:87:THR:HG21	1.79	0.63
1:B:85:GLN:HG3	1:B:114:VAL:HG21	1.82	0.59
1:C:49:ASN:OD1	1:C:72:ASP:HB2	2.03	0.57
1:C:48:LYS:NZ	1:C:229:MET:SD	2.76	0.55
1:A:305:GLU:HG3	1:A:306:ARG:HH21	1.72	0.54
1:B:45:ILE:HG21	1:B:78:LEU:HD12	1.90	0.53
1:E:2:ILE:HD12	1:E:307:SER:HB3	1.90	0.53
1:D:78:LEU:HD22	1:D:87:THR:HG21	1.90	0.53
1:F:85:GLN:HG3	1:F:114:VAL:HG21	1.90	0.53
1:D:34:ARG:HH11	1:D:37:GLN:HE22	1.58	0.52
1:C:292:ARG:HH21	1:C:296:GLN:HE22	1.58	0.52
1:D:54:ARG:CD	1:D:70:VAL:HG11	2.40	0.51
1:D:78:LEU:HD21	3:D:401:PTZ:C05	2.40	0.51

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:306:ARG:HH11	1:B:306:ARG:HG3	1.75	0.50
1:B:5:PHE:CE2	1:B:129:PRO:HB2	2.46	0.50
1:B:226:TRP:CZ2	1:B:272:GLU:HA	2.46	0.50
1:D:5:PHE:CE2	1:D:129:PRO:HB2	2.47	0.49
1:C:88:ALA:HB2	1:C:111:MET:SD	2.54	0.48
1:E:154:PHE:HE1	1:E:221:LEU:HD22	1.78	0.48
1:E:49:ASN:OD1	1:E:72:ASP:HB2	2.13	0.47
1:C:5:PHE:CE2	1:C:129:PRO:HB2	2.50	0.46
1:B:265:LEU:HD21	1:B:285:VAL:HG11	1.97	0.46
1:A:305:GLU:HG3	1:A:306:ARG:NH2	2.29	0.46
1:B:2:ILE:HD12	1:B:307:SER:HB3	1.98	0.46
1:C:78:LEU:HD22	1:C:87:THR:HG21	1.99	0.45
1:B:265:LEU:HD21	1:B:285:VAL:CG1	2.47	0.44
1:A:147:ARG:NH1	1:A:201:THR:O	2.50	0.44
1:D:38:GLY:N	1:D:39:PRO:HA	2.32	0.44
1:C:54:ARG:CD	1:C:70:VAL:HG11	2.48	0.44
1:C:54:ARG:HD3	1:C:70:VAL:HG11	1.99	0.44
1:F:5:PHE:CE2	1:F:129:PRO:HB2	2.52	0.44
1:E:154:PHE:CE1	1:E:221:LEU:HD22	2.52	0.43
1:A:5:PHE:CE2	1:A:129:PRO:HB2	2.53	0.43
1:E:54:ARG:HD3	1:E:70:VAL:HG11	2.01	0.43
1:E:5:PHE:CE2	1:E:129:PRO:HB2	2.54	0.42
1:A:51:THR:O	1:A:54:ARG:HB3	2.19	0.42
1:E:226:TRP:CZ2	1:E:272:GLU:HA	2.54	0.42
1:C:292:ARG:HH21	1:C:296:GLN:NE2	2.17	0.42
1:D:154:PHE:HE1	1:D:221:LEU:HD22	1.84	0.42
1:B:78:LEU:HD22	1:B:84:LEU:CD2	2.50	0.42
1:E:110:ALA:O	1:E:114:VAL:HG22	2.20	0.42
1:E:2:ILE:CD1	1:E:307:SER:HB3	2.50	0.41
1:B:306:ARG:HG3	1:B:306:ARG:NH1	2.34	0.41
1:F:54:ARG:NH1	1:F:58:GLU:OE1	2.54	0.41
1:B:78:LEU:CD2	1:B:87:THR:HG21	2.46	0.41
1:B:240:LYS:N	1:B:241:PRO:HD2	2.36	0.40
1:D:154:PHE:CE1	1:D:221:LEU:HD22	2.56	0.40
1:B:195:GLU:O	1:B:199:GLN:HG3	2.22	0.40
1:B:132:VAL:HG21	1:B:164:SER:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/309 (100%)	303 (98%)	4 (1%)	2 (1%)	25	10
1	B	295/309 (96%)	288 (98%)	6 (2%)	1 (0%)	41	23
1	C	307/309 (99%)	301 (98%)	5 (2%)	1 (0%)	41	23
1	D	295/309 (96%)	288 (98%)	6 (2%)	1 (0%)	41	23
1	E	297/309 (96%)	288 (97%)	8 (3%)	1 (0%)	41	23
1	F	299/309 (97%)	291 (97%)	7 (2%)	1 (0%)	41	23
All	All	1802/1854 (97%)	1759 (98%)	36 (2%)	7 (0%)	34	17

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	LEU
1	D	123	LEU
1	B	123	LEU
1	C	123	LEU
1	E	123	LEU
1	F	123	LEU
1	A	265	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/243 (99%)	234 (98%)	6 (2%)	47	26

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	231/243 (95%)	228 (99%)	3 (1%)	69	54
1	C	241/243 (99%)	237 (98%)	4 (2%)	60	43
1	D	229/243 (94%)	226 (99%)	3 (1%)	69	54
1	E	233/243 (96%)	228 (98%)	5 (2%)	53	33
1	F	235/243 (97%)	230 (98%)	5 (2%)	53	33
All	All	1409/1458 (97%)	1383 (98%)	26 (2%)	59	40

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	ARG
1	A	101	ARG
1	A	128	HIS
1	A	212[A]	HIS
1	A	212[B]	HIS
1	A	278	MET
1	B	78	LEU
1	B	101	ARG
1	B	128	HIS
1	C	101	ARG
1	C	128	HIS
1	C	147	ARG
1	C	247	LEU
1	D	37	GLN
1	D	101	ARG
1	D	147	ARG
1	E	8	ASP
1	E	55	PHE
1	E	101	ARG
1	E	128	HIS
1	E	306	ARG
1	F	101	ARG
1	F	128	HIS
1	F	247	LEU
1	F	250	ARG
1	F	278	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	296	GLN
1	D	37	GLN
1	D	212	HIS
1	E	199	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	E	503	-	4,4,4	0.78	0	6,6,6	0.59	0
3	PTZ	E	501	-	9,14,14	2.50	2 (22%)	9,19,19	1.26	1 (11%)
2	PO4	D	403	-	4,4,4	1.18	0	6,6,6	0.87	0
3	PTZ	F	502	-	9,14,14	2.85	2 (22%)	9,19,19	1.04	1 (11%)
2	PO4	F	501	-	4,4,4	0.83	0	6,6,6	0.54	0
2	PO4	A	401	-	4,4,4	0.93	0	6,6,6	0.42	0
3	PTZ	A	403	-	9,14,14	2.72	2 (22%)	9,19,19	1.20	1 (11%)
3	PTZ	D	401	-	9,14,14	2.73	2 (22%)	9,19,19	1.06	1 (11%)
2	PO4	A	402	-	4,4,4	0.89	0	6,6,6	0.82	0
2	PO4	C	401	-	4,4,4	0.78	0	6,6,6	0.81	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PTZ	C	402	-	9,14,14	2.68	2 (22%)	9,19,19	1.07	1 (11%)
2	PO4	D	404	-	4,4,4	0.82	0	6,6,6	0.58	0
2	PO4	E	502	-	4,4,4	0.74	0	6,6,6	0.93	0
2	PO4	D	402	-	4,4,4	0.90	0	6,6,6	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PTZ	E	501	-	-	2/4/4/4	0/2/2/2
3	PTZ	F	502	-	-	2/4/4/4	0/2/2/2
3	PTZ	D	401	-	-	2/4/4/4	0/2/2/2
3	PTZ	C	402	-	-	2/4/4/4	0/2/2/2
3	PTZ	A	403	-	-	2/4/4/4	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	502	PTZ	C11-C04	-7.54	1.40	1.49
3	A	403	PTZ	C11-C04	-7.30	1.40	1.49
3	D	401	PTZ	C11-C04	-7.28	1.40	1.49
3	C	402	PTZ	C11-C04	-7.05	1.41	1.49
3	E	501	PTZ	C11-C04	-6.44	1.41	1.49
3	C	402	PTZ	C02-N02	3.48	1.45	1.35
3	F	502	PTZ	C02-N02	3.42	1.45	1.35
3	E	501	PTZ	C02-N02	3.37	1.45	1.35
3	A	403	PTZ	C02-N02	3.26	1.44	1.35
3	D	401	PTZ	C02-N02	3.25	1.44	1.35

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	PTZ	N02-C02-N03	3.44	127.64	123.19
3	E	501	PTZ	N02-C02-N03	3.28	127.43	123.19
3	D	401	PTZ	N02-C02-N03	3.04	127.12	123.19
3	C	402	PTZ	N02-C02-N03	3.01	127.09	123.19
3	F	502	PTZ	N02-C02-N03	2.90	126.94	123.19

There are no chirality outliers.



All (10) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	403	PTZ	N03-C04-C11-C16
3	D	401	PTZ	N03-C04-C11-C16
3	F	502	PTZ	N03-C04-C11-C12
3	F	502	PTZ	N03-C04-C11-C16
3	A	403	PTZ	N03-C04-C11-C12
3	C	402	PTZ	N03-C04-C11-C16
3	D	401	PTZ	N03-C04-C11-C12
3	C	402	PTZ	N03-C04-C11-C12
3	E	501	PTZ	N03-C04-C11-C16
3	E	501	PTZ	N03-C04-C11-C12



There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	401	PTZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	308/309 (99%)	-0.24	3 (0%) 82 85	14, 22, 39, 62	0
1	B	297/309 (96%)	-0.14	8 (2%) 54 57	15, 29, 48, 79	0
1	C	307/309 (99%)	-0.38	2 (0%) 87 89	15, 26, 38, 57	0
1	D	298/309 (96%)	-0.37	3 (1%) 82 85	15, 23, 36, 69	0
1	E	300/309 (97%)	-0.35	3 (1%) 82 85	16, 24, 40, 54	0
1	F	301/309 (97%)	-0.33	5 (1%) 70 74	16, 24, 40, 74	0
All	All	1811/1854 (97%)	-0.30	24 (1%) 77 80	14, 25, 41, 79	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	238	ARG	5.9
1	B	1	VAL	5.6
1	B	76	THR	4.4
1	B	227	THR	4.3
1	B	238	ARG	3.8
1	B	78	LEU	3.7
1	C	229	MET	3.3
1	A	0	SER	3.2
1	A	73	SER	3.0
1	E	81	ASP	3.0
1	E	307	SER	2.9
1	A	51	THR	2.8
1	F	235	GLY	2.8
1	F	0	SER	2.8
1	D	239	VAL	2.7
1	F	81	ASP	2.6
1	F	236	LEU	2.6
1	B	307	SER	2.5
1	B	75	SER	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	230	GLY	2.4
1	B	239	VAL	2.3
1	E	1	VAL	2.2
1	F	227	THR	2.0
1	D	227	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PTZ	C	402	13/13	0.79	0.23	35,44,49,55	0
2	PO4	D	404	5/5	0.92	0.26	43,54,58,64	0
3	PTZ	A	403	13/13	0.93	0.09	26,29,34,36	0
3	PTZ	D	401	13/13	0.93	0.12	24,28,34,35	0
3	PTZ	E	501	13/13	0.94	0.09	26,29,34,37	0
2	PO4	A	402	5/5	0.95	0.07	26,35,36,36	0
2	PO4	E	503	5/5	0.95	0.21	39,41,46,51	0
2	PO4	D	403	5/5	0.96	0.07	28,33,37,40	0
2	PO4	F	501	5/5	0.96	0.21	42,49,53,56	0
2	PO4	A	401	5/5	0.96	0.14	32,37,40,40	0
3	PTZ	F	502	13/13	0.96	0.08	25,28,33,36	0
2	PO4	D	402	5/5	0.97	0.18	40,40,44,44	0
2	PO4	C	401	5/5	0.97	0.11	30,39,43,43	0
2	PO4	E	502	5/5	0.99	0.06	26,27,35,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around PTZ C 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around PTZ A 403:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around PTZ D 401:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around PTZ E 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.