



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 8, 2021 – 08:06 pm GMT

PDB ID : 7OJR
Title : Bacillus subtilis phosphoglucomutase GlmM (phosphate bound)
Authors : Pathania, M.; Grundling, A.G.; Freemont, P.
Deposited on : 2021-05-17
Resolution : 3.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4 (270009), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1754 (3.10-3.02)
Clashscore	141614	1864 (3.10-3.02)
Ramachandran outliers	138981	1794 (3.10-3.02)
Sidechain outliers	138945	1793 (3.10-3.02)
RSRZ outliers	127900	1713 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 6758 atoms, of which 3375 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphoglucosamine mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	445	6753	2114	3375	572	674	18	17	3	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	449	LEU	-	expression tag	UNP O34824
A	450	VAL	-	expression tag	UNP O34824
A	451	PRO	-	expression tag	UNP O34824
A	452	ARG	-	expression tag	UNP O34824
A	453	GLY	-	expression tag	UNP O34824
A	454	SER	-	expression tag	UNP O34824
A	455	SER	-	expression tag	UNP O34824
A	456	GLY	-	expression tag	UNP O34824
A	457	LEU	-	expression tag	UNP O34824
A	458	GLU	-	expression tag	UNP O34824
A	459	HIS	-	expression tag	UNP O34824
A	460	HIS	-	expression tag	UNP O34824
A	461	HIS	-	expression tag	UNP O34824
A	462	HIS	-	expression tag	UNP O34824
A	463	HIS	-	expression tag	UNP O34824
A	464	HIS	-	expression tag	UNP O34824

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoglucosamine mutase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	134.88Å 134.88Å 69.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.29 – 3.05 48.29 – 3.05	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.29-3.05) 98.6 (48.29-3.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.67 (at 3.07Å)	Xtrriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.203 , 0.254 0.203 , 0.254	Depositor DCC
R_{free} test set	738 reflections (5.32%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtrriage
Anisotropy	0.550	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6758	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	2/3438 (0.1%)	0.84	3/4632 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	430	CYS	CB-SG	-5.63	1.72	1.81
1	A	272	GLU	CG-CD	5.43	1.60	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	333	LEU	CA-CB-CG	-5.61	102.40	115.30
1	A	288	LEU	CB-CG-CD1	-5.37	101.86	111.00
1	A	60	LEU	CB-CG-CD1	-5.12	102.29	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3378	3375	3361	77	1
2	A	5	0	0	0	0
All	All	3383	3375	3361	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:ASN:O	1:A:232:ASN:ND2	2.04	0.91
1:A:2:GLY:N	1:A:132:ASP:OD2	2.16	0.79
1:A:82:VAL:O	1:A:86:THR:HG23	1.83	0.78
1:A:23:ALA:O	1:A:60:LEU:HD23	1.85	0.77
1:A:444:GLU:N	1:A:444:GLU:OE2	2.21	0.73
1:A:393:VAL:HG21	1:A:440:VAL:HG21	1.70	0.73
1:A:86:THR:HG21	1:A:94:GLY:HA3	1.71	0.71
1:A:268:HIS:NE2	1:A:272:GLU:OE1	2.26	0.69
1:A:45:ILE:HD11	1:A:97:ILE:HD11	1.75	0.67
1:A:274:ARG:O	1:A:321:ASN:ND2	2.27	0.67
1:A:162:PHE:O	1:A:165:GLN:HG3	1.97	0.64
1:A:100:SER:HB2	1:A:242:ASP:CG	2.18	0.64
1:A:122:GLU:O	1:A:125:ALA:HB3	2.00	0.62
1:A:192:LEU:HD21	1:A:346:ILE:HD11	1.83	0.61
1:A:408:LEU:HD22	1:A:409:VAL:N	2.15	0.61
1:A:237:LEU:HD13	1:A:345:ALA:HB1	1.84	0.59
1:A:249:VAL:HG22	1:A:255:ILE:HD13	1.86	0.57
1:A:315:MET:HE2	1:A:333:LEU:HD12	1.87	0.57
1:A:57:GLU:O	1:A:61:VAL:HG23	2.06	0.56
1:A:44:LEU:N	1:A:44:LEU:HD23	2.22	0.55
1:A:102:ASN:HB3	1:A:103:PRO:CD	2.38	0.54
1:A:387:ASN:HB3	1:A:390:VAL:HB	1.89	0.54
1:A:276:LYS:HE3	1:A:277:ASP:OD2	2.08	0.53
1:A:342:LEU:HD12	1:A:342:LEU:H	1.73	0.52
1:A:104:VAL:HG21	1:A:210:LEU:HA	1.91	0.52
1:A:381:LYS:HE2	1:A:415:GLU:O	2.09	0.52
1:A:251:GLU:N	1:A:251:GLU:OE1	2.42	0.52
1:A:211:ASN:OD1	1:A:211:ASN:N	2.39	0.51
1:A:242:ASP:OD1	1:A:244:ASP:OD1	2.29	0.51
1:A:250:ASP:OD1	1:A:254:ASN:N	2.42	0.51
1:A:394:ILE:O	1:A:398:GLU:HG3	2.11	0.49
1:A:22:LEU:O	1:A:26:VAL:HG23	2.13	0.49
1:A:181:ALA:HA	1:A:216:VAL:HG23	1.94	0.49
1:A:317:LYS:HG3	1:A:318:ASP:OD2	2.13	0.48
1:A:393:VAL:HG21	1:A:440:VAL:HG11	1.96	0.48
1:A:170:ASP:OD1	1:A:172:THR:HG22	2.13	0.47
1:A:406:ARG:NH1	1:A:423:GLU:OE2	2.47	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:379:THR:H	1:A:445:MET:HB3	1.79	0.47
1:A:150:ASN:OD1	1:A:150:ASN:N	2.48	0.47
1:A:179:ASP:C	1:A:179:ASP:OD1	2.52	0.47
1:A:378:VAL:HG22	1:A:417:LEU:HA	1.98	0.46
1:A:276:LYS:HB3	1:A:321:ASN:HB2	1.97	0.46
1:A:61:VAL:O	1:A:65:LEU:HG	2.14	0.46
1:A:98:SER:OG	1:A:99:ALA:N	2.49	0.45
1:A:126:GLU:O	1:A:130:LEU:HG	2.16	0.45
1:A:384:VAL:HG12	1:A:384:VAL:O	2.16	0.45
1:A:394:ILE:HG23	1:A:407:ILE:HG21	1.99	0.45
1:A:42:LYS:HG2	1:A:42:LYS:O	2.16	0.45
1:A:376:VAL:O	1:A:417:LEU:HB2	2.16	0.45
1:A:379:THR:O	1:A:380:ASP:CB	2.65	0.44
1:A:155:GLY:O	1:A:156:GLY:C	2.55	0.44
1:A:152:TYR:CD1	1:A:154:GLU:OE1	2.71	0.44
1:A:415:GLU:HB3	1:A:416:PRO:HD2	1.98	0.44
1:A:357:LYS:HB2	1:A:362:LEU:HD21	1.99	0.43
1:A:318:ASP:HB2	1:A:320:TYR:CE2	2.53	0.43
1:A:288:LEU:HD23	1:A:288:LEU:HA	1.80	0.43
1:A:396:GLU:HG2	1:A:397:VAL:N	2.33	0.43
1:A:357:LYS:HE3	1:A:361:GLU:OE2	2.19	0.43
1:A:100:SER:O	1:A:102:ASN:N	2.48	0.43
1:A:287:ASN:O	1:A:288:LEU:C	2.57	0.43
1:A:19:THR:HB	1:A:21:GLU:OE1	2.18	0.42
1:A:36:LYS:HA	1:A:36:LYS:HD3	1.78	0.42
1:A:80:PRO:HB3	1:A:340:ASP:OD2	2.20	0.42
1:A:255:ILE:HG22	1:A:256:VAL:N	2.35	0.42
1:A:47:ARG:O	1:A:76:VAL:HA	2.20	0.42
1:A:378:VAL:HA	1:A:445:MET:HB3	2.02	0.42
1:A:107:ASN:OD1	1:A:108:GLY:N	2.41	0.42
1:A:372:LEU:O	1:A:421:MET:HA	2.20	0.41
1:A:35:THR:O	1:A:35:THR:HG23	2.20	0.41
1:A:389:LYS:O	1:A:392:ALA:HB3	2.21	0.41
1:A:277:ASP:O	1:A:279:THR:HG23	2.21	0.41
1:A:44:LEU:HD23	1:A:44:LEU:H	1.86	0.41
1:A:52:SER:O	1:A:53:GLY:C	2.60	0.40
1:A:387:ASN:CB	1:A:390:VAL:HB	2.50	0.40
1:A:249:VAL:HG22	1:A:255:ILE:CD1	2.51	0.40
1:A:24:PHE:CE1	1:A:63:GLY:HA2	2.56	0.40
1:A:342:LEU:O	1:A:346:ILE:HG13	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:GLU:OE2	1:A:140:ARG:HH22[5_555]	1.57	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	446/464 (96%)	395 (89%)	42 (9%)	9 (2%)	7 27

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	37	ASP
1	A	100	SER
1	A	390	VAL
1	A	380	ASP
1	A	416	PRO
1	A	101	HIS
1	A	118	LYS
1	A	41	PRO
1	A	80	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	366/380 (96%)	325 (89%)	41 (11%)	6 21

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	SER
1	A	28	ARG
1	A	35	THR
1	A	40	ARG
1	A	44	LEU
1	A	51	ILE
1	A	52	SER
1	A	55	MET
1	A	60	LEU
1	A	92	GLU
1	A	98	SER
1	A	100	SER
1	A	101	HIS
1	A	105	GLN
1	A	120	SER
1	A	122	GLU
1	A	142	VAL
1	A	146	LEU
1	A	165	GLN
1	A	172	THR
1	A	187	SER
1	A	200	VAL
1	A	206	SER
1	A	218	SER
1	A	244	ASP
1	A	271	SER
1	A	303	GLN
1	A	304	THR
1	A	309	ARG
1	A	312	VAL
1	A	316	LYS
1	A	330	LEU
1	A	333	LEU
1	A	338	THR
1	A	348	LEU
1	A	373	LEU
1	A	375	ASN
1	A	387	ASN
1	A	399	LYS
1	A	402	ASN
1	A	408	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/464 (95%)	-0.15	2 (0%) 92 82	26, 44, 79, 104	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLY	5.1
1	A	132	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	501	5/5	0.96	0.12	64,65,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.