



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2025 – 07:40 am BST

PDB ID : 7OWP / pdb_00007owp
Title : HsNMT1 in complex with both MyrCoA and ACE-G-(L-ORN)SFSKPR
Authors : Dian, C.; Giglione, C.; Meinnel, T.
Deposited on : 2021-06-18
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9242 (1.84-1.80)
Clashscore	180529	1080 (1.82-1.82)
Ramachandran outliers	177936	1073 (1.82-1.82)
Sidechain outliers	177891	1073 (1.82-1.82)
RSRZ outliers	164620	9241 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	4% 91% 6%
1	B	402	4% 88% 9%
2	D	9	67% 33%
2	E	9	56% 44% 56%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ORN	D	2	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycylpeptide N-tetradecanoyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	5	0
			3208	2075	541	575	17			
1	B	392	Total	C	N	O	S	0	5	0
			3216	2079	544	576	17			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	GLY	-	expression tag	UNP P30419
A	96	GLY	-	expression tag	UNP P30419
A	97	SER	-	expression tag	UNP P30419
A	98	GLU	-	expression tag	UNP P30419
B	95	GLY	-	expression tag	UNP P30419
B	96	GLY	-	expression tag	UNP P30419
B	97	SER	-	expression tag	UNP P30419
B	98	GLU	-	expression tag	UNP P30419

- Molecule 2 is a protein called ACE-GLY-ORN-SER-PHE-SER-LYS-PRO-ARG.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	9	Total	C	N	O	0	0	0
			60	38	10	12			
2	D	9	Total	C	N	O	0	0	0
			60	38	10	12			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is TETRADECANOYL-COA (CCD ID: MYA) (formula: $C_{35}H_{62}N_7O_{17}P_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	S	0	0
			63	35	7	17	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	D	1	Total	C	N	O	P	S	0	0
			63	35	7	17	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	394	Total	O	0	0
			394	394		
5	B	381	Total	O	0	1
			381	381		
5	E	4	Total	O	0	0
			4	4		
5	D	11	Total	O	0	0
			11	11		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1




- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1



- Molecule 2: ACE-GLY-ORN-SER-PHE-SER-LYS-PRO-ARG



- Molecule 2: ACE-GLY-ORN-SER-PHE-SER-LYS-PRO-ARG



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	79.32Å 178.20Å 58.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.55 – 1.81 47.55 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.0 (47.55-1.81) 99.0 (47.55-1.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 1.81Å)	Xtriage
Refinement program	PHENIX 1.17_3644	Depositor
R, R_{free}	0.155 , 0.191 0.165 , 0.201	Depositor DCC
R_{free} test set	3735 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.901	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7472	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MYA, GOL, ACE, ORN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/3299	0.47	0/4488
1	B	0.26	0/3307	0.46	0/4498
2	D	0.23	0/50	0.51	0/63
2	E	0.20	0/50	0.40	0/63
All	All	0.26	0/6706	0.47	0/9112

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3208	0	3149	20	0
1	B	3216	0	3160	24	0
2	D	60	0	55	18	0
2	E	60	0	56	8	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	B	63	0	58	1	0
4	D	63	0	58	14	0
5	A	394	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	381	0	0	3	0
5	D	11	0	0	0	0
5	E	4	0	0	1	0
All	All	7472	0	6552	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:2:ORN:HG3	4:D:501:MYA:C2	1.35	1.51
2:D:2:ORN:HG3	4:D:501:MYA:S1	1.55	1.46
2:D:2:ORN:CG	4:D:501:MYA:S1	2.09	1.36
2:D:2:ORN:CG	4:D:501:MYA:C2	2.23	1.15
2:D:2:ORN:HG3	4:D:501:MYA:H2A	1.33	1.10
2:D:2:ORN:CB	4:D:501:MYA:S1	2.41	1.07
2:D:2:ORN:CG	4:D:501:MYA:H2A	1.87	1.01
1:B:403:LEU:HD23	2:E:0:ACE:H3	1.59	0.84
2:D:2:ORN:HG3	4:D:501:MYA:H2	1.60	0.83
1:B:144[A]:ASN:ND2	5:B:601:HOH:O	2.22	0.71
1:B:232[B]:ASN:ND2	5:B:604:HOH:O	2.26	0.69
2:D:2:ORN:HB3	4:D:501:MYA:S1	2.31	0.68
1:A:351:GLN:NE2	1:B:144[B]:ASN:OD1	2.34	0.60
2:D:2:ORN:CB	4:D:501:MYA:H2A	2.33	0.58
2:E:2:ORN:N	2:E:2:ORN:HD2	2.20	0.57
1:A:470:GLY:HA2	2:D:7:PRO:HA	1.87	0.56
2:D:2:ORN:CB	4:D:501:MYA:C2	2.82	0.56
2:D:2:ORN:HB2	4:D:501:MYA:S1	2.40	0.55
1:B:307:ILE:HD13	1:B:314:LEU:HD13	1.88	0.55
1:A:410:HIS:CD2	1:A:413:HIS:H	2.25	0.55
1:B:166:ARG:HD3	5:B:860:HOH:O	2.09	0.53
1:B:458:ASN:HA	1:B:461:PHE:CE2	2.45	0.52
1:A:130:GLU:O	1:A:289:LYS:NZ	2.36	0.52
1:B:246:ASN:OD1	2:E:2:ORN:HD3	2.11	0.51
1:B:115:ARG:HD2	1:B:117:TYR:CZ	2.47	0.50
1:B:170:LYS:HA	1:B:173[B]:TYR:CE1	2.48	0.49
1:B:420:TYR:CE2	2:E:0:ACE:H1	2.48	0.48
1:A:180:TYR:OH	2:D:2:ORN:HG2	2.13	0.48
1:A:405:SER:HB3	2:D:4:PHE:HE2	1.78	0.48
1:A:410:HIS:HD2	1:A:413:HIS:H	1.59	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:1:GLY:O	2:E:3:SER:N	2.46	0.48
2:D:2:ORN:HB2	4:D:501:MYA:H2A	1.97	0.46
2:E:1:GLY:HA3	5:E:103:HOH:O	2.15	0.46
2:D:2:ORN:HB2	4:D:501:MYA:C2	2.45	0.46
1:A:282:THR:HB	1:A:474:LEU:HD22	1.97	0.45
1:B:177[A]:ASN:O	1:B:189:ARG:HD2	2.17	0.45
1:A:417:LYS:HG2	1:A:447:PHE:CD1	2.52	0.44
1:A:351:GLN:HG3	1:B:143:ASP:HB3	1.98	0.44
1:A:141:ASP:OD2	5:A:601:HOH:O	2.21	0.44
1:A:246:ASN:OD1	2:D:2:ORN:HD3	2.18	0.44
1:B:128:LEU:HA	1:B:128:LEU:HD23	1.88	0.44
1:B:401:TYR:OH	2:E:0:ACE:H2	2.18	0.44
1:A:458:ASN:HA	1:A:461:PHE:CE2	2.53	0.44
1:A:458:ASN:O	1:A:462:LEU:HG	2.18	0.43
1:B:130:GLU:O	1:B:289:LYS:HE3	2.19	0.43
1:A:138:VAL:HG22	1:A:482:LYS:HG3	2.01	0.42
1:B:183:ASP:HB3	2:E:4:PHE:CD1	2.54	0.42
1:B:195:GLU:HB3	1:B:381:ILE:HD11	2.00	0.42
1:B:320:MET:O	1:B:324:MET:HG2	2.19	0.42
1:B:258:ARG:NH1	4:B:501:MYA:O7A	2.50	0.42
1:A:474:LEU:HD23	1:A:474:LEU:HA	1.92	0.42
1:B:234:HIS:HD2	1:B:239[B]:GLU:HG3	1.83	0.41
1:A:145:ILE:HD12	1:A:272:HIS:HB3	2.01	0.41
1:B:226:ILE:HD12	1:B:248:LEU:HD13	2.02	0.41
1:A:144[B]:ASN:OD1	1:B:351:GLN:NE2	2.54	0.41
1:A:435:MET:HG3	1:A:461:PHE:CE2	2.55	0.41
1:B:170:LYS:HG3	1:B:173[B]:TYR:CZ	2.55	0.41
1:A:209[B]:GLN:HG3	5:A:736:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/402 (99%)	388 (98%)	9 (2%)	0	100	100
1	B	395/402 (98%)	386 (98%)	9 (2%)	0	100	100
2	D	6/9 (67%)	6 (100%)	0	0	100	100
2	E	6/9 (67%)	6 (100%)	0	0	100	100
All	All	804/822 (98%)	786 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/362 (96%)	343 (98%)	6 (2%)	56	42
1	B	351/362 (97%)	344 (98%)	7 (2%)	50	35
2	D	5/6 (83%)	5 (100%)	0	100	100
2	E	5/6 (83%)	5 (100%)	0	100	100
All	All	710/736 (96%)	697 (98%)	13 (2%)	52	40

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	SER
1	A	143	ASP
1	A	331	GLU
1	A	405	SER
1	A	455	LEU
1	A	474	LEU
1	B	113	SER
1	B	114	LYS
1	B	115	ARG
1	B	314	LEU
1	B	315	SER
1	B	412	THR
1	B	455	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	234	HIS
1	A	351	GLN
1	A	410	HIS
1	A	458	ASN
1	B	147	GLN
1	B	234	HIS
1	B	351	GLN
1	B	380	ASN
1	B	410	HIS
1	B	458	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ORN	D	2	2	6,7,8	0.68	0	2,7,9	0.68	0
2	ORN	E	2	2	6,7,8	0.83	0	2,7,9	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ORN	D	2	2	-	2/5/6/8	-
2	ORN	E	2	2	-	1/5/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	2	ORN	NE-CD-CG-CB
2	D	2	ORN	NE-CD-CG-CB
2	D	2	ORN	CA-CB-CG-CD

There are no ring outliers.

2 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	ORN	16	0
2	E	2	ORN	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MYA	B	501	-	57,65,65	1.32	6 (10%)	68,91,91	1.38	7 (10%)
3	GOL	B	502	-	5,5,5	1.11	0	5,5,5	0.85	0
4	MYA	D	501	-	57,65,65	2.33	8 (14%)	68,91,91	2.16	11 (16%)
3	GOL	A	501	-	5,5,5	0.95	0	5,5,5	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MYA	B	501	-	-	1/60/80/80	0/3/3/3
3	GOL	B	502	-	-	0/4/4/4	-
4	MYA	D	501	-	-	4/60/80/80	0/3/3/3
3	GOL	A	501	-	-	0/4/4/4	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	501	MYA	O2M-C2M	13.76	1.43	1.21
4	D	501	MYA	C2A-N3A	5.23	1.40	1.32
4	B	501	MYA	C2A-N3A	5.07	1.40	1.32
4	D	501	MYA	O4X-C1X	4.21	1.46	1.41
4	D	501	MYA	C3M-C2M	4.04	1.55	1.50
4	B	501	MYA	O4X-C1X	3.62	1.46	1.41
4	D	501	MYA	C2A-N1A	3.42	1.40	1.33
4	B	501	MYA	C2A-N1A	3.15	1.39	1.33
4	B	501	MYA	C5A-C4A	-2.39	1.34	1.40
4	D	501	MYA	C5A-C4A	-2.35	1.34	1.40
4	B	501	MYA	C6A-C5A	-2.29	1.34	1.43
4	D	501	MYA	C6A-C5A	-2.26	1.34	1.43
4	D	501	MYA	P3X-O7A	-2.01	1.47	1.54
4	B	501	MYA	C3M-C2M	2.00	1.52	1.50

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	501	MYA	O2M-C2M-S1	-9.97	109.67	122.61
4	D	501	MYA	O2M-C2M-C3M	-9.70	112.53	123.99
4	B	501	MYA	N3A-C2A-N1A	-6.46	118.58	128.68
4	D	501	MYA	N3A-C2A-N1A	-6.30	118.83	128.68
4	B	501	MYA	C2X-C3X-C4X	-3.59	96.86	103.22
4	D	501	MYA	C2X-C3X-C4X	-3.35	97.28	103.22
4	D	501	MYA	P2A-O3A-P1A	-3.20	121.86	132.83
4	B	501	MYA	P2A-O3A-P1A	-2.94	122.73	132.83
4	B	501	MYA	O4X-C1X-C2X	-2.66	103.04	106.93
4	B	501	MYA	O2M-C2M-C3M	-2.64	120.87	123.99
4	D	501	MYA	C2-C3-N4	-2.26	107.67	112.42
4	D	501	MYA	C4A-C5A-N7A	-2.21	107.10	109.40
4	B	501	MYA	C4A-C5A-N7A	-2.20	107.11	109.40
4	D	501	MYA	O8A-P3X-O7A	2.18	115.98	107.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	501	MYA	C5M-C4M-C3M	-2.18	105.37	113.19
4	D	501	MYA	O3X-P3X-O9A	-2.16	101.05	109.39
4	D	501	MYA	O4X-C1X-C2X	-2.10	103.85	106.93
4	B	501	MYA	C4M-C3M-C2M	-2.03	107.83	112.33

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	501	MYA	C3-C2-S1-C2M
4	D	501	MYA	C3M-C2M-S1-C2
4	D	501	MYA	C5X-O5X-P1A-O3A
4	B	501	MYA	C5X-O5X-P1A-O2A
4	D	501	MYA	C5X-O5X-P1A-O2A


There are no ring outliers.

2 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	501	MYA	1	0
4	D	501	MYA	14	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	394/402 (98%)	-0.26	15 (3%) 44 44	12, 26, 62, 92	6 (1%)
1	B	392/402 (97%)	-0.19	15 (3%) 44 44	10, 25, 70, 96	5 (1%)
2	D	7/9 (77%)	0.59	0 100 100	37, 38, 49, 49	0
2	E	7/9 (77%)	2.98	5 (71%) 0 0	77, 79, 102, 108	0
All	All	800/822 (97%)	-0.19	35 (4%) 39 38	10, 26, 70, 108	11 (1%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	1	GLY	6.8
1	A	209[A]	GLN	5.4
1	B	107	LYS	4.4
1	B	106	ALA	4.0
1	A	313[A]	HIS	3.6
1	B	108	THR	3.5
1	A	108	THR	3.4
1	B	110	GLU	3.2
2	E	7	PRO	3.1
1	A	104	GLY	2.9
1	A	317	ASN	2.9
1	A	103	GLN	2.9
1	A	109	MET	2.8
1	B	109	MET	2.8
2	E	3	SER	2.8
2	E	5	SER	2.6
1	B	116	SER	2.6
1	B	412	THR	2.6
1	A	113	SER	2.5
1	B	472	GLY	2.5
1	B	105	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	4	PHE	2.4
1	B	317	ASN	2.4
1	B	314	LEU	2.3
1	B	111	GLU	2.3
1	B	415	SER	2.2
1	A	105	PRO	2.2
1	A	256	SER	2.2
1	B	254	LEU	2.2
1	A	106	ALA	2.2
1	A	112	ALA	2.1
1	A	258	ARG	2.1
1	A	134	THR	2.1
1	A	110	GLU	2.1
1	B	133	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ORN	E	2	8/9	0.68	0.23	76,79,85,92	0
2	ORN	D	2	8/9	0.92	0.14	30,45,82,83	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MYA	B	501	63/63	0.89	0.12	13,42,53,60	0
4	MYA	D	501	63/63	0.90	0.11	15,44,66,70	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	B	502	6/6	0.95	0.07	22,23,26,31	0
3	GOL	A	501	6/6	0.97	0.06	25,29,32,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.