



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 22, 2022 – 03:11 pm BST

PDB ID : 7OZ7
Title : Crystal structure of DPP8 in complex with a 4-oxo-b-lactam based inhibitor, L84
Authors : Ross, B.; Huber, R.
Deposited on : 2021-06-26
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.30
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.30


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	898	 87% 10%
1	B	898	 88% 10%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 13406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 8.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	804	Total 6533	C 4207	N 1087	O 1214	S 25	0	0	0
1	B	804	Total 6528	C 4204	N 1086	O 1212	S 26	0	1	0

- Molecule 2 is {N}-[3-[[4-(4-bromophenyl)methyl]piperazin-1-yl]methyl]phenyl]-2-ethyl-2-methanoyl-butanamide (three-letter code: MQ2) (formula: C₂₅H₃₂BrN₃O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	Br	C	N	O		
2	A	1	Total 31	Br 1	C 25	N 3	O 2	0	0
2	B	1	Total 31	Br 1	C 25	N 3	O 2	0	0

- Molecule 3 is trimethylamine oxide (three-letter code: TMO) (formula: C₃H₉NO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	A	1	5	3	1	1	0	0
3	B	1	5	3	1	1	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	159	159	159	0	0
4	B	114	114	114	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dipeptidyl peptidase 8

Chain A:  87% 10%



- Molecule 1: Dipeptidyl peptidase 8

Chain B:  88% 10%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	149.42Å 149.42Å 269.77Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.74 – 2.60 43.74 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.74-2.60) 100.0 (43.74-2.60)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.222 , 0.260 0.225 , 0.263	Depositor DCC
R_{free} test set	5217 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	65.9	Xtrriage
Anisotropy	0.267	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.229 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13406	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MQ2, TMO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/6715	0.59	0/9111
1	B	0.37	0/6712	0.58	0/9106
All	All	0.37	0/13427	0.58	0/18217

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6533	0	6343	12	0
1	B	6528	0	6347	7	0
2	A	31	0	0	1	0
2	B	31	0	0	1	0
3	A	5	0	9	0	0
3	B	5	0	9	0	0
4	A	159	0	0	0	0
4	B	114	0	0	0	0
All	All	13406	0	12708	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:ALA:HB1	1:B:104:MET:HB3	1.81	0.63
1:B:880:LEU:HD23	2:B:901:MQ2:BR	2.62	0.55
1:A:376:GLU:HG3	1:A:397:ARG:HB2	1.89	0.54
1:A:763:MET:SD	1:A:811:VAL:HG12	2.51	0.50
1:B:65:LEU:HD21	1:B:881:LEU:CD2	2.44	0.48
1:A:882:HIS:CE1	1:A:886:GLU:HG3	2.52	0.45
1:A:880:LEU:HD23	2:A:901:MQ2:BR	2.72	0.44
1:A:118:ILE:HD12	1:A:599:LEU:CD2	2.49	0.43
1:A:500:ALA:HB1	1:A:501:PRO:HD2	1.99	0.43
1:A:882:HIS:CE1	1:B:822:PRO:HG3	2.53	0.43
1:B:882:HIS:CE1	1:B:886:GLU:HG3	2.53	0.42
1:A:596:CYS:SG	1:A:623:SER:HB2	2.60	0.42
1:B:365:ILE:HG23	1:B:366:GLN:HG3	2.02	0.42
1:A:348:ILE:HG22	1:A:350:ILE:CD1	2.50	0.42
1:B:788:ASP:O	1:B:792:THR:HG22	2.20	0.42
1:A:302:LEU:HD22	1:A:390:ALA:HB1	2.01	0.41
1:A:779:ALA:HA	1:A:829:HIS:CD2	2.55	0.41
1:A:211:CYS:HB3	1:A:232:HIS:CD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	792/898 (88%)	773 (98%)	19 (2%)	0	100	100
1	B	793/898 (88%)	768 (97%)	25 (3%)	0	100	100
All	All	1585/1796 (88%)	1541 (97%)	44 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	714/795 (90%)	708 (99%)	6 (1%)	81	92
1	B	714/795 (90%)	708 (99%)	6 (1%)	81	92
All	All	1428/1590 (90%)	1416 (99%)	12 (1%)	81	92

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	ASP
1	A	376	GLU
1	A	473	LYS
1	A	596	CYS
1	A	614	LYS
1	A	718	PHE
1	B	97	ASP
1	B	340	LYS
1	B	350	ILE
1	B	718	PHE
1	B	811	VAL
1	B	829	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	GLN
1	A	579	GLN
1	A	654	HIS
1	A	829	HIS
1	A	837	HIS
1	A	882	HIS
1	B	111	ASN
1	B	123	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TMO	A	902	-	4,4,4	6.20	1 (25%)	6,6,6	0.22	0
2	MQ2	A	901	1	31,33,33	0.63	0	41,45,45	1.58	6 (14%)
2	MQ2	B	901	1	31,33,33	0.63	0	41,45,45	1.51	5 (12%)
3	TMO	B	902	-	4,4,4	6.04	1 (25%)	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MQ2	A	901	1	-	0/24/37/37	0/3/3/3
2	MQ2	B	901	1	-	3/24/37/37	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	TMO	OAE-NAC	-12.36	1.25	1.42
3	B	902	TMO	OAE-NAC	-12.03	1.26	1.42

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	MQ2	C24-C23-C3	4.96	120.17	114.38
2	B	901	MQ2	C22-C21-C3	4.77	119.95	114.38
2	A	901	MQ2	C23-C3-C21	-4.06	103.50	110.61
2	A	901	MQ2	C22-C21-C3	3.99	119.03	114.38
2	B	901	MQ2	C23-C3-C21	-3.76	104.03	110.61
2	B	901	MQ2	C3-C4-N5	3.50	120.62	115.43
2	A	901	MQ2	C3-C4-N5	3.46	120.56	115.43
2	B	901	MQ2	C24-C23-C3	3.21	118.12	114.38
2	A	901	MQ2	O19-C4-C3	-2.35	117.42	121.21
2	B	901	MQ2	O19-C4-C3	-2.26	117.57	121.21
2	A	901	MQ2	C7-C6-C11	-2.05	117.22	119.65

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	901	MQ2	O20-C2-C3-C23
2	B	901	MQ2	C24-C23-C3-C21
2	B	901	MQ2	C22-C21-C3-C4


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	MQ2	1	0
2	B	901	MQ2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)


Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.