



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 13, 2022 – 10:14 am BST

PDB ID : 7PB2
Title : Crystal structure of JDI TCR in complex with HLA-A*11:01 bound to KRAS G12D peptide (VVVGADGVGK)
Authors : Coles, C.H.; Karuppiah, V.; Robinson, R.A.
Deposited on : 2021-07-30
Resolution : 3.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.29
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1486 (3.50-3.34)
Clashscore	141614	1572 (3.50-3.34)
Ramachandran outliers	138981	1534 (3.50-3.34)
Sidechain outliers	138945	1535 (3.50-3.34)
RSRZ outliers	127900	1395 (3.50-3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 3% 94% 5%
1	F	276	 3% 93% 6%
2	B	100	 97%
2	G	100	 5% 95%
3	C	10	 90% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	10	
4	D	203	
4	I	203	
5	E	245	
5	J	245	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 12801 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	Total	C	N	O	S	0	0	0
			2240	1392	407	432	9			
1	F	274	Total	C	N	O	S	0	0	0
			2240	1392	407	432	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	276	PRO	-	expression tag	UNP A0A583ZB34
F	276	PRO	-	expression tag	UNP A0A583ZB34

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	98	Total	C	N	O	S	0	0	0
			820	523	139	156	2			
2	G	97	Total	C	N	O	S	0	0	0
			812	517	138	155	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
G	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called KRAS G12D peptide (VVVGADGVGK).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	10	Total	C	N	O	0	0	0
			63	39	11	13			
3	H	10	Total	C	N	O	0	0	0
			63	39	11	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	ASP	GLY	engineered mutation	UNP P01111
H	6	ASP	GLY	engineered mutation	UNP P01111

- Molecule 4 is a protein called TCR alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	163	Total	C	N	O	S	0	0	0
			1293	819	208	262	4			
4	I	179	Total	C	N	O	S	0	0	0
			1414	893	229	287	5			

- Molecule 5 is a protein called TCR beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	244	Total	C	N	O	S	0	0	0
			1928	1215	338	366	9			
5	J	244	Total	C	N	O	S	0	0	0
			1928	1215	338	366	9			

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen

Chain A: 



- Molecule 1: MHC class I antigen

Chain F: 




- Molecule 2: Beta-2-microglobulin

Chain B: 




- Molecule 2: Beta-2-microglobulin

Chain G: 




- Molecule 3: KRAS G12D peptide (VVVGADGVGK)



Chain C: 






- Molecule 3: KRAS G12D peptide (VVVGADGVGK)





• Molecule 4: TCR alpha




• Molecule 4: TCR alpha



• Molecule 5: TCR beta



• Molecule 5: TCR beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	208.37Å 208.37Å 124.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	147.77 – 3.41 147.34 – 3.41	Depositor EDS
% Data completeness (in resolution range)	99.9 (147.77-3.41) 99.6 (147.34-3.41)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 3.41Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.228 , 0.254 0.235 , 0.254	Depositor DCC
R_{free} test set	1859 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	117.7	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	12801	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/2301	0.72	0/3124
1	F	0.63	0/2301	0.71	0/3124
2	B	0.64	0/843	0.70	0/1142
2	G	0.64	0/835	0.70	0/1131
3	C	0.75	0/62	0.81	0/81
3	H	0.73	0/62	0.82	0/81
4	D	0.66	0/1321	0.73	0/1794
4	I	0.66	0/1444	0.74	0/1959
5	E	0.63	0/1982	0.72	0/2698
5	J	0.63	0/1982	0.72	0/2698
All	All	0.64	0/13133	0.72	0/17832

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2240	0	2085	5	0
1	F	2240	0	2085	7	0
2	B	820	0	785	0	0
2	G	812	0	771	1	0
3	C	63	0	69	1	0
3	H	63	0	69	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	1293	0	1225	7	0
4	I	1414	0	1334	10	0
5	E	1928	0	1838	4	0
5	J	1928	0	1838	4	0
All	All	12801	0	12099	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:ARG:NH2	1:A:102:ASP:OD1	2.30	0.64
4:I:190:CYS:HA	4:I:193:ALA:HB2	1.86	0.58
4:D:35:PHE:CE1	4:D:50:ARG:HG3	2.40	0.57
4:D:105:LEU:HD21	5:E:104:LEU:HD13	1.91	0.53
1:F:62:GLN:HE22	4:I:29:ASP:HA	1.77	0.50
4:I:105:LEU:HD21	5:J:104:LEU:HD13	1.93	0.49
1:F:5:MET:HB2	1:F:168:LEU:HD13	1.94	0.48
1:A:5:MET:HB2	1:A:168:LEU:HD13	1.96	0.48
1:F:9:TYR:OH	3:H:2:VAL:HG22	2.15	0.46
4:I:57:GLN:HG3	4:I:58:ASN:N	2.30	0.46
5:E:220:GLU:N	5:E:220:GLU:OE1	2.49	0.46
1:F:165:VAL:O	1:F:169:ARG:HG3	2.17	0.44
4:I:27:THR:HB	4:I:94:LEU:HD21	2.00	0.44
1:A:165:VAL:O	1:A:169:ARG:HG3	2.18	0.44
1:F:23:ILE:HG21	2:G:54:LEU:HB3	1.98	0.44
4:I:145:PHE:CE2	4:I:164:LYS:HD2	2.53	0.44
4:D:27:THR:HB	4:D:94:LEU:HD21	2.00	0.42
4:I:84:VAL:HG21	4:I:166:VAL:HG11	2.01	0.42
4:D:142:PHE:O	4:D:178:SER:HA	2.20	0.42
4:D:141:LEU:HD21	5:E:143:THR:HG21	2.01	0.42
1:A:62:GLN:HE22	4:D:29:ASP:HA	1.85	0.41
5:J:140:GLN:HA	5:J:140:GLN:OE1	2.21	0.41
4:D:129:LEU:HD21	5:E:143:THR:O	2.21	0.41
1:F:99:TYR:CZ	3:H:3:VAL:HG12	2.56	0.41
4:I:167:LEU:O	4:I:175:LYS:HA	2.21	0.40
1:A:80:THR:HG21	3:C:10:LYS:OXT	2.21	0.40
1:F:65:ARG:NH1	4:I:29:ASP:OD1	2.54	0.40
4:I:139:VAL:HG23	5:J:129:PHE:CE2	2.56	0.40
5:J:69:LEU:HD21	5:J:75:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/276 (99%)	254 (93%)	18 (7%)	0	100	100
1	F	272/276 (99%)	255 (94%)	17 (6%)	0	100	100
2	B	96/100 (96%)	89 (93%)	7 (7%)	0	100	100
2	G	95/100 (95%)	87 (92%)	8 (8%)	0	100	100
3	C	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
3	H	8/10 (80%)	6 (75%)	2 (25%)	0	100	100
4	D	157/203 (77%)	146 (93%)	11 (7%)	0	100	100
4	I	173/203 (85%)	164 (95%)	9 (5%)	0	100	100
5	E	242/245 (99%)	229 (95%)	13 (5%)	0	100	100
5	J	242/245 (99%)	227 (94%)	15 (6%)	0	100	100
All	All	1565/1668 (94%)	1464 (94%)	101 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/233 (100%)	226 (97%)	6 (3%)	46	74
1	F	232/233 (100%)	225 (97%)	7 (3%)	41	71

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	93/95 (98%)	92 (99%)	1 (1%)	73	87
2	G	92/95 (97%)	91 (99%)	1 (1%)	73	87
3	C	6/6 (100%)	6 (100%)	0	100	100
3	H	6/6 (100%)	6 (100%)	0	100	100
4	D	148/184 (80%)	138 (93%)	10 (7%)	16	48
4	I	161/184 (88%)	152 (94%)	9 (6%)	21	53
5	E	209/210 (100%)	197 (94%)	12 (6%)	20	53
5	J	209/210 (100%)	198 (95%)	11 (5%)	22	55
All	All	1388/1456 (95%)	1331 (96%)	57 (4%)	30	62

All (57) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ARG
1	A	75	ARG
1	A	170	ARG
1	A	183	ASP
1	A	224	GLN
1	A	275	GLU
2	B	70	PHE
4	D	44	GLU
4	D	51	ARG
4	D	60	ILE
4	D	69	GLN
4	D	105	LEU
4	D	118	ASN
4	D	120	GLN
4	D	130	ARG
4	D	133	LYS
4	D	164	LYS
5	E	27	MET
5	E	38	ASP
5	E	72	GLN
5	E	104	LEU
5	E	115	THR
5	E	119	ASN
5	E	147	LEU
5	E	176	GLN
5	E	194	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	200	THR
5	E	221	ASN
5	E	223	GLU
1	F	35	ARG
1	F	170	ARG
1	F	183	ASP
1	F	218	GLN
1	F	224	GLN
1	F	264	GLU
1	F	268	LYS
2	G	70	PHE
4	I	28	ARG
4	I	44	GLU
4	I	51	ARG
4	I	60	ILE
4	I	105	LEU
4	I	118	ASN
4	I	119	ILE
4	I	133	LYS
4	I	167	LEU
5	J	9	LYS
5	J	38	ASP
5	J	69	LEU
5	J	104	LEU
5	J	115	THR
5	J	147	LEU
5	J	176	GLN
5	J	194	ARG
5	J	200	THR
5	J	227	ASP
5	J	243	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	66	ASN
1	A	87	GLN
1	A	224	GLN
4	D	57	GLN
4	D	69	GLN
4	D	83	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	118	ASN
5	E	99	GLN
5	E	105	HIS
1	F	62	GLN
1	F	66	ASN
1	F	224	GLN
4	I	57	GLN
4	I	58	ASN
4	I	118	ASN
4	I	121	ASN
5	J	65	ASN
5	J	99	GLN
5	J	105	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/276 (99%)	0.53	7 (2%) 56 54	31, 75, 111, 129	0
1	F	274/276 (99%)	0.41	9 (3%) 46 46	63, 85, 141, 167	0
2	B	98/100 (98%)	0.02	0 100 100	31, 90, 115, 124	0
2	G	97/100 (97%)	0.16	5 (5%) 27 28	99, 128, 147, 154	0
3	C	10/10 (100%)	1.01	0 100 100	72, 75, 80, 84	0
3	H	10/10 (100%)	0.56	0 100 100	74, 77, 79, 85	0
4	D	163/203 (80%)	0.91	24 (14%) 2 3	71, 90, 127, 143	0
4	I	179/203 (88%)	0.95	20 (11%) 5 7	59, 79, 117, 137	0
5	E	244/245 (99%)	0.75	19 (7%) 13 16	62, 80, 118, 143	0
5	J	244/245 (99%)	0.69	10 (4%) 37 37	54, 66, 86, 103	0
All	All	1593/1668 (95%)	0.60	94 (5%) 22 24	31, 80, 130, 167	0

All (94) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	I	194	PHE	4.2
5	E	144	LEU	3.9
1	F	200	THR	3.7
5	J	209	PHE	3.5
1	F	257	TYR	3.4
4	D	91	PHE	3.3
4	I	145	PHE	3.2
5	E	156	VAL	3.2
4	D	79	ILE	3.2
4	D	90	TYR	3.2
5	J	43	LEU	3.1
5	E	43	LEU	3.1
4	I	21	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	I	75	PHE	3.0
5	E	240	ALA	2.9
5	J	160	TRP	2.9
4	D	66	TRP	2.9
4	D	145	PHE	2.9
5	E	209	PHE	2.9
4	I	142	PHE	2.8
5	J	164	GLY	2.8
4	D	59	GLU	2.8
1	F	248	VAL	2.8
4	I	66	TRP	2.8
5	E	160	TRP	2.8
4	I	120	GLN	2.7
4	D	49	ILE	2.7
4	I	49	ILE	2.7
4	I	37	TYR	2.7
4	I	141	LEU	2.6
4	I	35	PHE	2.6
1	A	172	LEU	2.6
4	I	91	PHE	2.6
4	D	10	GLU	2.6
4	I	113	LEU	2.6
1	F	247	VAL	2.6
5	J	156	VAL	2.6
4	D	37	TYR	2.5
1	F	201	LEU	2.5
2	G	2	GLN	2.5
5	E	195	LEU	2.5
1	F	194	ILE	2.5
4	I	107	PHE	2.5
1	F	78	LEU	2.5
1	F	199	ALA	2.5
5	E	147	LEU	2.5
5	E	191	LEU	2.5
4	I	159	VAL	2.5
4	D	68	PHE	2.4
1	A	179	LEU	2.4
2	G	69	GLU	2.4
4	D	141	LEU	2.4
1	A	130	LEU	2.4
4	I	111	THR	2.4
5	E	5	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	201	LEU	2.3
2	G	82	VAL	2.3
5	E	201	PHE	2.3
5	J	215	PHE	2.3
4	D	45	LEU	2.3
4	D	164	LYS	2.3
4	I	77	PHE	2.3
5	E	196	ARG	2.3
4	D	35	PHE	2.3
5	E	205	PRO	2.3
5	E	90	PHE	2.3
5	E	128	VAL	2.3
1	A	160	LEU	2.2
5	E	241	TRP	2.2
4	I	79	ILE	2.2
4	D	93	ALA	2.2
1	A	156	GLN	2.2
4	D	107	PHE	2.2
4	D	180	VAL	2.1
4	D	33	TYR	2.1
4	D	127	TYR	2.1
4	D	60	ILE	2.1
5	J	213	VAL	2.1
5	J	147	LEU	2.1
5	J	195	LEU	2.1
4	I	126	VAL	2.1
5	E	162	VAL	2.1
4	D	34	LEU	2.1
5	E	215	PHE	2.1
5	E	145	VAL	2.1
1	A	81	LEU	2.0
4	D	21	LEU	2.0
2	G	47	GLU	2.0
5	J	148	ALA	2.0
4	D	36	TRP	2.0
2	G	39	LEU	2.0
1	F	109	PHE	2.0
4	I	117	PRO	2.0
4	D	80	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.