



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 28, 2025 – 10:12 am BST

PDB ID : 7PEI / pdb_00007pei
Title : OXA-48_E185A/R186A/R206A with carbamylated Lys73 in complex with chloride ions at neutral pH
Authors : Zhou, Q.; He, Y.; Jin, Y.
Deposited on : 2021-08-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	260	 89% 7%
1	BBB	260	 89% 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8196 atoms, of which 3921 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	242	Total	C	H	N	O	S	47	2	0
			3917	1262	1939	344	364	8			
1	BBB	242	Total	C	H	N	O	S	51	5	0
			3940	1270	1950	345	367	8			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	6	MET	-	initiating methionine	UNP A0A482LRD5
AAA	7	HIS	-	expression tag	UNP A0A482LRD5
AAA	8	HIS	-	expression tag	UNP A0A482LRD5
AAA	9	HIS	-	expression tag	UNP A0A482LRD5
AAA	10	HIS	-	expression tag	UNP A0A482LRD5
AAA	11	HIS	-	expression tag	UNP A0A482LRD5
AAA	12	HIS	-	expression tag	UNP A0A482LRD5
AAA	13	SER	-	expression tag	UNP A0A482LRD5
AAA	14	ALA	-	expression tag	UNP A0A482LRD5
AAA	15	GLY	-	expression tag	UNP A0A482LRD5
AAA	16	GLU	-	expression tag	UNP A0A482LRD5
AAA	17	ASN	-	expression tag	UNP A0A482LRD5
AAA	18	LEU	-	expression tag	UNP A0A482LRD5
AAA	19	TYR	-	expression tag	UNP A0A482LRD5
AAA	20	PHE	-	expression tag	UNP A0A482LRD5
AAA	21	GLN	-	expression tag	UNP A0A482LRD5
AAA	22	GLY	-	expression tag	UNP A0A482LRD5
AAA	185	ALA	GLU	engineered mutation	UNP A0A482LRD5
AAA	186	ALA	ARG	engineered mutation	UNP A0A482LRD5
AAA	206	ALA	ARG	engineered mutation	UNP A0A482LRD5
BBB	6	MET	-	initiating methionine	UNP A0A482LRD5
BBB	7	HIS	-	expression tag	UNP A0A482LRD5
BBB	8	HIS	-	expression tag	UNP A0A482LRD5
BBB	9	HIS	-	expression tag	UNP A0A482LRD5
BBB	10	HIS	-	expression tag	UNP A0A482LRD5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
BBB	11	HIS	-	expression tag	UNP A0A482LRD5
BBB	12	HIS	-	expression tag	UNP A0A482LRD5
BBB	13	SER	-	expression tag	UNP A0A482LRD5
BBB	14	ALA	-	expression tag	UNP A0A482LRD5
BBB	15	GLY	-	expression tag	UNP A0A482LRD5
BBB	16	GLU	-	expression tag	UNP A0A482LRD5
BBB	17	ASN	-	expression tag	UNP A0A482LRD5
BBB	18	LEU	-	expression tag	UNP A0A482LRD5
BBB	19	TYR	-	expression tag	UNP A0A482LRD5
BBB	20	PHE	-	expression tag	UNP A0A482LRD5
BBB	21	GLN	-	expression tag	UNP A0A482LRD5
BBB	22	GLY	-	expression tag	UNP A0A482LRD5
BBB	185	ALA	GLU	engineered mutation	UNP A0A482LRD5
BBB	186	ALA	ARG	engineered mutation	UNP A0A482LRD5
BBB	206	ALA	ARG	engineered mutation	UNP A0A482LRD5

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	H	O	2	0
			14	3	8	3		
2	AAA	1	Total	C	H	O	2	0
			14	3	8	3		
2	BBB	1	Total	C	H	O	2	0
			14	3	8	3		
2	BBB	1	Total	C	H	O	2	0
			14	3	8	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	O	S	0	0
			5	4	1		
3	AAA	1	Total	O	S	0	0
			5	4	1		
3	AAA	1	Total	O	S	0	0
			5	4	1		
3	BBB	1	Total	O	S	0	0
			5	4	1		
3	BBB	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	145	Total	O	0	1
			146	146		
4	BBB	112	Total	O	0	0
			112	112		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase

Chain AAA:  89% 7%



- Molecule 1: Beta-lactamase

Chain BBB:  89% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	146.19Å 146.19Å 54.89Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.69 – 1.90 19.69 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.69-1.90) 99.9 (19.69-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.177 , 0.211 0.176 , 0.210	Depositor DCC
R_{free} test set	2657 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	0.838	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.030 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	8196	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.16	0/2020	1.30	5/2733 (0.2%)
1	BBB	1.14	0/2042	1.30	3/2763 (0.1%)
All	All	1.15	0/4062	1.30	8/5496 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	132	GLU	CB-CA-C	-5.82	101.75	110.88
1	BBB	151	GLY	CA-C-O	-5.40	117.06	121.76
1	AAA	119	VAL	CA-C-N	5.35	125.56	120.43
1	AAA	119	VAL	C-N-CA	5.35	125.56	120.43
1	AAA	131	GLY	CA-C-N	5.32	127.36	120.44
1	AAA	131	GLY	C-N-CA	5.32	127.36	120.44
1	BBB	119	VAL	CA-C-N	5.13	125.36	120.43
1	BBB	119	VAL	C-N-CA	5.13	125.36	120.43

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1978	1939	1932	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	BBB	1990	1950	1945	4	0
2	AAA	12	16	16	2	0
2	BBB	12	16	16	1	0
3	AAA	15	0	0	1	0
3	BBB	10	0	0	0	0
4	AAA	146	0	0	2	0
4	BBB	112	0	0	2	0
All	All	4275	3921	3909	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:BBB:302:GOL:O1	4:BBB:401:HOH:O	2.03	0.76
3:AAA:305:SO4:O4	4:AAA:401:HOH:O	2.07	0.70
1:AAA:134:ARG:O	1:AAA:138:MET:HG2	2.00	0.62
2:AAA:302:GOL:H11	4:AAA:486:HOH:O	2.07	0.54
1:AAA:204:ILE:HG21	2:AAA:302:GOL:H12	1.92	0.50
1:AAA:32:ASN:ND2	1:AAA:57:ASN:HD22	2.13	0.46
1:BBB:140[B]:HIS:CD2	1:BBB:147:GLU:OE1	2.67	0.46
1:BBB:32:ASN:ND2	1:BBB:57:ASN:HD22	2.15	0.45
1:BBB:106:ASN:ND2	4:BBB:411:HOH:O	2.48	0.44
1:BBB:120:VAL:N	1:BBB:121:PRO:HD2	2.32	0.44
1:AAA:120:VAL:N	1:AAA:121:PRO:HD2	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	241/260 (93%)	236 (98%)	5 (2%)	0	100	100
1	BBB	244/260 (94%)	239 (98%)	5 (2%)	0	100	100
All	All	485/520 (93%)	475 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	209/222 (94%)	209 (100%)	0	100	100
1	BBB	212/222 (96%)	212 (100%)	0	100	100
All	All	421/444 (95%)	421 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	BBB	73	1	9,11,12	0.98	1 (11%)	5,12,14	0.85	0
1	KCX	AAA	73	1	9,11,12	0.95	1 (11%)	5,12,14	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	BBB	73	1	-	0/9/10/12	-
1	KCX	AAA	73	1	-	0/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	BBB	73	KCX	CX-NZ	2.21	1.39	1.35
1	AAA	73	KCX	CX-NZ	2.17	1.38	1.35

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	AAA	301	-	5,5,5	0.14	0	5,5,5	0.36	0
2	GOL	BBB	302	-	5,5,5	0.05	0	5,5,5	0.21	0
3	SO4	BBB	303	-	4,4,4	0.43	0	6,6,6	0.20	0
2	GOL	BBB	301	-	5,5,5	0.18	0	5,5,5	0.26	0
3	SO4	AAA	303	-	4,4,4	0.27	0	6,6,6	0.19	0
3	SO4	AAA	305	-	4,4,4	0.38	0	6,6,6	0.11	0
2	GOL	AAA	302	-	5,5,5	0.18	0	5,5,5	0.52	0
3	SO4	BBB	304	-	4,4,4	0.44	0	6,6,6	0.09	0
3	SO4	AAA	304	-	4,4,4	0.26	0	6,6,6	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	AAA	302	-	-	2/4/4/4	-
2	GOL	BBB	301	-	-	2/4/4/4	-
2	GOL	AAA	301	-	-	2/4/4/4	-
2	GOL	BBB	302	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	302	GOL	C1-C2-C3-O3
2	BBB	301	GOL	C1-C2-C3-O3
2	BBB	302	GOL	C1-C2-C3-O3
2	AAA	301	GOL	C1-C2-C3-O3
2	AAA	302	GOL	O2-C2-C3-O3
2	BBB	302	GOL	O2-C2-C3-O3
2	AAA	301	GOL	O2-C2-C3-O3
2	BBB	301	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	BBB	302	GOL	1	0
3	AAA	305	SO4	1	0
2	AAA	302	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	241/260 (92%)	-0.75	0 100 100	17, 35, 58, 81	3 (1%)
1	BBB	241/260 (92%)	-0.50	0 100 100	18, 40, 61, 80	9 (3%)
All	All	482/520 (92%)	-0.62	0 100 100	17, 37, 61, 81	12 (2%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	AAA	73	12/13	0.96	0.06	24,29,33,38	1
1	KCX	BBB	73	12/13	0.96	0.07	30,34,43,43	1

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	AAA	304	5/5	0.82	0.12	42,42,51,56	5
2	GOL	AAA	302	6/6	0.86	0.15	38,45,49,49	14
2	GOL	BBB	302	6/6	0.88	0.13	31,44,49,49	14
3	SO4	BBB	304	5/5	0.88	0.14	41,47,56,56	5
2	GOL	BBB	301	6/6	0.92	0.18	45,48,56,56	14
3	SO4	BBB	303	5/5	0.94	0.08	38,42,46,49	5
2	GOL	AAA	301	6/6	0.95	0.10	41,46,52,52	14
3	SO4	AAA	303	5/5	0.96	0.07	38,42,46,52	5
3	SO4	AAA	305	5/5	0.96	0.06	41,49,55,57	5

6.5 Other polymers [i](#)

There are no such residues in this entry.