



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 11, 2022 – 10:06 am BST

PDB ID : 7QK9  
Title : Crystal structure of the ALDH1A3-ATP complex  
Authors : Castellvi, A.; Farres, J.  
Deposited on : 2021-12-17  
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	489	 15% 93% 7%
1	B	489	 11% 92% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PEG	B	904	-	-	-	X
4	GOL	A	603	-	-	X	-
4	GOL	B	901	-	-	-	X

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 8199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldehyde dehydrogenase family 1 member A3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	489	Total	C	N	O	S	0	3	0
			3721	2375	637	688	21			
1	B	489	Total	C	N	O	S	0	5	0
			3722	2377	635	688	22			

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is O-ACETALDEHYDYL-HEXAETHYLENE GLYCOL (three-letter code: P4C) (formula:  $C_{14}H_{28}O_8$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 13 8 5	0	0
5	B	1	Total C O 13 8 5	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	308	Total	O	0	3
			311	311		
7	B	303	Total	O	0	3
			306	306		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldehyde dehydrogenase family 1 member A3



- Molecule 1: Aldehyde dehydrogenase family 1 member A3





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.11Å 90.01Å 158.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.19 – 1.78 48.19 – 1.78	Depositor EDS
% Data completeness (in resolution range)	99.2 (48.19-1.78) 99.2 (48.19-1.78)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.03 (at 1.78Å)	Xtriage
Refinement program	BUSTER 2.10.3 (18-SEP-2020)	Depositor
R, $R_{free}$	0.193 , 0.215 0.198 , 0.223	Depositor DCC
$R_{free}$ test set	5474 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.2	Xtriage
Anisotropy	0.678	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8199	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: P4C, GOL, ATP, PEG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3799	0.57	0/5157
1	B	0.42	0/3800	0.57	0/5160
All	All	0.41	0/7599	0.57	0/10317

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3721	0	3673	17	0
1	B	3722	0	3660	23	0
2	A	31	0	12	0	0
2	B	31	0	12	0	0
3	A	14	0	20	0	0
3	B	14	0	20	0	0
4	A	12	0	16	6	0
4	B	6	0	8	1	0
5	A	13	0	14	0	0
5	B	13	0	14	2	0
6	B	5	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	311	0	0	1	0
7	B	306	0	0	0	0
All	All	8199	0	7449	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:377:ALA:HB2	1:B:405:MET:HE1	1.68	0.76
1:B:177:ILE:HD12	1:B:253:ILE:HD11	1.71	0.71
1:A:505:ILE:HG12	1:B:465:ILE:HD12	1.84	0.59
4:A:603:GOL:C3	1:B:266:LYS:HZ1	2.17	0.57
1:B:269:ALA:HB1	1:B:275:LYS:HG3	1.88	0.55
1:B:377:ALA:HB2	1:B:405:MET:CE	2.37	0.54
1:A:373:LYS:HE2	1:A:379:LEU:HD22	1.90	0.54
1:A:266:LYS:NZ	4:A:603:GOL:H2	2.24	0.53
1:A:161:ASN:ND2	1:A:506:LYS:NZ	2.57	0.51
1:A:214:LEU:HD21	1:A:234:PRO:HG3	1.93	0.51
4:A:603:GOL:C3	1:B:266:LYS:NZ	2.74	0.51
1:B:373:LYS:HE2	1:B:379:LEU:HD22	1.92	0.50
1:A:177:ILE:HD12	1:A:253:ILE:HD11	1.94	0.50
1:A:465:ILE:HD12	1:B:505:ILE:HG12	1.94	0.49
1:B:368:LEU:HD13	1:B:412:ILE:HG12	1.94	0.48
1:B:29:VAL:HG21	1:B:211:LEU:HD22	1.96	0.47
1:B:325:GLN:NE2	1:B:325:GLN:H	2.14	0.46
1:B:310:GLN:HG3	1:B:353:GLN:HG3	1.98	0.46
1:A:377:ALA:HB2	1:A:405:MET:CE	2.46	0.45
1:A:377:ALA:HB2	1:A:405:MET:HE3	1.98	0.45
1:A:266:LYS:HZ1	4:A:603:GOL:H2	1.81	0.45
1:B:253:ILE:HG23	1:B:277:VAL:HG13	1.98	0.44
1:A:263:LYS:HE3	1:A:482:MET:HE1	2.00	0.44
1:B:154:LYS:NZ	4:B:901:GOL:H12	2.33	0.44
1:B:343:VAL:HG22	1:B:353:GLN:HB3	1.99	0.44
1:B:35:PHE:CZ	1:B:38:ASN:HA	2.53	0.43
1:B:461:GLY:HA3	1:B:478:GLY:O	2.19	0.43
1:A:322:VAL:HG21	1:A:330:PHE:CD1	2.54	0.43
1:B:177:ILE:CD1	1:B:253:ILE:HD11	2.46	0.42
1:A:244:ILE:HG23	1:A:250:ILE:HD13	2.01	0.42
4:A:603:GOL:H31	1:B:266:LYS:NZ	2.35	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:LEU:HD21	1:B:234:PRO:HG3	2.00	0.42
1:A:269:ALA:HB1	1:A:275:LYS:HG3	2.01	0.41
4:A:603:GOL:H31	1:B:266:LYS:HZ1	1.85	0.41
1:A:179:PRO:HD3	1:A:256:THR:HB	2.01	0.41
1:A:161:ASN:HD21	1:A:506:LYS:NZ	2.18	0.41
1:A:79:GLN:NE2	7:A:710:HOH:O	2.54	0.40
1:B:341:ARG:HA	1:B:342:PRO:HD3	1.91	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	490/489 (100%)	474 (97%)	15 (3%)	1 (0%)	47	32
1	B	492/489 (101%)	475 (96%)	17 (4%)	0	100	100
All	All	982/978 (100%)	949 (97%)	32 (3%)	1 (0%)	51	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	438	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/405 (95%)	376 (98%)	9 (2%)	50	34
1	B	383/405 (95%)	372 (97%)	11 (3%)	42	25
All	All	768/810 (95%)	748 (97%)	20 (3%)	46	29

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	GLU
1	A	102	GLN
1	A	133	ASP
1	A	196	CYS
1	A	203	LEU
1	A	348	ASP
1	A	350	LYS
1	A	404	ASN
1	A	413	PHE
1	B	28	GLU
1	B	29	VAL
1	B	102	GLN
1	B	118	GLU
1	B	121	ASP
1	B	133	ASP
1	B	196	CYS
1	B	203	LEU
1	B	234	PRO
1	B	252	LYS
1	B	413	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	HIS
1	A	161	ASN
1	B	152	GLN
1	B	325	GLN
1	B	356	GLN

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEG	B	903	-	6,6,6	0.09	0	5,5,5	0.18	0
3	PEG	B	904	-	6,6,6	0.24	0	5,5,5	0.19	0
5	P4C	A	606	-	12,12,21	0.59	0	11,11,20	0.59	0
5	P4C	B	906	-	12,12,21	0.56	0	11,11,20	0.73	1 (9%)
3	PEG	A	602	-	6,6,6	0.11	0	5,5,5	0.18	0
6	SO4	B	905	-	4,4,4	0.18	0	6,6,6	0.10	0
4	GOL	B	901	-	5,5,5	0.03	0	5,5,5	0.25	0
2	ATP	B	902	-	26,33,33	0.60	0	31,52,52	0.77	1 (3%)
4	GOL	A	603	-	5,5,5	0.09	0	5,5,5	0.35	0
4	GOL	A	604	-	5,5,5	0.03	0	5,5,5	0.17	0
3	PEG	A	605	-	6,6,6	0.23	0	5,5,5	0.21	0
2	ATP	A	601	-	26,33,33	0.65	0	31,52,52	0.72	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	B	903	-	-	0/4/4/4	-
3	PEG	B	904	-	-	2/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	P4C	A	606	-	-	3/9/10/19	-
5	P4C	B	906	-	-	4/9/10/19	-
3	PEG	A	602	-	-	1/4/4/4	-
4	GOL	B	901	-	-	1/4/4/4	-
2	ATP	B	902	-	-	3/18/38/38	0/3/3/3
4	GOL	A	603	-	-	0/4/4/4	-
4	GOL	A	604	-	-	0/4/4/4	-
3	PEG	A	605	-	-	3/4/4/4	-
2	ATP	A	601	-	-	2/18/38/38	0/3/3/3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	902	ATP	C5-C6-N6	2.30	123.85	120.35
5	B	906	P4C	O22-C21-C20	-2.18	119.82	126.39
2	A	601	ATP	C5-C6-N6	2.17	123.64	120.35

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	606	P4C	O13-C14-C15-O16
3	B	904	PEG	O1-C1-C2-O2
5	A	606	P4C	O10-C11-C12-O13
5	B	906	P4C	O13-C14-C15-O16
3	A	605	PEG	O2-C3-C4-O4
5	B	906	P4C	O10-C11-C12-O13
3	A	605	PEG	O1-C1-C2-O2
2	A	601	ATP	PB-O3A-PA-O5'
5	B	906	P4C	C17-C18-O19-C20
2	A	601	ATP	PB-O3B-PG-O3G
5	A	606	P4C	C17-C18-O19-C20
3	A	605	PEG	C4-C3-O2-C2
3	B	904	PEG	C4-C3-O2-C2
2	B	902	ATP	PG-O3B-PB-O2B
2	B	902	ATP	PB-O3A-PA-O2A
4	B	901	GOL	O1-C1-C2-C3
2	B	902	ATP	PG-O3B-PB-O1B
3	A	602	PEG	C1-C2-O2-C3

*Continued on next page...*

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	906	P4C	C14-C15-O16-C17

There are no ring outliers.


3 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	906	P4C	2	0
4	B	901	GOL	1	0
4	A	603	GOL	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	489/489 (100%)	0.88	71 (14%) <b>2</b> <b>2</b>	28, 41, 67, 82	0
1	B	489/489 (100%)	0.71	55 (11%) <b>5</b> <b>5</b>	28, 41, 71, 87	0
All	All	978/978 (100%)	0.80	126 (12%) <b>3</b> <b>3</b>	28, 41, 70, 87	0

All (126) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	389	LYS	7.9
1	A	386	MET	6.6
1	B	21	PRO	6.3
1	B	388	ASP	6.3
1	B	343	VAL	5.8
1	A	23	PRO	5.6
1	A	343	VAL	5.4
1	B	25	ARG	5.4
1	B	377	ALA	5.3
1	A	20	LEU	5.3
1	B	24	ILE	5.2
1	A	389	LYS	5.1
1	A	22	ARG	4.9
1	A	21	PRO	4.9
1	A	383	GLY	4.8
1	A	337	TYR	4.8
1	A	377	ALA	4.7
1	B	404	ASN	4.5
1	B	352	GLU	4.4
1	A	388	ASP	4.4
1	A	348	ASP	4.2
1	A	406	ARG	4.2
1	B	372	GLY	4.1
1	B	20	LEU	4.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	347	PHE	3.9
1	B	349	VAL	3.8
1	A	382	GLY	3.8
1	B	327	TYR	3.8
1	A	376	GLY	3.8
1	A	276[A]	ARG	3.7
1	B	22	ARG	3.7
1	B	382	GLY	3.7
1	B	346	PRO	3.7
1	A	404	ASN	3.6
1	A	58	GLU	3.5
1	A	390	GLY	3.5
1	A	24	ILE	3.5
1	A	350	LYS	3.5
1	A	379	LEU	3.5
1	B	383	GLY	3.5
1	B	338	ALA	3.4
1	B	390	GLY	3.3
1	A	26	ASN	3.3
1	A	362	PHE	3.3
1	A	365	ILE	3.3
1	B	353	GLN	3.2
1	A	156	ILE	3.2
1	A	253	ILE	3.2
1	A	185	LEU	3.2
1	A	374	LYS	3.2
1	A	492	TYR	3.2
1	A	493	ALA	3.1
1	B	344	GLY	3.1
1	A	349	VAL	3.1
1	A	384	SER	3.1
1	B	398	PHE	3.0
1	A	387	GLU	3.0
1	B	23	PRO	3.0
1	B	361	GLN	3.0
1	B	492	TYR	3.0
1	B	402	THR	3.0
1	A	385	ALA	2.9
1	B	26	ASN	2.9
1	A	335	VAL	2.9
1	B	386	MET	2.9
1	A	254	ALA	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	391	LEU	2.8
1	B	153	GLY	2.8
1	B	500	VAL	2.8
1	A	153	GLY	2.7
1	A	347	PHE	2.7
1	A	279	LEU	2.7
1	A	144	PHE	2.7
1	A	166	THR	2.7
1	A	494	LEU	2.7
1	A	475	ALA	2.6
1	B	490	GLY	2.6
1	A	500	VAL	2.6
1	A	261	VAL	2.6
1	A	397	VAL	2.6
1	B	156	ILE	2.6
1	B	39	GLU	2.6
1	A	255	PHE	2.6
1	B	28	GLU	2.6
1	A	369	ILE	2.5
1	A	327	TYR	2.5
1	A	490	GLY	2.5
1	A	498	THR	2.5
1	B	393	ILE	2.5
1	B	276[A]	ARG	2.5
1	A	372	GLY	2.5
1	B	342	PRO	2.5
1	B	387	GLU	2.5
1	A	90	ARG	2.5
1	A	351	THR	2.5
1	B	341	ARG	2.4
1	B	439	LEU	2.4
1	A	179	PRO	2.4
1	B	351	THR	2.4
1	A	143	TYR	2.4
1	A	342	PRO	2.4
1	B	55	SER	2.4
1	A	314	CYS	2.4
1	A	497	TYR	2.3
1	A	147	TRP	2.3
1	B	364	LYS	2.3
1	A	237	GLY	2.3
1	B	379	LEU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	348	ASP	2.3
1	A	371	SER	2.3
1	A	489	LEU	2.2
1	A	407	ILE	2.2
1	B	277	VAL	2.2
1	B	112	ALA	2.2
1	B	413	PHE	2.1
1	B	44	LYS	2.1
1	B	357	ILE	2.1
1	A	189	TRP	2.1
1	B	478	GLY	2.1
1	A	25	ARG	2.1
1	A	332	ARG	2.1
1	A	437	TYR	2.0
1	B	42	GLU	2.0
1	A	27	LEU	2.0
1	A	307	PHE	2.0
1	B	295	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PEG	B	904	7/7	0.51	0.47	72,72,72,72	0
4	GOL	A	603	6/6	0.64	0.40	57,58,59,59	0
4	GOL	B	901	6/6	0.68	0.51	85,86,86,86	0
3	PEG	A	605	7/7	0.71	0.30	51,54,56,56	0
5	P4C	A	606	13/22	0.78	0.19	55,60,64,65	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	604	6/6	0.79	0.19	96,97,97,97	0
2	ATP	B	902	31/31	0.84	0.14	36,38,71,71	0
2	ATP	A	601	31/31	0.86	0.13	38,40,65,66	0
5	P4C	B	906	13/22	0.86	0.13	53,58,62,62	0
6	SO4	B	905	5/5	0.87	0.14	108,108,108,108	0
3	PEG	B	903	7/7	0.92	0.12	42,45,48,49	0
3	PEG	A	602	7/7	0.93	0.14	41,44,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around ATP B 902:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around ATP A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around P4C B 906:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.