



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 09:24 AM EDT

PDB ID : 7TYG  
Title : Structure of the human leucine rich repeat protein SHOC2, residues 80-582  
Authors : Dhembi, A.; Clark, K.; King, D.A.  
Deposited on : 2022-02-13  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	 3% 90% 7% ..
1	B	505	 2% 89% 8% ..

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8414 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leucine-rich repeat protein SHOC-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	494	3897	2467	671	742	17	0	0	0
1	B	493	3891	2465	669	739	18	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLY	-	expression tag	UNP Q9UQ13
A	79	PRO	-	expression tag	UNP Q9UQ13
B	78	GLY	-	expression tag	UNP Q9UQ13
B	79	PRO	-	expression tag	UNP Q9UQ13

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	280	Total	O	0	0
			280	280		
3	B	345	Total	O	0	0
			345	345		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leucine-rich repeat protein SHOC-2



- Molecule 1: Leucine-rich repeat protein SHOC-2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.35Å 102.88Å 120.83Å 90.00° 101.74° 90.00°	Depositor
Resolution (Å)	41.44 – 1.90 41.44 – 1.90	Depositor EDS
% Data completeness (in resolution range)	94.7 (41.44-1.90) 94.7 (41.44-1.90)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.188 , 0.226 0.186 , 0.225	Depositor DCC
$R_{free}$ test set	4119 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.3	Xtrriage
Anisotropy	0.184	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8414	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/3954	0.58	0/5359
1	B	0.37	0/3948	0.61	0/5350
All	All	0.36	0/7902	0.60	0/10709

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3897	0	4059	25	0
1	B	3891	0	4057	21	0
2	B	1	0	0	0	0
3	A	280	0	0	2	1
3	B	345	0	0	2	1
All	All	8414	0	8116	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:ASP:OD1	1:B:190:ARG:NH1	2.17	0.76
1:A:371:PRO:HD2	1:A:374:ILE:HG12	1.78	0.65
1:A:327:LEU:HB2	1:A:353:GLN:HB2	1.82	0.62
1:B:327:LEU:HB2	1:B:353:GLN:HB2	1.81	0.61
1:B:371:PRO:HD2	1:B:374:ILE:HG12	1.83	0.60
1:A:228:LYS:NZ	1:A:249:GLN:OE1	2.33	0.59
1:B:371:PRO:O	1:B:374:ILE:HG13	2.03	0.58
1:B:486:LEU:HD13	1:B:490:ILE:HD13	1.87	0.57
1:B:209:LYS:NZ	3:B:703:HOH:O	2.28	0.56
1:B:89:GLU:HG3	1:B:91:ILE:H	1.70	0.56
1:A:350:GLY:O	1:A:353:GLN:HG2	2.06	0.56
1:A:182:ARG:NH2	3:A:610:HOH:O	2.40	0.55
1:B:200:ARG:HB2	1:B:223:ARG:HG3	1.88	0.55
1:A:488:ARG:NE	1:A:512:GLU:OE2	2.41	0.53
1:A:371:PRO:O	1:A:374:ILE:HG13	2.08	0.53
1:B:549:GLU:H	1:B:549:GLU:CD	2.11	0.53
1:B:119:ILE:HA	1:B:122:LEU:HD13	1.90	0.52
1:A:120:LYS:HB3	1:A:141:GLU:HB3	1.93	0.51
1:B:522:LEU:O	1:B:546:MET:HA	2.13	0.49
1:A:511:GLU:O	1:A:536:GLU:HG3	2.13	0.49
1:B:200:ARG:HA	1:B:223:ARG:O	2.13	0.48
1:A:147:ASN:HA	1:A:170:LYS:HE3	1.94	0.48
1:A:155:GLU:OE1	1:B:366:ARG:HD2	2.13	0.48
1:A:146:VAL:HG13	1:A:167:ASN:O	2.15	0.47
1:B:120:LYS:HB3	1:B:141:GLU:HB3	1.96	0.47
1:A:522:LEU:O	1:A:546:MET:HA	2.15	0.47
1:A:226:LYS:NZ	1:B:272:GLU:HG2	2.30	0.46
1:B:222:ILE:CG1	1:B:245:VAL:HG12	2.46	0.46
1:A:87:ASN:N	1:A:87:ASN:OD1	2.47	0.45
1:B:253:LEU:HD11	1:B:268:LEU:HD11	1.98	0.45
1:A:486:LEU:HD13	1:A:490:ILE:HD13	1.99	0.45
1:A:510:PRO:HB2	1:A:512:GLU:HG2	1.99	0.45
1:A:173:MET:O	1:A:173:MET:HG3	2.16	0.45
1:A:579:ARG:HH11	1:A:579:ARG:HB2	1.82	0.44
1:B:409:LEU:HB2	1:B:432:LEU:HD23	2.00	0.44
1:A:327:LEU:CB	1:A:353:GLN:HB2	2.46	0.44
1:B:579:ARG:H	1:B:579:ARG:HG2	1.38	0.44
1:A:222:ILE:CG1	1:A:245:VAL:HG12	2.48	0.43
1:A:327:LEU:HB2	1:A:353:GLN:CB	2.49	0.42
1:A:134:LYS:HB3	1:A:134:LYS:HE3	1.79	0.42
1:B:358:TYR:CD2	1:B:380:VAL:HG13	2.55	0.42
1:B:559:GLN:HB3	3:B:996:HOH:O	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:482:GLN:NE2	3:A:601:HOH:O	2.19	0.41
1:A:571:PHE:CE1	1:A:575:GLN:HG3	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:603:HOH:O	3:B:746:HOH:O[2_455]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/505 (97%)	460 (94%)	32 (6%)	0	100	100
1	B	491/505 (97%)	461 (94%)	29 (6%)	1 (0%)	47	38
All	All	983/1010 (97%)	921 (94%)	61 (6%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	166	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	460/468 (98%)	446 (97%)	14 (3%)	41	33
1	B	459/468 (98%)	445 (97%)	14 (3%)	40	32
All	All	919/936 (98%)	891 (97%)	28 (3%)	41	33

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	173	MET
1	A	223	ARG
1	A	369	LYS
1	A	374	ILE
1	A	420	ASP
1	A	436	LEU
1	A	512	GLU
1	A	513	ILE
1	A	533	LEU
1	A	536	GLU
1	A	561	VAL
1	A	570	GLN
1	A	579	ARG
1	B	98	ARG
1	B	100	GLU
1	B	101	ASN
1	B	121	GLU
1	B	163	ASP
1	B	166	ASP
1	B	201	PHE
1	B	206	THR
1	B	261	THR
1	B	272	GLU
1	B	274	LEU
1	B	374	ILE
1	B	539	LEU
1	B	579	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	494/505 (97%)	0.04	16 (3%) 47 50	26, 40, 69, 101	0
1	B	493/505 (97%)	-0.06	9 (1%) 68 71	24, 35, 57, 83	0
All	All	987/1010 (97%)	-0.01	25 (2%) 57 60	24, 38, 63, 101	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	562	ALA	5.3
1	A	561	VAL	5.1
1	A	567	PHE	4.5
1	A	560	ILE	4.5
1	A	571	PHE	4.0
1	A	88	ALA	3.8
1	A	559	GLN	3.3
1	B	90	VAL	3.0
1	A	579	ARG	3.0
1	B	92	LYS	2.9
1	B	97	CYS	2.9
1	A	576	GLY	2.8
1	B	113	HIS	2.7
1	A	556	LEU	2.7
1	B	348	VAL	2.6
1	A	535	PHE	2.6
1	A	575	GLN	2.6
1	B	227	ILE	2.4
1	B	102	SER	2.4
1	A	549	GLU	2.4
1	A	557	PRO	2.4
1	A	558	PRO	2.4
1	B	98	ARG	2.2
1	B	204	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	574	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	601	1/1	0.97	0.12	30,30,30,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.