



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 15, 2022 – 01:27 am BST

PDB ID : 7ZMF
Title : Dehydratase domain of module 3 from Brevibacillus Brevis orphan BGC11
Authors : Tittes, Y.U.; Herbst, D.A.; Jakob, R.P.; Maier, T.
Deposited on : 2022-04-19
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.30
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.30


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	
1	B	304	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	B	1501	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8604 atoms, of which 4238 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative polyketide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	273	4217	1344	2096	359	405	13	0	0	0
1	B	279	4293	1368	2134	365	412	14	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1133	GLY	-	expression tag	UNP C0ZGQ6
A	1134	PRO	-	expression tag	UNP C0ZGQ6
A	1436	ALA	-	expression tag	UNP C0ZGQ6
B	1133	GLY	-	expression tag	UNP C0ZGQ6
B	1134	PRO	-	expression tag	UNP C0ZGQ6
B	1436	ALA	-	expression tag	UNP C0ZGQ6

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			14	3	8	3		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	43	Total	O	0	0
			43	43		
4	B	35	Total	O	0	0
			35	35		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative polyketide synthase



- Molecule 1: Putative polyketide synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	72.38Å 197.36Å 39.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.68 – 2.21 48.68 – 2.21	Depositor EDS
% Data completeness (in resolution range)	98.9 (48.68-2.21) 98.9 (48.68-2.21)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.20rc4_4425	Depositor
R, R_{free}	0.224 , 0.246 0.222 , 0.245	Depositor DCC
R_{free} test set	1999 reflections (6.90%)	wwPDB-VP
Wilson B-factor (Å ²)	44.4	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8604	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2160	0.51	0/2918
1	B	0.33	0/2200	0.50	0/2972
All	All	0.32	0/4360	0.50	0/5890

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2121	2096	2096	9	0
1	B	2159	2134	2134	6	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	6	8	8	0	0
4	A	43	0	0	1	0
4	B	35	0	0	0	0
All	All	4366	4238	4238	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1276:SER:OG	1:B:1415:ASN:OD1	2.09	0.71
1:A:1323:GLN:OE1	1:A:1391:ARG:NH1	2.24	0.70
1:A:1272:PRO:O	4:A:1601:HOH:O	2.12	0.68
1:A:1366:SER:N	1:A:1430:ASN:OD1	2.28	0.67
1:A:1276:SER:OG	1:A:1415:ASN:OD1	2.14	0.65
1:A:1155:GLU:OE1	1:A:1157:ARG:NH2	2.30	0.64
1:A:1240:GLU:HG2	1:A:1267:VAL:HG12	1.98	0.45
1:B:1369:LEU:HD12	1:B:1428:ILE:HD13	2.00	0.43
1:B:1297:TYR:CZ	1:B:1312:ARG:HA	2.54	0.43
1:A:1404:LYS:HA	1:A:1423:PHE:O	2.19	0.42
1:B:1358:LEU:HD11	1:B:1390:ILE:HG22	2.02	0.41
1:A:1297:TYR:CZ	1:A:1312:ARG:HA	2.56	0.41
1:B:1404:LYS:HA	1:B:1423:PHE:O	2.21	0.41
1:B:1370:PRO:HA	1:B:1425:THR:HG22	2.01	0.41
1:A:1206:LEU:N	1:A:1206:LEU:HD12	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/304 (87%)	258 (97%)	7 (3%)	0	100	100
1	B	273/304 (90%)	266 (97%)	7 (3%)	0	100	100
All	All	538/608 (88%)	524 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/259 (91%)	234 (100%)	1 (0%)	91	95
1	B	238/259 (92%)	237 (100%)	1 (0%)	91	95
All	All	473/518 (91%)	471 (100%)	2 (0%)	91	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1252	ASP
1	B	1218	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	1502	-	5,5,5	0.64	0	5,5,5	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1502	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/304 (89%)	0.97	37 (13%) 3 2	34, 52, 103, 136	0
1	B	279/304 (91%)	1.03	33 (11%) 4 3	36, 59, 96, 152	0
All	All	552/608 (90%)	1.00	70 (12%) 3 3	34, 54, 97, 152	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1250	LYS	7.3
1	B	1140	GLY	7.1
1	A	1428	ILE	6.7
1	A	1402	VAL	6.2
1	B	1237	ALA	6.2
1	A	1425	THR	5.5
1	B	1364	ASP	5.1
1	B	1365	PHE	5.0
1	A	1255	GLN	4.8
1	A	1430	ASN	4.7
1	B	1363	SER	4.4
1	A	1429	ALA	4.3
1	A	1431	VAL	4.2
1	B	1298	ASP	3.9
1	B	1284	GLU	3.7
1	A	1251	GLN	3.6
1	B	1286	CYS	3.6
1	A	1199	GLY	3.6
1	A	1252	ASP	3.5
1	A	1206	LEU	3.5
1	B	1251	GLN	3.4
1	B	1154	MET	3.4
1	A	1296	CYS	3.3
1	B	1311	HIS	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1423	PHE	3.1
1	A	1432	SER	3.0
1	A	1366	SER	3.0
1	B	1358	LEU	2.9
1	B	1394	GLU	2.9
1	B	1360	MET	2.9
1	A	1358	LEU	2.8
1	B	1383	SER	2.7
1	B	1396	SER	2.7
1	A	1359	MET	2.7
1	A	1142	ILE	2.7
1	B	1398	ALA	2.6
1	B	1306	ASP	2.6
1	A	1372	ALA	2.6
1	A	1368	ILE	2.6
1	B	1393	ARG	2.6
1	B	1282	LEU	2.5
1	B	1299	ALA	2.5
1	A	1163	THR	2.4
1	B	1232	GLU	2.4
1	A	1198	GLY	2.3
1	A	1140	GLY	2.3
1	A	1284	GLU	2.3
1	B	1278	ASP	2.3
1	A	1228	GLN	2.3
1	A	1403	GLU	2.3
1	B	1141	VAL	2.2
1	A	1357	GLY	2.2
1	B	1325	LEU	2.2
1	B	1412	GLU	2.2
1	A	1216	TRP	2.2
1	A	1338	GLN	2.2
1	B	1269	CYS	2.2
1	A	1392	TYR	2.1
1	B	1381	ASP	2.1
1	A	1321	GLN	2.1
1	A	1256	THR	2.1
1	A	1286	CYS	2.1
1	A	1302	LYS	2.1
1	B	1371	PHE	2.1
1	B	1431	VAL	2.1
1	A	1422	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1404	LYS	2.1
1	B	1297	TYR	2.1
1	A	1250	LYS	2.0
1	B	1283	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	B	1501	1/1	-0.35	0.47	69,69,69,69	0
2	MG	A	1501	1/1	0.45	0.32	111,111,111,111	0
3	GOL	B	1502	6/6	0.73	0.17	55,66,76,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.