



# Full wwPDB X-ray Structure Validation Report i

Jun 27, 2022 – 07:09 pm BST

PDB ID : 7ZV3  
Title : Crystal structure of indoleamine 2,3-dioxygenase 1 (IDO1) in complex with ferric heme and MMG-0472  
Authors : Roehrig, U.F.; Reynaud, A.; Pojer, F.; Michelin, O.; Zoete, V.  
Deposited on : 2022-05-13  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1279 (2.58-2.54)
Clashscore	141614	1327 (2.58-2.54)
Ramachandran outliers	138981	1312 (2.58-2.54)
Sidechain outliers	138945	1312 (2.58-2.54)
RSRZ outliers	127900	1269 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Indoleamine 2,3-dioxygenase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C 2964	N 1905	O 505	S 537	17	0	0
1	B	372	Total	C 2951	N 1897	O 504	S 533	17	0	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP P14902
A	-18	GLY	-	expression tag	UNP P14902
A	-17	SER	-	expression tag	UNP P14902
A	-16	SER	-	expression tag	UNP P14902
A	-15	HIS	-	expression tag	UNP P14902
A	-14	HIS	-	expression tag	UNP P14902
A	-13	HIS	-	expression tag	UNP P14902
A	-12	HIS	-	expression tag	UNP P14902
A	-11	HIS	-	expression tag	UNP P14902
A	-10	HIS	-	expression tag	UNP P14902
A	-9	SER	-	expression tag	UNP P14902
A	-8	SER	-	expression tag	UNP P14902
A	-7	GLY	-	expression tag	UNP P14902
A	-6	LEU	-	expression tag	UNP P14902
A	-5	VAL	-	expression tag	UNP P14902
A	-4	PRO	-	expression tag	UNP P14902
A	-3	ARG	-	expression tag	UNP P14902
A	-2	GLY	-	expression tag	UNP P14902
A	-1	SER	-	expression tag	UNP P14902
A	0	HIS	-	expression tag	UNP P14902
A	3	SER	HIS	conflict	UNP P14902
B	-19	MET	-	initiating methionine	UNP P14902
B	-18	GLY	-	expression tag	UNP P14902
B	-17	SER	-	expression tag	UNP P14902
B	-16	SER	-	expression tag	UNP P14902

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP P14902
B	-14	HIS	-	expression tag	UNP P14902
B	-13	HIS	-	expression tag	UNP P14902
B	-12	HIS	-	expression tag	UNP P14902
B	-11	HIS	-	expression tag	UNP P14902
B	-10	HIS	-	expression tag	UNP P14902
B	-9	SER	-	expression tag	UNP P14902
B	-8	SER	-	expression tag	UNP P14902
B	-7	GLY	-	expression tag	UNP P14902
B	-6	LEU	-	expression tag	UNP P14902
B	-5	VAL	-	expression tag	UNP P14902
B	-4	PRO	-	expression tag	UNP P14902
B	-3	ARG	-	expression tag	UNP P14902
B	-2	GLY	-	expression tag	UNP P14902
B	-1	SER	-	expression tag	UNP P14902
B	0	HIS	-	expression tag	UNP P14902
B	3	SER	HIS	conflict	UNP P14902

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 4-(3-chloro-2-phenethylphenyl)-1H-1,2,3-triazole (three-letter code: RCT) (formula: C<sub>16</sub>H<sub>14</sub>ClN<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	Cl	N		
3	A	1	20	16	1	3	0	0
3	B	1	20	16	1	3	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	26	Total O 26 26	0	0
6	B	26	Total O 26 26	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Indoleamine 2,3-dioxygenase 1



- Molecule 1: Indoleamine 2,3-dioxygenase 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.65 Å    97.42 Å    132.00 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	48.71 – 2.55 48.71 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.71-2.55) 95.0 (48.71-2.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	0.85 (at 2.54 Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
$R$ , $R_{free}$	0.191 , 0.241 0.191 , 0.240	Depositor DCC
$R_{free}$ test set	1819 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.8	Xtriage
Anisotropy	0.370	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6109	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4, RCT, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.46	0/3032	0.59	0/4098
1	B	0.45	0/3019	0.59	0/4082
All	All	0.46	0/6051	0.59	0/8180

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2964	0	2978	37	0
1	B	2951	0	2966	22	0
2	A	43	0	30	4	0
2	B	43	0	30	3	0
3	A	20	0	0	1	0
3	B	20	0	0	1	0
4	A	6	0	8	2	0
5	A	5	0	0	0	0
5	B	5	0	0	1	0
6	A	26	0	0	0	0
6	B	26	0	0	1	0
All	All	6109	0	6012	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:ILE:HG13	1:A:357:PRO:HD3	1.62	0.82
1:A:58:ARG:HH22	1:A:61:LYS:HD3	1.52	0.74
1:A:337:LYS:HG3	1:A:399:LEU:HD21	1.79	0.64
1:A:21:VAL:HG13	1:A:24:ALA:HB3	1.80	0.62
1:A:175:ALA:HA	1:A:178:ILE:HD12	1.81	0.62
1:A:99:VAL:HG11	1:A:243:LEU:HD11	1.82	0.61
2:A:501:HEM:HBB2	2:A:501:HEM:HMB2	1.83	0.60
1:A:12:SER:OG	1:A:14:GLU:OE2	2.20	0.60
1:A:227:PHE:O	1:A:231:ARG:NH1	2.35	0.60
1:B:227:PHE:O	1:B:231:ARG:NH1	2.35	0.59
2:B:501:HEM:HMB2	2:B:501:HEM:HBB2	1.83	0.59
1:A:109:VAL:O	1:A:113:GLN:HG3	2.04	0.58
1:A:253:TRP:CG	1:A:257:LYS:HE2	2.40	0.57
1:A:321:LEU:HD21	1:A:400:LEU:HD22	1.84	0.57
2:B:501:HEM:NA	3:B:502:RCT:N12	2.52	0.57
1:B:32:LEU:HD12	1:B:40:MET:HG2	1.87	0.55
1:B:175:ALA:HA	1:B:178:ILE:HD12	1.89	0.55
1:A:76:GLN:HB3	1:A:114:LEU:HD11	1.88	0.54
1:B:385:MET:HA	1:B:385:MET:HE2	1.90	0.53
1:A:383:ASP:OD2	1:A:383:ASP:N	2.42	0.53
1:A:72:ASP:O	1:A:76:GLN:HG3	2.09	0.52
1:B:76:GLN:HB3	1:B:114:LEU:HD11	1.91	0.52
1:B:42:ILE:HD12	1:B:62:LEU:HD11	1.93	0.51
1:A:287:HIS:H	4:A:503:GOL:H2	1.75	0.51
1:B:280:GLN:HB3	1:B:283:ALA:HB2	1.91	0.51
1:B:280:GLN:HB3	1:B:283:ALA:CB	2.41	0.51
1:A:58:ARG:NH2	1:A:61:LYS:HD3	2.25	0.50
1:A:267:SER:OG	1:A:269:VAL:HG22	2.11	0.50
1:B:384:LEU:HD11	2:B:501:HEM:HMA2	1.95	0.48
1:B:286:GLY:HA3	5:B:503:SO4:O4	2.12	0.48
1:A:217:ILE:HD11	2:A:501:HEM:HAB	1.95	0.48
1:B:105:ARG:HA	1:B:108:ALA:HB3	1.96	0.48
1:A:280:GLN:HB3	1:A:283:ALA:HB3	1.96	0.47
1:A:97:GLY:O	1:A:99:VAL:HG23	2.15	0.47
1:B:80:ARG:NH2	1:B:128:ASP:OD2	2.39	0.47
1:A:177:ALA:HB2	1:A:206:CYS:HB2	1.96	0.46
1:B:178:ILE:HD11	1:B:269:VAL:HG23	1.97	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:VAL:HG13	6:B:610:HOH:O	2.16	0.46
1:A:287:HIS:N	4:A:503:GOL:H2	2.31	0.45
1:A:238:LYS:HG3	1:A:258:GLU:HB3	1.98	0.45
1:A:384:LEU:HD12	1:A:385:MET:N	2.33	0.44
1:B:90:TYR:CE2	1:B:104:PRO:HD3	2.53	0.44
1:A:100:ARG:HD3	1:A:100:ARG:HA	1.75	0.43
1:A:253:TRP:CE3	1:A:257:LYS:HG3	2.54	0.43
1:B:181:ILE:HG21	1:B:310:LEU:HD21	2.01	0.43
1:A:348:GLN:O	1:A:351:THR:HB	2.19	0.43
1:B:57:GLU:O	1:B:61:LYS:HG2	2.18	0.42
1:B:58:ARG:HE	1:B:58:ARG:HB3	1.60	0.42
1:A:279:ILE:HD13	1:A:395:THR:HG23	2.01	0.42
1:B:188:MET:SD	1:B:316:VAL:HG22	2.60	0.42
2:A:501:HEM:NA	3:A:502:RCT:N12	2.66	0.42
1:A:27:ASN:O	1:A:74:LYS:NZ	2.51	0.42
1:A:248:VAL:HG22	1:A:256:PRO:HB3	2.02	0.41
1:A:214:PHE:HD1	1:A:214:PHE:O	2.03	0.41
1:B:279:ILE:HD13	1:B:395:THR:HG23	2.03	0.41
1:A:230:LEU:HD12	1:A:230:LEU:HA	1.90	0.41
1:A:269:VAL:O	1:A:272:CYS:HB3	2.20	0.41
1:A:328:LEU:HD23	1:A:328:LEU:HA	1.89	0.41
1:B:384:LEU:HD12	1:B:384:LEU:HA	1.78	0.41
1:A:345:TYR:O	1:A:348:GLN:HB2	2.21	0.40
1:A:217:ILE:CD1	2:A:501:HEM:HAB	2.52	0.40
1:A:339:LEU:HD23	1:A:339:LEU:HA	1.92	0.40
1:B:339:LEU:HD23	1:B:339:LEU:HA	1.88	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/423 (88%)	350 (95%)	20 (5%)	0	100	100
1	B	368/423 (87%)	354 (96%)	14 (4%)	0	100	100
All	All	738/846 (87%)	704 (95%)	34 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	323/365 (88%)	309 (96%)	14 (4%)	29	39
1	B	322/365 (88%)	309 (96%)	13 (4%)	31	42
All	All	645/730 (88%)	618 (96%)	27 (4%)	30	40

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ILE
1	A	13	LYS
1	A	14	GLU
1	A	21	VAL
1	A	58	ARG
1	A	80	ARG
1	A	125	VAL
1	A	176	SER
1	A	214	PHE
1	A	312	SER
1	A	317	ARG
1	A	349	ILE
1	A	356	ILE
1	A	401	LYS
1	B	13	LYS
1	B	55	LEU
1	B	58	ARG
1	B	125	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	135	LYS
1	B	149	ASP
1	B	176	SER
1	B	183	THR
1	B	214	PHE
1	B	242	GLN
1	B	269	VAL
1	B	311	GLU
1	B	398	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	503	-	5,5,5	0.70	0	5,5,5	1.12	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	RCT	B	502	2	22,22,22	0.99	1 (4%)	25,29,29	1.63	4 (16%)
2	HEM	B	501	3,1	41,50,50	1.57	5 (12%)	45,82,82	1.85	13 (28%)
5	SO4	B	503	-	4,4,4	0.14	0	6,6,6	0.19	0
3	RCT	A	502	2	22,22,22	0.93	2 (9%)	25,29,29	1.49	3 (12%)
2	HEM	A	501	3,1	41,50,50	1.63	7 (17%)	45,82,82	1.66	11 (24%)
5	SO4	A	504	-	4,4,4	0.11	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	503	-	-	4/4/4/4	-
3	RCT	B	502	2	-	2/8/9/9	0/3/3/3
2	HEM	B	501	3,1	-	4/12/54/54	-
3	RCT	A	502	2	-	2/8/9/9	0/3/3/3
2	HEM	A	501	3,1	-	4/12/54/54	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	FE-ND	4.73	2.20	1.96
2	B	501	HEM	C3C-C2C	-4.24	1.34	1.40
2	B	501	HEM	FE-ND	3.81	2.15	1.96
2	A	501	HEM	C3C-CAC	3.69	1.55	1.47
2	A	501	HEM	C3C-C2C	-3.55	1.35	1.40
2	B	501	HEM	C3C-CAC	3.48	1.54	1.47
2	A	501	HEM	CAA-C2A	2.90	1.56	1.52
2	B	501	HEM	CAB-C3B	2.81	1.55	1.47
2	A	501	HEM	CAB-C3B	2.79	1.55	1.47
3	B	502	RCT	C8-CL2	2.39	1.79	1.73
2	B	501	HEM	CMD-C2D	2.26	1.55	1.50
3	A	502	RCT	C8-CL2	2.08	1.78	1.73
2	A	501	HEM	CMD-C2D	2.05	1.55	1.50
2	A	501	HEM	CHA-C4D	2.05	1.40	1.35
3	A	502	RCT	C9-N13	-2.01	1.31	1.36

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	RCT	C1-C2-C3	-5.29	102.73	112.49
3	A	502	RCT	N13-N12-N11	-4.74	105.34	111.24
3	B	502	RCT	N13-N12-N11	-4.41	105.76	111.24
2	B	501	HEM	CAD-CBD-CCD	-4.28	104.38	113.60
2	B	501	HEM	C4C-CHD-C1D	4.21	128.11	122.56
2	A	501	HEM	C1B-NB-C4B	4.08	109.29	105.07
2	B	501	HEM	C4D-ND-C1D	4.01	109.22	105.07
3	A	502	RCT	C1-C2-C3	-3.87	105.35	112.49
2	A	501	HEM	C4D-ND-C1D	3.56	108.75	105.07
2	B	501	HEM	C1B-NB-C4B	3.20	108.38	105.07
2	B	501	HEM	C4A-C3A-C2A	3.19	109.22	107.00
3	B	502	RCT	C4-C9-N13	2.79	125.37	120.93
2	B	501	HEM	C3D-C4D-ND	-2.74	107.12	110.17
3	A	502	RCT	C4-C9-N13	2.70	125.24	120.93
2	A	501	HEM	C3D-C4D-ND	-2.67	107.20	110.17
2	A	501	HEM	CAA-CBA-CGA	-2.59	106.49	113.76
2	A	501	HEM	C4C-CHD-C1D	2.57	125.96	122.56
2	B	501	HEM	C2D-C1D-ND	-2.53	106.86	109.88
2	B	501	HEM	CMA-C3A-C4A	-2.47	124.67	128.46
2	A	501	HEM	C4B-CHC-C1C	2.45	125.79	122.56
3	B	502	RCT	C2-C1-C14	2.40	121.70	113.28
2	A	501	HEM	CMA-C3A-C4A	-2.36	124.83	128.46
2	A	501	HEM	C3B-C2B-C1B	2.34	108.22	106.49
2	A	501	HEM	C2B-C1B-NB	-2.33	107.08	109.84
2	B	501	HEM	C3C-C4C-NC	-2.28	106.64	110.94
2	B	501	HEM	CBA-CAA-C2A	-2.26	108.77	112.62
2	B	501	HEM	C3B-C2B-C1B	2.25	108.16	106.49
2	A	501	HEM	C2D-C1D-ND	-2.14	107.31	109.88
2	B	501	HEM	CMB-C2B-C1B	-2.14	121.78	125.04
2	B	501	HEM	CHD-C1D-C2D	2.12	128.30	124.98
2	A	501	HEM	C4A-C3A-C2A	2.05	108.42	107.00

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	RCT	C3-C4-C9-C10
3	B	502	RCT	C3-C4-C9-C10
4	A	503	GOL	O1-C1-C2-C3
4	A	503	GOL	C1-C2-C3-O3
4	A	503	GOL	O1-C1-C2-O2
4	A	503	GOL	O2-C2-C3-O3
3	B	502	RCT	C5-C4-C9-N13

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	501	HEM	CAA-CBA-CGA-O1A
2	A	501	HEM	CAD-CBD-CGD-O1D
2	A	501	HEM	CAD-CBD-CGD-O2D
2	B	501	HEM	CAD-CBD-CGD-O1D
2	B	501	HEM	CAA-CBA-CGA-O1A
2	B	501	HEM	CAA-CBA-CGA-O2A
2	A	501	HEM	CAA-CBA-CGA-O2A
3	A	502	RCT	C5-C4-C9-N13
2	B	501	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.


6 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	GOL	2	0
3	B	502	RCT	1	0
2	B	501	HEM	3	0
5	B	503	SO4	1	0
3	A	502	RCT	1	0
2	A	501	HEM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	374/423 (88%)	0.07	5 (1%) 77 82	41, 59, 81, 112	0
1	B	372/423 (87%)	0.10	6 (1%) 72 78	43, 58, 79, 95	0
All	All	746/846 (88%)	0.08	11 (1%) 73 80	41, 59, 80, 112	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	382	THR	4.8
1	B	67	ILE	4.7
1	A	380	GLY	3.2
1	B	70	LEU	3.2
1	B	69	HIS	3.0
1	A	355	LEU	2.6
1	B	32	LEU	2.4
1	B	35	PHE	2.3
1	B	33	PRO	2.3
1	A	354	ILE	2.2
1	A	241	PRO	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i


There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	A	504	5/5	0.79	0.29	81,82,86,89	5
4	GOL	A	503	6/6	0.87	0.19	78,79,82,84	0
2	HEM	A	501	43/43	0.94	0.18	55,62,69,76	43
3	RCT	B	502	20/20	0.95	0.16	46,57,62,64	0
3	RCT	A	502	20/20	0.96	0.14	55,65,68,68	0
2	HEM	B	501	43/43	0.96	0.21	46,52,59,62	43
5	SO4	B	503	5/5	0.96	0.22	62,65,69,71	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.