



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2022 – 10:24 am BST

PDB ID : 8AD7
Title : Flavin-dependent tryptophan 6-halogenase Thal in complex with D-Trp
Authors : Moritzer, A.C.; Niemann, H.H.
Deposited on : 2022-07-08
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	 2% 91% 7% .
1	B	534	 9% 89% 9% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8691 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan 6-halogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	528	4216	2693	723	780	20	0	0	0
1	B	525	4185	2671	720	774	20	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP A1E280
A	-1	ALA	-	expression tag	UNP A1E280
A	0	MET	-	expression tag	UNP A1E280
A	1	GLY	-	expression tag	UNP A1E280
B	-2	GLY	-	expression tag	UNP A1E280
B	-1	ALA	-	expression tag	UNP A1E280
B	0	MET	-	expression tag	UNP A1E280
B	1	GLY	-	expression tag	UNP A1E280

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is D-TRYPTOPHAN (three-letter code: DTR) (formula: $C_{11}H_{12}N_2O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			15	11	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	154	Total	O	0	0
			154	154		
5	B	70	Total	O	0	0
			70	70		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tryptophan 6-halogenase



- Molecule 1: Tryptophan 6-halogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	138.91Å 138.91Å 142.59Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.80 – 2.33 49.75 – 2.33	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.80-2.33) 100.0 (49.75-2.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.192 , 0.237 0.196 , 0.235	Depositor DCC
R_{free} test set	3342 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	63.9	Xtrriage
Anisotropy	0.105	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.036 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8691	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL, DTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/4331	0.83	0/5880
1	B	0.68	0/4297	0.82	0/5831
All	All	0.68	0/8628	0.82	0/11711

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4216	0	4071	22	0
1	B	4185	0	4048	23	0
2	A	15	0	0	2	0
3	A	24	0	32	2	0
3	B	12	0	16	0	0
4	A	15	0	11	0	0
5	A	154	0	0	0	0
5	B	70	0	0	1	0
All	All	8691	0	8178	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54:ASN:ND2	1:B:366:TYR:OH	2.17	0.75
1:A:337:ARG:NH1	2:A:602:PO4:O4	2.31	0.64
1:B:130:LYS:NZ	5:B:701:HOH:O	2.27	0.63
1:A:248:SER:HA	1:A:251:LEU:O	2.00	0.60
1:B:248:SER:HA	1:B:251:LEU:O	2.02	0.59
1:B:301:ASP:HA	1:B:304:ALA:O	2.04	0.58
1:A:301:ASP:HA	1:A:304:ALA:O	2.04	0.58
1:A:58:ALA:HB1	1:A:366:TYR:CZ	2.41	0.56
1:A:250:HIS:HE2	2:A:603:PO4:P	2.32	0.53
1:A:20:THR:HG21	1:A:226:VAL:HG11	1.93	0.51
1:B:105:THR:HG22	1:B:107:THR:HG23	1.92	0.50
1:A:168:TYR:H	3:A:604:GOL:H12	1.77	0.50
1:B:445:ASN:HB3	1:B:464:ASN:OD1	2.11	0.49
1:B:80:MET:CE	1:B:166:THR:HG21	2.43	0.49
1:A:231:PHE:CZ	1:A:331:PHE:HB3	2.48	0.48
1:A:80:MET:CE	1:A:166:THR:HG21	2.45	0.47
1:B:20:THR:HG21	1:B:226:VAL:HG11	1.97	0.46
1:B:306:GLN:O	1:B:310:THR:OG1	2.28	0.46
1:A:4:ARG:HD3	1:B:490:ALA:HB1	1.97	0.46
1:A:431:ILE:O	1:A:435:VAL:HG23	2.16	0.45
1:B:18:TRP:CE3	1:B:182:ARG:HA	2.50	0.45
1:A:100:LEU:HD12	1:A:104:HIS:HB2	1.98	0.45
1:A:18:TRP:CE3	1:A:182:ARG:HA	2.52	0.44
1:A:306:GLN:O	1:A:310:THR:OG1	2.28	0.44
1:B:185:ALA:HA	1:B:189:GLN:HG3	1.99	0.44
1:A:138:GLU:OE2	1:A:507:LYS:NZ	2.51	0.44
1:B:231:PHE:CZ	1:B:331:PHE:HB3	2.53	0.44
1:A:428:ALA:O	1:A:431:ILE:HG22	2.18	0.43
1:B:405:GLN:NE2	1:B:426:LYS:O	2.48	0.43
1:A:237:ASN:O	1:A:241:GLU:HA	2.18	0.43
1:A:343:ASN:HA	1:A:381:PHE:CE2	2.54	0.43
1:B:23:TYR:CG	1:B:369:ILE:CG2	3.02	0.43
1:B:431:ILE:O	1:B:435:VAL:HG23	2.19	0.42
1:A:439:ARG:HE	1:A:483:ARG:HG3	1.85	0.42
1:B:282:GLY:HA2	1:B:418:PHE:CE2	2.56	0.41
1:A:23:TYR:CG	1:A:369:ILE:CG2	3.04	0.41
1:B:51:THR:HG21	1:B:55:LEU:HD22	2.01	0.41
1:B:121:ILE:HA	1:B:122:PRO:HD3	1.95	0.41
1:B:339:ALA:O	1:B:346:SER:OG	2.30	0.41
1:B:439:ARG:HA	1:B:482:PRO:HA	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:454:TYR:CE2	1:A:461:GLU:HG3	2.57	0.40
1:A:349:LEU:HD13	3:A:607:GOL:H31	2.03	0.40
1:B:237:ASN:O	1:B:241:GLU:HA	2.21	0.40
1:B:428:ALA:O	1:B:431:ILE:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	526/534 (98%)	504 (96%)	22 (4%)	0	100	100
1	B	521/534 (98%)	499 (96%)	21 (4%)	1 (0%)	47	55
All	All	1047/1068 (98%)	1003 (96%)	43 (4%)	1 (0%)	51	62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	46	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/445 (100%)	439 (99%)	4 (1%)	78	87

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	440/445 (99%)	434 (99%)	6 (1%)	67	78
All	All	883/890 (99%)	873 (99%)	10 (1%)	73	83

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ILE
1	A	147	ASP
1	A	439	ARG
1	A	484	ASN
1	B	54	ASN
1	B	147	ASP
1	B	268	ASN
1	B	292	ARG
1	B	307	ASP
1	B	484	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	133	GLN
1	A	327	ASN
1	A	464	ASN
1	B	30	ASN
1	B	54	ASN
1	B	133	GLN
1	B	189	GLN
1	B	268	ASN
1	B	327	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	602	-	5,5,5	0.14	0	5,5,5	0.33	0
3	GOL	A	604	-	5,5,5	0.16	0	5,5,5	0.53	0
2	PO4	A	602	-	4,4,4	0.49	0	6,6,6	0.63	0
3	GOL	A	605	-	5,5,5	0.13	0	5,5,5	0.36	0
3	GOL	A	606	-	5,5,5	0.34	0	5,5,5	0.52	0
3	GOL	A	607	-	5,5,5	0.39	0	5,5,5	0.74	0
3	GOL	B	601	-	5,5,5	0.16	0	5,5,5	0.40	0
2	PO4	A	603	-	4,4,4	0.54	0	6,6,6	0.54	0
2	PO4	A	601	-	4,4,4	1.26	0	6,6,6	0.67	0
4	DTR	A	608	-	14,16,16	0.73	0	16,22,22	1.00	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	602	-	-	1/4/4/4	-
3	GOL	A	604	-	-	2/4/4/4	-
3	GOL	A	606	-	-	2/4/4/4	-
3	GOL	A	605	-	-	3/4/4/4	-
3	GOL	A	607	-	-	2/4/4/4	-
3	GOL	B	601	-	-	0/4/4/4	-
4	DTR	A	608	-	-	5/7/8/8	0/2/2/2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	608	DTR	OXT-C-CA	2.28	121.14	113.38

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	605	GOL	O1-C1-C2-C3
3	A	606	GOL	O1-C1-C2-C3
3	A	607	GOL	C1-C2-C3-O3
4	A	608	DTR	O-C-CA-N
4	A	608	DTR	CA-CB-CG-CD1
4	A	608	DTR	OXT-C-CA-N
3	A	604	GOL	O1-C1-C2-O2
3	A	604	GOL	O1-C1-C2-C3
3	A	607	GOL	O2-C2-C3-O3
3	A	605	GOL	O1-C1-C2-O2
3	B	602	GOL	C1-C2-C3-O3
3	A	606	GOL	O1-C1-C2-O2
3	A	605	GOL	C1-C2-C3-O3
4	A	608	DTR	OXT-C-CA-CB
4	A	608	DTR	O-C-CA-CB


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	604	GOL	1	0
2	A	602	PO4	1	0
3	A	607	GOL	1	0
2	A	603	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	528/534 (98%)	0.05	9 (1%) 70 78	47, 62, 93, 137	0
1	B	525/534 (98%)	0.50	49 (9%) 8 14	51, 81, 122, 153	0
All	All	1053/1068 (98%)	0.27	58 (5%) 25 34	47, 70, 112, 153	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	326	PHE	8.4
1	B	319	LEU	6.1
1	B	45	ILE	5.6
1	B	269	GLY	5.1
1	B	317	TRP	5.0
1	B	270	VAL	4.6
1	B	42	ILE	4.4
1	B	311	LEU	4.3
1	B	44	ARG	4.2
1	B	324	THR	4.2
1	B	313	PHE	4.2
1	B	458	PHE	4.2
1	B	521	TYR	4.0
1	B	43	PRO	3.9
1	B	160	LEU	3.9
1	B	291	GLY	3.8
1	B	266	GLU	3.8
1	B	457	ASN	3.7
1	B	323	ASN	3.6
1	B	529	GLY	3.6
1	A	326	PHE	3.5
1	B	307	ASP	3.4
1	B	268	ASN	3.4
1	B	525	ARG	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	321	PRO	3.4
1	B	318	GLY	3.3
1	B	327	ASN	3.2
1	B	316	LEU	3.2
1	B	325	GLU	3.2
1	B	290	LEU	3.1
1	A	521	TYR	3.1
1	B	267	LYS	3.1
1	B	46	GLY	3.1
1	B	262	PRO	3.0
1	A	525	ARG	3.0
1	A	290	LEU	3.0
1	B	47	VAL	2.9
1	B	259	THR	2.9
1	B	264	ASP	2.9
1	B	159	PHE	2.9
1	B	322	ASP	2.9
1	B	41	THR	2.8
1	A	311	LEU	2.7
1	B	524	LEU	2.7
1	A	322	ASP	2.6
1	A	323	ASN	2.5
1	B	527	LEU	2.5
1	B	321	PRO	2.4
1	B	330	ARG	2.4
1	A	529	GLY	2.2
1	B	281	ALA	2.2
1	B	310	THR	2.2
1	B	523	LEU	2.2
1	B	514	VAL	2.1
1	B	202	LEU	2.0
1	B	452	GLY	2.0
1	B	95	PRO	2.0
1	B	450	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	605	6/6	0.65	0.29	96,114,120,121	0
3	GOL	B	601	6/6	0.70	0.24	79,86,90,97	0
3	GOL	A	607	6/6	0.72	0.25	78,91,95,103	0
2	PO4	A	602	5/5	0.72	0.34	67,71,80,84	5
2	PO4	A	603	5/5	0.79	0.27	87,95,111,116	5
3	GOL	B	602	6/6	0.79	0.40	102,106,115,115	0
3	GOL	A	606	6/6	0.80	0.21	65,87,95,99	0
4	DTR	A	608	15/15	0.85	0.21	58,67,72,76	15
3	GOL	A	604	6/6	0.91	0.19	60,73,77,77	0
2	PO4	A	601	5/5	0.97	0.10	62,65,71,75	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.