



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 24, 2022 – 12:25 am GMT

PDB ID : 8B2J
Title : Crystal structure of human STING in complex with ADU-S100
Authors : Nawrotek, A.; Vuillard, L.; Miallau, L.
Deposited on : 2022-09-14
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.3
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	 6% 67% 7% 26%
1	B	242	 5% 65% 8% 26%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Stimulator of interferon genes protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	179	1459	920	260	273	6	0	1	0
1	B	179	1451	913	259	273	6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	ARG	HIS	variant	UNP Q86WV6
B	232	ARG	HIS	variant	UNP Q86WV6

- Molecule 2 is (1 {R},3 {S},6 {R},8 {R},9 {R},10 {S},12 {S},15 {R},17 {R},18 {R})-8,17-bis(6-aminopurin-9-yl)-3,12-bis(oxidanylidene)-3,12-bis(sulfanyl)-2,4,7,11,13,16-hexaoxa-3 λ^1 {5},12 λ^1 {5}-diphosphatricyclo[13.2.1.0 \wedge {6,10}]octadecane-9,18-diol (three-letter code: GJF) (formula: C₂₀H₂₄N₁₀O₁₀P₂S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	44	20	10	10	2	2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	Na	0	0
			4	4		
4	B	4	Total	Na	0	0
			4	4		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	26	Total	O	0	0
			26	26		
5	B	29	Total	O	0	0
			29	29		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.36Å 67.37Å 136.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	68.16 – 2.17 68.16 – 2.17	Depositor EDS
% Data completeness (in resolution range)	86.0 (68.16-2.17) 86.1 (68.16-2.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 2.18Å)	Xtrriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.242 , 0.285 0.237 , 0.283	Depositor DCC
R_{free} test set	1494 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å ²)	41.8	Xtrriage
Anisotropy	0.022	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.470 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3019	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA, GJF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/1486	0.58	0/2010
1	B	0.37	0/1477	0.59	0/1998
All	All	0.37	0/2963	0.58	0/4008

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1459	0	1438	6	0
1	B	1451	0	1431	9	0
2	A	44	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0
5	A	26	0	0	0	0
5	B	29	0	0	0	0
All	All	3019	0	2869	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:PHE:HA	1:A:290:LEU:HD23	1.92	0.52
1:B:188:ASN:C	1:B:188:ASN:HD22	2.14	0.50
1:B:188:ASN:ND2	1:B:190:LEU:H	2.13	0.47
1:B:206:CYS:SG	1:B:316:GLU:OE2	2.69	0.44
1:B:156:ALA:HB2	1:B:287:GLN:HG3	1.99	0.44
1:B:199:TYR:HD2	1:B:312:ILE:HD11	1.83	0.43
1:A:201:LEU:HD22	1:A:325:LEU:HD11	2.02	0.42
1:B:188:ASN:HD22	1:B:190:LEU:H	1.66	0.41
1:A:239:VAL:HB	1:B:239:VAL:HB	2.01	0.41
1:B:211:ASN:HB3	1:B:214:MET:HG2	2.03	0.41
1:A:233:ALA:HB1	1:B:261:TYR:HB2	2.02	0.40
1:A:199:TYR:HD2	1:A:312:ILE:HD11	1.86	0.40
1:A:204:LEU:HB2	1:A:315:GLN:HG3	2.01	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/242 (73%)	171 (97%)	4 (2%)	1 (1%)	25	24
1	B	175/242 (72%)	168 (96%)	6 (3%)	1 (1%)	25	24
All	All	351/484 (72%)	339 (97%)	10 (3%)	2 (1%)	25	24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	279	PHE
1	B	279	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/210 (75%)	150 (96%)	7 (4%)	27	32
1	B	156/210 (74%)	145 (93%)	11 (7%)	14	14
All	All	313/420 (74%)	295 (94%)	18 (6%)	20	21

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	220	ARG
1	A	232	ARG
1	A	258	VAL
1	A	263	THR
1	A	293	ARG
1	A	315	GLN
1	A	330	LEU
1	B	155	VAL
1	B	175	LEU
1	B	188	ASN
1	B	220	ARG
1	B	229	THR
1	B	263	THR
1	B	287	GLN
1	B	290	LEU
1	B	299	LEU
1	B	334	ARG
1	B	337	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	184	GLN
1	A	196	GLN
1	A	315	GLN
1	A	332	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	188	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 10 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GJF	A	401	-	39,50,50	2.20	2 (5%)	46,78,78	1.45	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GJF	A	401	-	-	5/18/62/62	0/6/7/7

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	GJF	P2-O9	12.61	1.78	1.57
2	A	401	GJF	P2-O6	3.19	1.62	1.57

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	GJF	P2-O6-C9	4.09	133.08	120.16
2	A	401	GJF	P1-O7-C15	3.86	132.34	120.16
2	A	401	GJF	P2-O9-C17	-2.81	114.14	121.13
2	A	401	GJF	O3-P1-O7	2.65	109.64	102.22
2	A	401	GJF	O9-C17-C18	-2.60	102.25	111.68
2	A	401	GJF	O3-P1-O4	-2.41	105.52	114.42
2	A	401	GJF	O9-C17-C16	2.39	118.72	110.08
2	A	401	GJF	C11-C14-N7	2.27	123.80	120.35
2	A	401	GJF	C2-C3-N8	2.24	123.75	120.35

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	GJF	C9-O6-P2-O5
2	A	401	GJF	C5-O3-P1-O7
2	A	401	GJF	C15-O7-P1-O4
2	A	401	GJF	C5-O3-P1-O4
2	A	401	GJF	C9-O6-P2-O9

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/242 (73%)	0.73	14 (7%) 13 13	31, 42, 73, 90	0
1	B	179/242 (73%)	0.73	12 (6%) 17 18	30, 42, 72, 93	0
All	All	358/484 (73%)	0.73	26 (7%) 15 15	30, 42, 73, 93	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	323	PHE	4.4
1	A	337	GLU	4.3
1	B	323	PHE	3.5
1	A	285	LEU	3.3
1	B	338	LYS	3.1
1	B	302	ALA	3.0
1	B	285	LEU	3.0
1	B	281	ARG	2.9
1	B	305	SER	2.8
1	B	280	SER	2.8
1	B	284	ARG	2.7
1	A	338	LYS	2.6
1	A	279	PHE	2.6
1	A	185	HIS	2.6
1	A	304	GLU	2.4
1	B	337	GLU	2.3
1	A	312	ILE	2.3
1	B	303	PRO	2.3
1	A	292	CYS	2.2
1	A	329	VAL	2.2
1	A	303	PRO	2.2
1	B	304	GLU	2.1
1	A	205	ASP	2.1
1	B	292	CYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	180	ARG	2.0
1	A	325	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NA	A	404	1/1	0.81	0.14	52,52,52,52	0
4	NA	B	405	1/1	0.81	0.23	49,49,49,49	0
4	NA	B	404	1/1	0.87	0.20	42,42,42,42	0
4	NA	A	406	1/1	0.91	0.23	58,58,58,58	0
2	GJF	A	401	44/44	0.93	0.17	40,41,42,44	0
4	NA	B	402	1/1	0.95	0.51	46,46,46,46	0
4	NA	A	403	1/1	0.97	0.26	44,44,44,44	0
4	NA	A	405	1/1	0.98	0.06	37,37,37,37	0
4	NA	B	403	1/1	0.98	0.14	40,40,40,40	0
3	CA	A	402	1/1	0.99	0.22	36,36,36,36	0
3	CA	B	401	1/1	0.99	0.16	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.