



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 1, 2022 – 06:24 pm GMT

PDB ID : 8BGM
Title : Crystal structure of the OrfX1-OrfX3 complex from the PMP1 neurotoxin gene cluster
Authors : Kosenina, S.; Stenmark, P.
Deposited on : 2022-10-28
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.31.3
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	
1	C	155	
2	B	491	
2	D	491	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 9696 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	130	1082	702	181	193	6	0	0	0
1	C	131	1089	707	180	196	6	0	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	initiating methionine	UNP A0A5P3XKM0
A	-4	HIS	-	expression tag	UNP A0A5P3XKM0
A	-3	HIS	-	expression tag	UNP A0A5P3XKM0
A	-2	HIS	-	expression tag	UNP A0A5P3XKM0
A	-1	HIS	-	expression tag	UNP A0A5P3XKM0
A	0	HIS	-	expression tag	UNP A0A5P3XKM0
A	1	HIS	-	expression tag	UNP A0A5P3XKM0
A	2	GLU	-	expression tag	UNP A0A5P3XKM0
A	3	ASN	-	expression tag	UNP A0A5P3XKM0
A	4	LEU	-	expression tag	UNP A0A5P3XKM0
A	5	TYR	-	expression tag	UNP A0A5P3XKM0
A	6	PHE	-	expression tag	UNP A0A5P3XKM0
A	7	GLN	-	expression tag	UNP A0A5P3XKM0
A	8	GLY	-	expression tag	UNP A0A5P3XKM0
C	-5	MET	-	initiating methionine	UNP A0A5P3XKM0
C	-4	HIS	-	expression tag	UNP A0A5P3XKM0
C	-3	HIS	-	expression tag	UNP A0A5P3XKM0
C	-2	HIS	-	expression tag	UNP A0A5P3XKM0
C	-1	HIS	-	expression tag	UNP A0A5P3XKM0
C	0	HIS	-	expression tag	UNP A0A5P3XKM0
C	1	HIS	-	expression tag	UNP A0A5P3XKM0
C	2	GLU	-	expression tag	UNP A0A5P3XKM0
C	3	ASN	-	expression tag	UNP A0A5P3XKM0
C	4	LEU	-	expression tag	UNP A0A5P3XKM0
C	5	TYR	-	expression tag	UNP A0A5P3XKM0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	PHE	-	expression tag	UNP A0A5P3XKM0
C	7	GLN	-	expression tag	UNP A0A5P3XKM0
C	8	GLY	-	expression tag	UNP A0A5P3XKM0


- Molecule 2 is a protein called Toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	475	Total	C	N	O	S	0	0	0
			3758	2416	600	727	15			
2	D	476	Total	C	N	O	S	0	0	0
			3767	2422	602	728	15			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Toxin




- Molecule 1: Toxin




- Molecule 2: Toxin



- Molecule 2: Toxin





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.02Å 126.57Å 154.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.37 – 2.70 63.29 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (63.37-2.70) 99.9 (63.29-2.70)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.222 , 0.263 0.221 , 0.262	Depositor DCC
R_{free} test set	2543 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	57.2	Xtrriage
Anisotropy	0.991	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9696	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.09 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3876e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1102	0.77	0/1475
1	C	0.49	0/1109	0.75	0/1484
2	B	0.47	0/3837	0.72	0/5205
2	D	0.49	0/3846	0.72	0/5216
All	All	0.48	0/9894	0.73	0/13380

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1082	0	1095	9	0
1	C	1089	0	1101	7	0
2	B	3758	0	3702	25	0
2	D	3767	0	3715	20	0
All	All	9696	0	9613	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:189:VAL:HG11	2:D:209:VAL:HG11	1.68	0.74
1:A:81:SER:HA	1:A:145:LEU:HD11	1.70	0.71
1:C:16:PHE:CZ	1:C:18:ASP:HB3	2.27	0.69
1:A:47:PHE:HZ	1:A:133:LEU:HD21	1.61	0.66
1:C:11:GLU:HB2	1:C:26:LEU:HD21	1.79	0.63
2:B:126:SER:HA	2:B:179:ILE:HD13	1.83	0.61
1:C:28:CYS:SG	1:C:48:CYS:SG	3.01	0.58
2:D:368:GLY:C	2:D:370:GLY:N	2.56	0.57
2:B:435:ILE:H	2:B:435:ILE:HD12	1.69	0.57
2:D:368:GLY:C	2:D:370:GLY:H	2.09	0.56
1:A:16:PHE:CZ	1:A:18:ASP:HB3	2.41	0.55
2:B:407:LEU:O	2:B:411:VAL:HG13	2.07	0.54
1:C:58:VAL:O	1:C:58:VAL:HG12	2.08	0.53
1:C:144:LEU:O	1:C:147:ILE:HG12	2.09	0.53
2:B:175:LEU:O	2:B:179:ILE:HG13	2.09	0.52
2:B:126:SER:HA	2:B:179:ILE:CD1	2.41	0.51
2:B:372:GLN:OE1	2:B:453:ILE:HD11	2.11	0.50
2:D:56:ILE:HD11	2:D:186:PHE:HE2	1.77	0.49
2:B:230:MET:HA	2:B:230:MET:HE2	1.96	0.48
2:B:105:TYR:CZ	2:B:122:LEU:HD13	2.49	0.47
2:D:5:ARG:HH21	2:D:246:HIS:HA	1.76	0.47
2:D:269:LEU:N	2:D:270:PRO:HD2	2.29	0.47
1:A:57:LYS:N	2:B:157:GLU:OE2	2.47	0.47
1:C:32:ILE:HD12	1:C:42:TYR:HB3	1.97	0.47
2:D:471:ILE:HD12	2:D:471:ILE:O	2.14	0.47
1:A:80:TYR:O	1:A:80:TYR:CG	2.67	0.47
2:B:135:VAL:HG11	2:B:179:ILE:HD11	1.95	0.47
2:D:260:GLU:HG3	2:D:481:GLY:O	2.15	0.47
2:B:260:GLU:HG3	2:B:481:GLY:O	2.15	0.46
2:D:117:VAL:HG11	2:D:190:ASN:HD22	1.79	0.46
1:A:129:LEU:O	1:A:133:LEU:HB2	2.16	0.46
1:A:32:ILE:HD12	1:A:42:TYR:HB3	1.98	0.46
2:D:105:TYR:CZ	2:D:122:LEU:HD13	2.50	0.46
2:B:269:LEU:N	2:B:270:PRO:HD2	2.31	0.46
2:D:288:ASN:HA	2:D:297:GLN:O	2.15	0.45
2:B:12:TRP:HA	2:B:230:MET:CE	2.47	0.45
2:B:288:ASN:HA	2:B:297:GLN:O	2.17	0.44
2:B:105:TYR:HA	2:B:121:ASP:O	2.18	0.44
2:B:11:ASN:O	2:B:239:ARG:NH1	2.45	0.44
2:B:220:LEU:O	2:B:221:ASP:HB2	2.18	0.44
2:D:113:LYS:HA	2:D:113:LYS:HE2	1.99	0.44
2:D:274:MET:CE	2:D:438:THR:C	2.86	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:402:ALA:O	2:B:406:ILE:HG12	2.19	0.43
2:D:105:TYR:HA	2:D:121:ASP:O	2.18	0.43
2:D:272:LEU:HB2	2:D:273:PRO:HD3	2.01	0.43
2:B:65:ILE:N	2:B:65:ILE:HD12	2.33	0.43
2:D:135:VAL:HG12	2:D:136:VAL:HG23	2.01	0.43
2:B:272:LEU:HD23	2:B:284:TYR:CG	2.54	0.43
2:B:262:ARG:HA	2:B:262:ARG:NE	2.34	0.42
1:C:63:CYS:SG	2:D:85:ILE:HD12	2.59	0.42
2:D:402:ALA:O	2:D:406:ILE:HG12	2.20	0.42
2:B:5:ARG:NH2	2:B:248:VAL:O	2.51	0.42
2:B:57:ILE:HD11	2:B:66:ARG:HB3	2.02	0.41
2:B:135:VAL:HG12	2:B:136:VAL:HG23	2.02	0.41
2:B:453:ILE:HD13	2:B:474:LEU:CD2	2.50	0.41
1:A:147:ILE:HD12	1:A:147:ILE:N	2.36	0.41
2:D:118:GLU:CB	2:D:193:LEU:HD11	2.51	0.41
1:A:95:MET:SD	1:A:97:ILE:HD12	2.60	0.41
2:D:205:ARG:HA	2:D:206:PRO:HD3	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/155 (81%)	120 (95%)	6 (5%)	0	100	100
1	C	127/155 (82%)	119 (94%)	8 (6%)	0	100	100
2	B	469/491 (96%)	449 (96%)	20 (4%)	0	100	100
2	D	470/491 (96%)	451 (96%)	19 (4%)	0	100	100
All	All	1192/1292 (92%)	1139 (96%)	53 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/145 (83%)	108 (89%)	13 (11%)	6	15
1	C	122/145 (84%)	114 (93%)	8 (7%)	16	38
2	B	417/430 (97%)	388 (93%)	29 (7%)	15	35
2	D	418/430 (97%)	404 (97%)	14 (3%)	38	67
All	All	1078/1150 (94%)	1014 (94%)	64 (6%)	19	43

All (64) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	18	ASP
1	A	23	MET
1	A	27	PHE
1	A	33	LYS
1	A	64	GLU
1	A	91	SER
1	A	103	HIS
1	A	120	ASP
1	A	122	ASN
1	A	124	MET
1	A	133	LEU
1	A	146	ARG
2	B	16	PHE
2	B	23	VAL
2	B	42	SER
2	B	93	GLU
2	B	114	TYR
2	B	115	LYS
2	B	126	SER
2	B	152	LEU
2	B	157	GLU
2	B	179	ILE
2	B	180	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	190	ASN
2	B	222	LYS
2	B	238	LEU
2	B	239	ARG
2	B	250	GLU
2	B	272	LEU
2	B	314	GLU
2	B	334	ASP
2	B	348	MET
2	B	356	THR
2	B	393	GLU
2	B	394	ASN
2	B	399	VAL
2	B	407	LEU
2	B	435	ILE
2	B	449	THR
2	B	458	GLU
2	B	473	GLU
1	C	22	SER
1	C	27	PHE
1	C	33	LYS
1	C	53	SER
1	C	83	GLU
1	C	103	HIS
1	C	120	ASP
1	C	124	MET
2	D	16	PHE
2	D	42	SER
2	D	115	LYS
2	D	126	SER
2	D	172	LYS
2	D	180	SER
2	D	193	LEU
2	D	222	LYS
2	D	280	THR
2	D	310	LEU
2	D	356	THR
2	D	399	VAL
2	D	437	LYS
2	D	446	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	39	ASN
1	A	122	ASN
2	B	278	ASN
1	C	24	ASN
1	C	39	ASN
2	D	190	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	130/155 (83%)	0.74	7 (5%) 25 24	37, 50, 71, 95	0
1	C	131/155 (84%)	0.64	4 (3%) 49 49	40, 52, 75, 93	0
2	B	475/491 (96%)	0.78	27 (5%) 23 22	37, 51, 72, 105	0
2	D	476/491 (96%)	0.73	12 (2%) 57 59	37, 49, 72, 104	0
All	All	1212/1292 (93%)	0.74	50 (4%) 37 36	37, 50, 73, 105	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	81	ASP	3.5
2	B	51	PHE	3.2
2	B	194	TYR	3.2
2	B	89	ASN	3.1
2	D	155	PHE	3.0
2	D	115	LYS	3.0
2	D	419	ALA	3.0
2	B	82	LYS	2.9
2	B	354	CYS	2.9
1	A	62	LEU	2.7
2	B	182	PHE	2.7
2	B	106	PHE	2.7
2	B	181	LEU	2.6
2	B	170	LEU	2.6
1	C	18	ASP	2.6
2	B	48	LYS	2.5
1	A	123	TYR	2.5
2	D	287	ILE	2.5
2	B	323	TYR	2.4
2	D	268	LEU	2.4
2	D	81	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	331	LEU	2.4
1	A	128	ILE	2.4
1	A	121	GLU	2.3
2	B	349	GLY	2.2
2	B	441	ILE	2.2
2	B	394	ASN	2.2
2	B	92	PHE	2.2
1	A	115	ILE	2.2
2	B	43	SER	2.2
2	B	29	ASP	2.2
2	B	38	GLU	2.2
2	B	282	GLU	2.2
1	C	142	ARG	2.2
2	B	297	GLN	2.1
2	B	399	VAL	2.1
2	D	441	ILE	2.1
1	A	102	ILE	2.1
2	B	313	VAL	2.1
2	D	229	CYS	2.1
1	C	94	PHE	2.1
2	B	440	ASP	2.1
1	C	54	ILE	2.1
2	B	286	VAL	2.0
2	D	348	MET	2.0
2	D	65	ILE	2.0
2	B	151	PHE	2.0
2	B	422	PHE	2.0
2	D	151	PHE	2.0
1	A	135	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.