

Oct 16, 2024 - 12:07 AM JST

			8HFF EMD-34719
Ti	itle	:	Cryo-EM structure of human norepinephrine transporter NET in the presence
			of norepinephrine in an inward-open state at resolution of 2.9 angstrom.
Auth	ors	:	Tan, J.; Xiao, Y.; Kong, F.; Lei, J.; Yuan, Y.; Yan, C.
Deposited	on	:	2022-11-10
Resolut	ion	:	2.86 Å(reported)
Th	is is	a I	Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.86 Å.

Sidechain outliers

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



206894

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

16415

Mol	Chain	Length	Quality of chain		
1	А	617	74%	15%	11%



2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 4399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Sodium-dependent noradrenaline transporter.

Mol	Chain	Residues	Atoms			AltConf	Trace		
1	А	547	Total 4373	C 2937	N 679	0 737	S 20	0	0

• Molecule 2 is L-NOREPINEPHRINE (three-letter code: LNR) (formula: $C_8H_{11}NO_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	AltConf
0	Δ	1	Total C N O	0
	A	1	12 8 1 3	0
0	Δ	1	Total C N O	0
	A	1	12 8 1 3	0

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	AltConf
3	А	2	Total O 2 2	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sodium-dependent noradrenaline transporter





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	304363	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	1400	Depositor
Magnification	Not provided	
Image detector	GATAN K3 $(6k \times 4k)$	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LNR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.29	0/4514	0.44	0/6156

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4373	0	4364	51	0
2	А	24	0	20	0	0
3	А	2	0	0	0	0
All	All	4399	0	4384	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:337:ASN:OD1	1:A:338:ASN:N	2.29	0.66

Continued on next page...



Continued from prev		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:238:LEU:HD13	1:A:458:LEU:HD11	1.80	0.63
1:A:238:LEU:HD13	1:A:458:LEU:CD1	2.29	0.62
1:A:278:LEU:HD12	1:A:356:VAL:HG13	1.83	0.60
1:A:256:VAL:HG22	1:A:425:GLU:OE2	2.02	0.59
1:A:175:ASP:N	1:A:175:ASP:OD1	2.37	0.58
1:A:89:ASN:HB3	1:A:307:VAL:HG22	1.86	0.56
1:A:118:GLN:NE2	1:A:508:MET:SD	2.80	0.55
1:A:289:ASN:OD1	1:A:290:GLY:N	2.40	0.55
1:A:131:CYS:HG	1:A:134:PHE:HE1	1.56	0.54
1:A:148:VAL:HG21	1:A:323:PHE:CE1	2.43	0.54
1:A:148:VAL:HG21	1:A:323:PHE:HE1	1.74	0.53
1:A:66:LEU:O	1:A:70:VAL:HG22	2.10	0.51
1:A:325:VAL:HA	1:A:426:ALA:HB2	1.94	0.50
1:A:328:ALA:O	1:A:331:SER:OG	2.26	0.48
1:A:278:LEU:O	1:A:282:VAL:HG23	2.13	0.48
1:A:256:VAL:HG21	1:A:332:TYR:CE2	2.48	0.48
1:A:101:PHE:CZ	1:A:532:VAL:HG21	2.49	0.47
1:A:436:GLN:HA	1:A:436:GLN:OE1	2.15	0.47
1:A:374:VAL:HG22	1:A:375:ASN:N	2.31	0.46
1:A:75:ASP:OD1	1:A:76:LEU:N	2.48	0.46
1:A:368:MET:SD	1:A:390:LEU:HD12	2.56	0.46
1:A:547:ASP:OD1	1:A:547:ASP:N	2.39	0.46
1:A:578:LEU:C	1:A:578:LEU:HD12	2.36	0.46
1:A:362:PHE:HA	1:A:365:LEU:HG	1.99	0.45
1:A:500:ARG:NH2	1:A:597:GLU:OE2	2.49	0.45
1:A:138:GLY:O	1:A:141:VAL:HG22	2.17	0.45
1:A:305:ALA:HB1	1:A:535:VAL:HG11	1.99	0.44
1:A:534:VAL:O	1:A:538:ILE:HG12	2.16	0.44
1:A:612:GLN:OE1	1:A:612:GLN:N	2.48	0.44
1:A:283:THR:O	1:A:283:THR:OG1	2.33	0.44
1:A:365:LEU:HB2	1:A:379:VAL:HG21	1.98	0.44
1:A:75:ASP:OD1	1:A:419:SER:OG	2.14	0.43
1:A:438:LEU:HD22	1:A:445:PHE:CG	2.53	0.43
1:A:262:VAL:HG12	1:A:262:VAL:O	2.19	0.43
1:A:608:GLN:O	1:A:614:TRP:NE1	2.48	0.43
1:A:116:LEU:HD11	1:A:492:VAL:HG21	2.00	0.43
1:A:154:VAL:HG21	1:A:460:CYS:HB2	2.01	0.42
1:A:266:THR:HG21	1:A:418:ASP:OD2	2.20	0.42
1:A:177:GLY:N	1:A:181:ASN:OD1	2.53	0.42
1:A:424:MET:CE	1:A:449:VAL:HG12	2.50	0.42
1:A:361:ILE:HD13	1:A:390:LEU:HD23	2.02	0.41

Continued from previous page...

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:221:LEU:CD1	1:A:466:ILE:HG22	2.51	0.41
1:A:126:THR:O	1:A:126:THR:HG22	2.20	0.41
1:A:461:ILE:O	1:A:461:ILE:CG2	2.68	0.41
1:A:107:MET:HB3	1:A:108:PRO:HD3	2.02	0.41
1:A:301:ARG:HG2	1:A:307:VAL:HG21	2.03	0.41
1:A:482:LEU:HD13	1:A:532:VAL:HG12	2.03	0.41
1:A:525:SER:HB2	1:A:526:PRO:HD3	2.02	0.41
1:A:68:SER:HA	1:A:263:VAL:HG21	2.03	0.40
1:A:571:ILE:O	1:A:574:ILE:HG12	2.20	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles		
1	А	543/617~(88%)	528~(97%)	15 (3%)	0	100	100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	461/519 (89%)	447 (97%)	14 (3%)	36 62		



Mol	Chain	Res	Type
1	А	164	PHE
1	А	242	MET
1	А	253	TRP
1	А	257	LYS
1	А	300	TYR
1	А	317	PHE
1	А	336	ASP
1	А	338	ASN
1	А	363	SER
1	А	396	SER
1	А	424	MET
1	А	459	PHE
1	А	498	VAL
1	А	547	ASP

All (14) residues with a non-rotameric sidechain are listed below:

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type	
1	А	296	HIS	

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain Res		Link	Bond lengths			Bond angles			
	Type	Unain	Res	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	LNR	А	701	-	12,12,12	0.39	0	15, 16, 16	0.42	0
2	LNR	А	702	-	12,12,12	0.43	0	15, 16, 16	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LNR	А	701	-	-	4/6/6/6	0/1/1/1
2	LNR	А	702	-	-	1/6/6/6	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	701	LNR	CAG-CAK-CAL-CAH
2	А	701	LNR	CAF-CAK-CAL-CAH
2	А	701	LNR	CAF-CAK-CAL-OAD
2	А	701	LNR	CAG-CAK-CAL-OAD
2	А	702	LNR	CAF-CAK-CAL-OAD


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the



average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

