



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2024 – 07:21 pm BST

PDB ID : 8PJF  
Title : Human Leukocyte Antigen class II allotype DR1 presenting P11T->R modified influenza A virus haemagglutinin (HA)306-318 PKYVKQNTLKLAR  
Authors : MacLachlan, B.J.; Wall, A.; Greenshields-Watson, A.L.; Cole, D.K.; Rizkallah, P.J.; Godkin, A.J.  
Deposited on : 2023-06-23  
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	 3% 87% 10%
2	B	194	 2% 89% 10%
3	C	13	 8% 85% 15%

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 3654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1531	1000	245	280	6	0	9	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	MET	-	initiating methionine	UNP P01903
A	-2	GLY	-	expression tag	UNP P01903
A	-1	SER	-	expression tag	UNP P01903
A	0	MET	-	expression tag	UNP P01903

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DRB1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	191	1613	1019	288	299	7	0	7	0

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	MET	-	initiating methionine	UNP P01911
B	-2	GLY	-	expression tag	UNP P01911
B	-1	SER	-	expression tag	UNP P01911
B	0	MET	-	expression tag	UNP P01911
B	11	LEU	PRO	variant	UNP P01911
B	13	PHE	ARG	variant	UNP P01911
B	26	LEU	PHE	variant	UNP P01911
B	28	GLU	ASP	variant	UNP P01911
B	30	CYS	TYR	variant	UNP P01911
B	31	ILE	PHE	variant	UNP P01911
B	47	TYR	PHE	variant	UNP P01911
B	67	LEU	ILE	variant	UNP P01911

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	71	ARG	ALA	variant	UNP P01911
B	86	GLY	VAL	variant	UNP P01911
B	96	GLU	GLN	variant	UNP P01911
B	133	ARG	LEU	variant	UNP P01911
B	142	VAL	MET	variant	UNP P01911


- Molecule 3 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	13	110	71	21	18	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	13	ARG	THR	engineered mutation	UNP P03435

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0
4	A	1	5	4	1	0	0
4	B	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	150	Total O 150 150	0	0
6	B	148	Total O 148 148	0	0
6	C	8	Total O 8 8	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain A: 




- Molecule 2: HLA class II histocompatibility antigen, DRB1 beta chain

Chain B: 



- Molecule 3: Hemagglutinin HA2 chain

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.35Å 134.43Å 40.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.14 – 1.48 54.14 – 1.48	Depositor EDS
% Data completeness (in resolution range)	99.7 (54.14-1.48) 99.7 (54.14-1.48)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.46 (at 1.48Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.167 , 0.192 0.165 , 0.190	Depositor DCC
$R_{free}$ test set	4216 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.0	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3654	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/1604	0.85	0/2183
2	B	0.70	0/1676	0.87	1/2273 (0.0%)
3	C	0.65	0/111	0.82	0/145
All	All	0.71	0/3391	0.86	1/4601 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	158	LEU	CA-CB-CG	7.29	132.07	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	94[A]	ARG	Sidechain
2	B	94[B]	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1531	0	1502	14	1
2	B	1613	0	1563	12	1
3	C	110	0	127	2	0
4	A	10	0	0	0	0
4	B	10	0	0	1	0
4	C	10	0	0	0	0
5	A	20	0	30	2	0
5	B	44	0	66	5	0
6	A	150	0	0	1	0
6	B	148	0	0	3	0
6	C	8	0	0	0	0
All	All	3654	0	3288	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:55[A]:ARG:NH1	4:B:202:SO4:O3	2.20	0.74
1:A:55:GLU:OE1	3:C:2:LYS:HD2	2.00	0.62
1:A:156:SER:OG	1:A:158:GLU:HG2	2.00	0.61
5:B:206:EDO:H11	6:B:306:HOH:O	2.02	0.60
1:A:111[B]:LYS:HG2	1:A:140:ARG:CZ	2.37	0.55
2:B:67:LEU:HD12	6:B:346:HOH:O	2.07	0.54
1:A:39:LYS:HB3	1:A:60:LEU:HD11	1.92	0.52
1:A:39:LYS:HD3	1:A:60:LEU:HD13	1.92	0.51
2:B:99:VAL:HB	5:B:206:EDO:H12	1.94	0.49
1:A:138:LEU:HB2	1:A:146[B]:ARG:HG2	1.96	0.48
2:B:162[B]:GLU:HB3	5:B:210:EDO:O1	2.13	0.48
1:A:96:PRO:HB3	5:A:205:EDO:H12	1.98	0.46
2:B:147:LEU:HB2	5:B:212:EDO:H21	1.99	0.45
2:B:9:TRP:CH2	2:B:30:CYS:HB3	2.52	0.45
3:C:2:LYS:HB3	3:C:2:LYS:HE3	1.69	0.44
2:B:103:PRO:HD3	2:B:188[A]:TRP:CZ3	2.54	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:ASP:O	6:A:301:HOH:O	2.21	0.43
1:A:111[B]:LYS:HB3	1:A:111[B]:LYS:HE3	1.87	0.43
2:B:18:PHE:HB2	2:B:23:ARG:HB3	2.00	0.42
2:B:40:PHE:HB2	2:B:47:TYR:CE1	2.54	0.42
2:B:52:GLU:HG2	2:B:55[A]:ARG:HE	1.84	0.42
2:B:59:GLU:HG3	6:B:376:HOH:O	2.19	0.42
1:A:97:VAL:HG21	1:A:178:TRP:HZ2	1.85	0.41
1:A:105:LEU:HG	1:A:153:PHE:CE1	2.56	0.41
1:A:129:THR:O	1:A:132:VAL:HG22	2.21	0.41
2:B:99:VAL:CB	5:B:206:EDO:H12	2.51	0.40
1:A:97:VAL:O	5:A:205:EDO:H11	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ARG:NH2	2:B:106:THR:OG1[1_556]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	187/186 (100%)	186 (100%)	1 (0%)	0	100	100
2	B	196/194 (101%)	191 (97%)	5 (3%)	0	100	100
3	C	11/13 (85%)	11 (100%)	0	0	100	100
All	All	394/393 (100%)	388 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/169 (103%)	174 (100%)	0	100	100
2	B	179/174 (103%)	178 (99%)	1 (1%)	86	72
3	C	12/12 (100%)	11 (92%)	1 (8%)	11	1
All	All	365/355 (103%)	363 (100%)	2 (0%)	86	77

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	19	ASN
3	C	5	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
2	B	19	ASN
2	B	112	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	B	208	-	3,3,3	0.44	0	2,2,2	0.30	0
5	EDO	A	203	-	3,3,3	0.35	0	2,2,2	0.76	0
4	SO4	A	201	-	4,4,4	0.16	0	6,6,6	0.28	0
5	EDO	B	206	-	3,3,3	0.54	0	2,2,2	0.90	0
5	EDO	A	205	-	3,3,3	0.52	0	2,2,2	0.23	0
4	SO4	C	102	-	4,4,4	0.17	0	6,6,6	0.13	0
4	SO4	B	202	-	4,4,4	0.15	0	6,6,6	0.12	0
5	EDO	B	211	-	3,3,3	0.53	0	2,2,2	0.22	0
5	EDO	B	210	-	3,3,3	0.58	0	2,2,2	0.24	0
4	SO4	A	202	-	4,4,4	0.18	0	6,6,6	0.06	0
5	EDO	B	205	-	3,3,3	0.65	0	2,2,2	0.15	0
5	EDO	A	204	-	3,3,3	0.56	0	2,2,2	0.02	0
5	EDO	B	207	-	3,3,3	0.46	0	2,2,2	0.52	0
4	SO4	C	101	-	4,4,4	0.15	0	6,6,6	0.16	0
5	EDO	B	203	-	3,3,3	0.57	0	2,2,2	0.16	0
5	EDO	B	204	-	3,3,3	0.42	0	2,2,2	0.25	0
5	EDO	B	212	-	3,3,3	0.45	0	2,2,2	0.11	0
4	SO4	B	201	-	4,4,4	0.17	0	6,6,6	0.12	0
5	EDO	A	206	-	3,3,3	0.65	0	2,2,2	0.16	0
5	EDO	A	207	-	3,3,3	0.44	0	2,2,2	0.75	0
5	EDO	B	209	-	3,3,3	0.44	0	2,2,2	0.44	0
5	EDO	B	213	-	3,3,3	0.44	0	2,2,2	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	210	-	-	0/1/1/1	-
5	EDO	B	208	-	-	1/1/1/1	-
5	EDO	B	204	-	-	0/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	205	-	-	0/1/1/1	-
5	EDO	B	212	-	-	0/1/1/1	-
5	EDO	A	204	-	-	0/1/1/1	-
5	EDO	A	203	-	-	1/1/1/1	-
5	EDO	B	206	-	-	1/1/1/1	-
5	EDO	B	207	-	-	1/1/1/1	-
5	EDO	A	206	-	-	0/1/1/1	-
5	EDO	A	207	-	-	1/1/1/1	-
5	EDO	A	205	-	-	0/1/1/1	-
5	EDO	B	209	-	-	1/1/1/1	-
5	EDO	B	203	-	-	0/1/1/1	-
5	EDO	B	213	-	-	0/1/1/1	-
5	EDO	B	211	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	208	EDO	O1-C1-C2-O2
5	A	207	EDO	O1-C1-C2-O2
5	B	209	EDO	O1-C1-C2-O2
5	B	206	EDO	O1-C1-C2-O2
5	B	207	EDO	O1-C1-C2-O2
5	B	211	EDO	O1-C1-C2-O2
5	A	203	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	206	EDO	3	0
5	A	205	EDO	2	0
4	B	202	SO4	1	0
5	B	210	EDO	1	0
5	B	212	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	180/186 (96%)	-0.32	5 (2%) 53 57	16, 24, 48, 77	0
2	B	191/194 (98%)	-0.31	3 (1%) 72 75	16, 26, 49, 59	0
3	C	13/13 (100%)	-0.30	1 (7%) 13 14	22, 32, 46, 76	0
All	All	384/393 (97%)	-0.31	9 (2%) 60 65	16, 25, 49, 77	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	LYS	5.5
1	A	158	GLU	3.6
2	B	106	THR	3.2
1	A	100	ARG	3.2
3	C	13	ARG	2.8
1	A	181	ASP	2.6
2	B	109	LEU	2.5
2	B	105	LYS	2.5
1	A	99	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	B	212	4/4	0.66	0.31	54,54,55,55	0
5	EDO	B	208	4/4	0.68	0.25	62,62,63,63	0
5	EDO	A	205	4/4	0.70	0.21	61,61,61,62	0
5	EDO	B	207	4/4	0.70	0.20	42,44,46,47	0
5	EDO	B	209	4/4	0.71	0.27	59,60,60,60	0
4	SO4	A	201	5/5	0.72	0.15	107,107,107,107	0
5	EDO	A	204	4/4	0.75	0.25	57,57,58,58	0
5	EDO	B	210	4/4	0.75	0.15	38,38,40,40	0
5	EDO	B	205	4/4	0.75	0.16	52,52,53,53	0
5	EDO	B	211	4/4	0.78	0.19	60,60,60,61	0
4	SO4	B	201	5/5	0.81	0.33	101,102,102,102	0
5	EDO	B	213	4/4	0.82	0.30	59,59,59,60	0
5	EDO	A	206	4/4	0.84	0.15	34,35,35,35	0
5	EDO	A	207	4/4	0.84	0.23	40,42,43,44	0
5	EDO	B	206	4/4	0.85	0.24	41,43,44,45	0
4	SO4	A	202	5/5	0.86	0.28	89,89,89,89	0
4	SO4	C	101	5/5	0.87	0.44	96,97,97,97	0
5	EDO	B	203	4/4	0.89	0.16	37,38,39,39	0
4	SO4	C	102	5/5	0.91	0.20	80,81,81,81	0
4	SO4	B	202	5/5	0.92	0.50	96,96,96,96	0
5	EDO	A	203	4/4	0.93	0.12	53,53,54,54	0
5	EDO	B	204	4/4	0.97	0.14	48,48,48,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.