



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 2, 2025 – 01:41 am GMT

PDB ID : 8RVH  
Title : Structure of the binding domain of BoNT/A mutant Y1117V/H1253K in complex with the GD1a ganglioside receptor  
Authors : Masuyer, G.; Stenmark, P.  
Deposited on : 2024-02-01  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	
2	G	4	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Botulinum neurotoxin A heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	421	Total	C	N	O	S	0	3	0
			3460	2203	597	645	15			

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	854	MET	-	initiating methionine	UNP P0DPI0
A	855	GLY	-	expression tag	UNP P0DPI0
A	856	SER	-	expression tag	UNP P0DPI0
A	857	SER	-	expression tag	UNP P0DPI0
A	858	HIS	-	expression tag	UNP P0DPI0
A	859	HIS	-	expression tag	UNP P0DPI0
A	860	HIS	-	expression tag	UNP P0DPI0
A	861	HIS	-	expression tag	UNP P0DPI0
A	862	HIS	-	expression tag	UNP P0DPI0
A	863	HIS	-	expression tag	UNP P0DPI0
A	864	SER	-	expression tag	UNP P0DPI0
A	865	SER	-	expression tag	UNP P0DPI0
A	866	GLY	-	expression tag	UNP P0DPI0
A	867	LEU	-	expression tag	UNP P0DPI0
A	868	VAL	-	expression tag	UNP P0DPI0
A	869	PRO	-	expression tag	UNP P0DPI0
A	870	ARG	-	expression tag	UNP P0DPI0
A	871	GLY	-	expression tag	UNP P0DPI0
A	872	SER	-	expression tag	UNP P0DPI0
A	873	HIS	-	expression tag	UNP P0DPI0
A	874	MET	-	expression tag	UNP P0DPI0
A	875	ASP	-	expression tag	UNP P0DPI0
A	1117	VAL	TYR	engineered mutation	UNP P0DPI0
A	1253	LYS	HIS	engineered mutation	UNP P0DPI0
A	1297	GLN	-	expression tag	UNP P0DPI0


- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-b

eta-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	4	Total	C	N	O	0	0	0
			48	26	1	21			

- Molecule 3 is N-acetyl-alpha-neuraminic acid (three-letter code: SIA) (formula:  $C_{11}H_{19}NO_9$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			20	11	1	8		
3	A	1	Total	C	N	O	0	0
			20	11	1	8		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	428	Total	O	0	0
			428	428		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Botulinum neurotoxin A heavy chain



- Molecule 2: beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.64Å 43.58Å 133.21Å 90.00° 95.71° 90.00°	Depositor
Resolution (Å)	69.84 – 1.80 69.84 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.8 (69.84-1.80) 94.8 (69.84-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.92 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.167 , 0.207 0.175 , 0.214	Depositor DCC
$R_{free}$ test set	2068 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.5	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3976	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NGA, SIA, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/3533	0.76	2/4774 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1061	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	A	1076	ASP	CB-CG-OD1	5.12	122.91	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1138	GLY	Peptide,Mainchain

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3460	0	3412	12	0
2	G	48	0	42	5	0
3	A	40	0	34	5	0
4	A	428	0	0	2	1
All	All	3976	0	3488	17	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1301:SIA:C2	2:G:4:GAL:O3	1.98	1.10
1:A:1236:LYS:NZ	1:A:1280:CYS:SG	2.43	0.92
3:A:1301:SIA:C2	2:G:4:GAL:HO3	1.88	0.84
1:A:876:THR:HG21	1:A:1077:LYS:HD2	1.69	0.73
1:A:960:ASN:HD21	1:A:1061:ARG:H	1.45	0.62
3:A:1301:SIA:C2	2:G:4:GAL:C3	2.86	0.54
1:A:1003:GLN:HA	1:A:1011:ILE:HD11	1.91	0.52
1:A:996[A]:ARG:NH2	4:A:1405:HOH:O	2.43	0.48
1:A:1120:ASN:ND2	4:A:1409:HOH:O	2.48	0.47
3:A:1302:SIA:C2	2:G:2:GAL:O3	2.63	0.46
1:A:960:ASN:ND2	1:A:1061:ARG:H	2.12	0.44
1:A:1026:ASN:HD22	1:A:1040:PRO:HA	1.82	0.44
3:A:1302:SIA:O6	2:G:2:GAL:O3	2.35	0.44
1:A:880:ASN:OD1	1:A:882:ARG:CZ	2.66	0.43
1:A:913:GLN:HG2	1:A:1070:LYS:HD3	2.01	0.42
1:A:1227:ASN:N	1:A:1231:ILE:O	2.50	0.41
1:A:1158:THR:HG23	1:A:1185:VAL:O	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1574:HOH:O	4:A:1574:HOH:O[2_555]	1.94	0.26



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/444 (95%)	400 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/408 (94%)	384 (100%)	1 (0%)	91	90

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1065	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	954	ASN
1	A	960	ASN
1	A	991	GLN
1	A	1026	ASN
1	A	1038	GLN
1	A	1046	ASN
1	A	1172	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1233	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	BGC	G	1	2	12,12,12	0.65	0	17,17,17	1.04	1 (5%)
2	GAL	G	2	2	11,11,12	0.29	0	15,15,17	1.65	4 (26%)
2	NGA	G	3	2	14,14,15	0.30	0	17,19,21	1.26	1 (5%)
2	GAL	G	4	2	11,11,12	0.39	0	15,15,17	1.66	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	G	1	2	-	0/2/22/22	0/1/1/1
2	GAL	G	2	2	-	0/2/19/22	0/1/1/1
2	NGA	G	3	2	-	0/6/23/26	0/1/1/1
2	GAL	G	4	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	4	GAL	C2-C3-C4	3.99	117.80	110.89
2	G	3	NGA	O3-C3-C2	-3.79	101.62	109.47
2	G	2	GAL	O5-C1-C2	3.26	115.81	110.77
2	G	2	GAL	O3-C3-C2	-3.17	103.93	109.99
2	G	2	GAL	C1-O5-C5	3.06	116.34	112.19
2	G	4	GAL	O3-C3-C2	-2.90	104.45	109.99
2	G	4	GAL	O5-C5-C6	2.77	111.55	107.20
2	G	1	BGC	C1-O5-C5	2.70	118.77	113.66
2	G	2	GAL	C1-C2-C3	2.45	112.68	109.67

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	4	GAL	3	0
2	G	2	GAL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SIA	A	1301	-	20,20,21	1.04	1 (5%)	24,28,31	1.80	6 (25%)
3	SIA	A	1302	-	20,20,21	1.09	2 (10%)	24,28,31	1.90	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIA	A	1301	-	-	8/18/34/38	0/1/1/1
3	SIA	A	1302	-	-	6/18/34/38	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1302	SIA	C4-C5	2.53	1.55	1.53
3	A	1302	SIA	C2-C1	-2.41	1.50	1.52
3	A	1301	SIA	O6-C6	-2.04	1.40	1.44

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1302	SIA	O6-C2-C3	5.27	117.72	110.46
3	A	1302	SIA	O6-C2-C1	4.55	116.62	107.70
3	A	1302	SIA	C3-C4-C5	4.44	116.83	111.46
3	A	1301	SIA	O6-C2-C1	4.09	115.72	107.70
3	A	1301	SIA	C11-C10-N5	3.14	121.41	116.10
3	A	1301	SIA	O4-C4-C3	-2.78	103.06	109.94
3	A	1301	SIA	C8-C7-C6	2.37	117.53	113.03
3	A	1301	SIA	C3-C4-C5	2.13	114.04	111.46
3	A	1301	SIA	C6-C5-N5	2.06	114.33	110.91

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1301	SIA	C5-C6-C7-C8
3	A	1301	SIA	C5-C6-C7-O7
3	A	1301	SIA	O6-C6-C7-C8
3	A	1301	SIA	O6-C6-C7-O7
3	A	1302	SIA	C5-C6-C7-O7
3	A	1302	SIA	O6-C6-C7-O7
3	A	1301	SIA	C11-C10-N5-C5
3	A	1301	SIA	O10-C10-N5-C5
3	A	1302	SIA	O8-C8-C9-O9
3	A	1301	SIA	C7-C8-C9-O9
3	A	1301	SIA	O8-C8-C9-O9
3	A	1302	SIA	C7-C8-C9-O9
3	A	1302	SIA	C5-C6-C7-C8
3	A	1302	SIA	C4-C5-N5-C10


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1301	SIA	3	0
3	A	1302	SIA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	421/444 (94%)	-0.16	19 (4%) 39 36	9, 19, 44, 104	3 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1297	GLN	5.1
1	A	1228	ASP	4.9
1	A	1229	GLN	4.7
1	A	1227	ASN	4.3
1	A	1232	THR	4.0
1	A	1231	ILE	3.7
1	A	1234	LYS	3.5
1	A	1274	SER	3.4
1	A	1277	THR	3.4
1	A	1278	LEU	3.2
1	A	1233	ASN	3.0
1	A	1230	GLY	3.0
1	A	1296	LEU	2.9
1	A	1275	SER	2.7
1	A	1168	GLY	2.7
1	A	1276	ARG	2.6
1	A	874	MET	2.3
1	A	959	ASN	2.3
1	A	1235	CYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SIA	A	1302	20/21	0.46	0.24	80,99,116,118	0
3	SIA	A	1301	20/21	0.83	0.14	37,43,56,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around SIA A 1302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SIA A 1301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.