

Full wwPDB X-ray Structure Validation Report (i)

Aug 5, 2025 – 06:10 PM EDT

PDB ID : 8TD7 / pdb 00008td7

Title: Structure of PYCR1 complexed with 2S-hydroxy-3-methylbutyric acid

Authors: Tanner, J.J.; Meeks, K.R.

Deposited on : 2023-07-02

Resolution : 1.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 2.0rc1

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.45.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \text{Entries})$	Similar resolution $(\#\text{Entries, resolution range}(\mathring{A}))$
R_{free}	164625	6077 (1.64-1.60)
Clashscore	180529	6617 (1.64-1.60)
Ramachandran outliers	177936	6498 (1.64-1.60)
Sidechain outliers	177891	6497 (1.64-1.60)
RSRZ outliers	164620	6075 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	316	84%	00/ 110/
	11	310	4%	6% 11%
1	В	316	82%	6% 12%
1	С	316	81%	5% 13%
1	D	316	7% 84%	• 12%
1	E	316	5% 82%	• 13%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10783 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Pyrroline-5-carboxylate reductase 1, mitochondrial.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
1	A	282	Total	С	N	О	S	0	4	0
1	A	202	2053	1297	361	380	15	U		
1	В	279	Total	С	N	О	S	0	3	0
1	Ъ	219	2026	1281	355	377	13	0	3	
1	C	275	Total	С	N	О	S	0	5	0
1		210	1986	1252	352	369	13	0		
1	D	277	Total	С	N	О	S	0	2	0
1	D	211	1956	1236	341	366	13	0	<u> </u>	
1	Е	274	Total	С	N	О	S	0	2	0
1	ינו	214	1948	1229	344	362	13	U	2	

There are 110 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP P32322
A	-20	HIS	-	expression tag	UNP P32322
A	-19	HIS	-	expression tag	UNP P32322
A	-18	HIS	-	expression tag	UNP P32322
A	-17	HIS	-	expression tag	UNP P32322
A	-16	HIS	_	expression tag	UNP P32322
A	-15	HIS	-	expression tag	UNP P32322
A	-14	SER	-	expression tag	UNP P32322
A	-13	SER	-	expression tag	UNP P32322
A	-12	GLY	-	expression tag	UNP P32322
A	-11	VAL	_	expression tag	UNP P32322
A	-10	ASP	-	expression tag	UNP P32322
A	-9	LEU	-	expression tag	UNP P32322
A	-8	GLY	-	expression tag	UNP P32322
A	-7	THR	-	expression tag	UNP P32322
A	-6	GLU	-	expression tag	UNP P32322
A	-5	ASN	-	expression tag	UNP P32322
A	-4	LEU	-	expression tag	UNP P32322
A	-3	TYR	-	expression tag	UNP P32322



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	-2	PHE	-	expression tag	UNP P32322
A	-1	GLN	-	expression tag	UNP P32322
A	0	SER	_	expression tag	UNP P32322
В	-21	MET	-	initiating methionine	UNP P32322
В	-20	HIS	-	expression tag	UNP P32322
В	-19	HIS	-	expression tag	UNP P32322
В	-18	HIS	-	expression tag	UNP P32322
В	-17	HIS	-	expression tag	UNP P32322
В	-16	HIS	-	expression tag	UNP P32322
В	-15	HIS	-	expression tag	UNP P32322
В	-14	SER	-	expression tag	UNP P32322
В	-13	SER	-	expression tag	UNP P32322
В	-12	GLY	-	expression tag	UNP P32322
В	-11	VAL	-	expression tag	UNP P32322
В	-10	ASP	-	expression tag	UNP P32322
В	-9	LEU	-	expression tag	UNP P32322
В	-8	GLY	-	expression tag	UNP P32322
В	-7	THR	-	expression tag	UNP P32322
В	-6	GLU	-	expression tag	UNP P32322
В	-5	ASN	-	expression tag	UNP P32322
В	-4	LEU	-	expression tag	UNP P32322
В	-3	TYR	-	expression tag	UNP P32322
В	-2	PHE	_	expression tag	UNP P32322
В	-1	GLN	-	expression tag	UNP P32322
В	0	SER	-	expression tag	UNP P32322
С	-21	MET	_	initiating methionine	UNP P32322
С	-20	HIS	-	expression tag	UNP P32322
С	-19	HIS	-	expression tag	UNP P32322
С	-18	HIS	_	expression tag	UNP P32322
С	-17	HIS	-	expression tag	UNP P32322
С	-16	HIS	-	expression tag	UNP P32322
С	-15	HIS	_	expression tag	UNP P32322
С	-14	SER	-	expression tag	UNP P32322
С	-13	SER	_	expression tag	UNP P32322
С	-12	GLY	-	expression tag	UNP P32322
С	-11	VAL	-	expression tag	UNP P32322
С	-10	ASP	-	expression tag	UNP P32322
С	-9	LEU	-	expression tag	UNP P32322
С	-8	GLY		expression tag	UNP P32322
С	-7	THR	-	expression tag	UNP P32322
С	-6	GLU	-	expression tag	UNP P32322
С	-5	ASN	-	expression tag	UNP P32322



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
С	-4	LEU	-	expression tag	UNP P32322
С	-3	TYR	-	expression tag	UNP P32322
С	-2	PHE	-	expression tag	UNP P32322
С	-1	GLN	-	expression tag	UNP P32322
С	0	SER	-	expression tag	UNP P32322
D	-21	MET	-	initiating methionine	UNP P32322
D	-20	HIS	-	expression tag	UNP P32322
D	-19	HIS	-	expression tag	UNP P32322
D	-18	HIS	-	expression tag	UNP P32322
D	-17	HIS	-	expression tag	UNP P32322
D	-16	HIS	-	expression tag	UNP P32322
D	-15	HIS	-	expression tag	UNP P32322
D	-14	SER	-	expression tag	UNP P32322
D	-13	SER	-	expression tag	UNP P32322
D	-12	GLY	-	expression tag	UNP P32322
D	-11	VAL	-	expression tag	UNP P32322
D	-10	ASP	-	expression tag	UNP P32322
D	-9	LEU	-	expression tag	UNP P32322
D	-8	GLY	-	expression tag	UNP P32322
D	-7	THR	-	expression tag	UNP P32322
D	-6	GLU	-	expression tag	UNP P32322
D	-5	ASN	-	expression tag	UNP P32322
D	-4	LEU	-	expression tag	UNP P32322
D	-3	TYR	-	expression tag	UNP P32322
D	-2	PHE	-	expression tag	UNP P32322
D	-1	GLN	-	expression tag	UNP P32322
D	0	SER	-	expression tag	UNP P32322
Е	-21	MET	-	initiating methionine	UNP P32322
Е	-20	HIS	-	expression tag	UNP P32322
Е	-19	HIS	-	expression tag	UNP P32322
Е	-18	HIS	-	expression tag	UNP P32322
Е	-17	HIS	-	expression tag	UNP P32322
Е	-16	HIS	-	expression tag	UNP P32322
Е	-15	HIS	-	expression tag	UNP P32322
Е	-14	SER	-	expression tag	UNP P32322
Е	-13	SER		expression tag	UNP P32322
Е	-12	GLY		expression tag	UNP P32322
Е	-11	VAL		expression tag	UNP P32322
Е	-10	ASP	-	expression tag	UNP P32322
Е	-9	LEU	-	expression tag	UNP P32322
Е	-8	GLY	-	expression tag	UNP P32322
Е	-7	THR	-	expression tag	UNP P32322



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-6	GLU	-	expression tag	UNP P32322
Е	-5	ASN	-	expression tag	UNP P32322
E	-4	LEU	-	expression tag	UNP P32322
E	-3	TYR	-	expression tag	UNP P32322
E	-2	PHE	-	expression tag	UNP P32322
Е	-1	GLN	-	expression tag	UNP P32322
Е	0	SER	-	expression tag	UNP P32322


• Molecule 2 is (2S)-2-hydroxy-3-methylbutanoic acid (CCD ID: LHV) (formula: $C_5H_{10}O_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 8 5 3	0	0
2	В	1	Total C O 8 5 3	0	0
2	С	1	Total C O 8 5 3	0	0
2	D	1	Total C O 8 5 3	0	0
2	E	1	Total C O 8 5 3	0	0

• Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O_4S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	С	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	Е	1	Total O S 5 4 1	0	0





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	D	1	Total 7	C 4	O 3	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	191	Total O 191 191	0	0
5	В	139	Total O 139 139	0	0
5	С	138	Total O 138 138	0	0
5	D	128	Total O 128 128	0	0
5	Е	146	Total O 146 146	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial





• Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	109.85Å 179.35Å 87.99Å	Donositor
a, b, c, α , β , γ	90.00° 106.67° 90.00°	Depositor
Resolution (Å)	90.76 - 1.61	Depositor
Resolution (A)	90.76 - 1.61	EDS
% Data completeness	98.0 (90.76-1.61)	Depositor
(in resolution range)	98.0 (90.76-1.61)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.68 (at 1.61Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
D D.	0.173 , 0.194	Depositor
R, R_{free}	0.173 , 0.194	DCC
R_{free} test set	10473 reflections $(5.09%)$	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	0.324	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 39.2	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10783	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, LHV, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.30	0/2097	0.50	0/2844	
1	В	0.30	0/2068	0.49	0/2810	
1	С	0.27	0/2025	0.46	0/2754	
1	D	0.26	0/1993	0.44	0/2716	
1	Е	0.29	0/1984	0.46	0/2696	
All	All	0.28	0/10167	0.47	0/13820	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2053	0	2079	9	0
1	В	2026	0	2036	11	0
1	С	1986	0	1973	8	0
1	D	1956	0	1914	7	0
1	Е	1948	0	1952	10	0
2	A	8	0	0	0	0
2	В	8	0	0	0	0
2	С	8	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	8	0	0	0	0
2	Е	8	0	0	0	0
3	A	5	0	0	0	0
3	В	5	0	0	1	0
3	С	5	0	0	0	0
3	D	5	0	0	0	0
3	Е	5	0	0	0	0
4	D	7	0	10	1	0
5	A	191	0	0	0	0
5	В	139	0	0	0	0
5	С	138	0	0	0	0
5	D	128	0	0	0	0
5	Е	146	0	0	1	0
All	All	10783	0	9964	44	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance } (\text{\AA}) \end{array}$	Clash overlap (Å)
1:A:133:THR:HG21	1:A:153:LEU:HD13	1.71	0.70
1:E:118:ILE:HD13	1:E:149[B]:MET:HG2	1.80	0.63
1:C:210:LEU:HD13	1:D:210:LEU:HD11	1.82	0.60
1:B:268:LEU:HA	1:B:271:MET:HE2	1.85	0.58
1:B:-4:LEU:HD21	1:B:151:GLN:HG2	1.86	0.58
1:A:93:VAL:HG21	1:A:149[B]:MET:HE2	1.88	0.55
1:C:93:VAL:HG21	1:C:149[A]:MET:HE1	1.88	0.55
1:C:133:THR:HG21	1:C:153:LEU:HD13	1.90	0.54
1:E:93:VAL:HG21	1:E:149[A]:MET:HE1	1.90	0.54
1:A:93:VAL:HG21	1:A:149[A]:MET:HE1	1.90	0.53
1:E:87:GLU:OE1	1:E:89:ARG:NH1	2.44	0.50
1:B:54:PRO:HG2	1:B:55:HIS:CE1	2.47	0.50
1:B:74:ILE:HG22	1:B:78:ILE:HG12	1.93	0.49
1:A:1[A]:MET:HE1	1:A:148:LEU:HD11	1.96	0.48
1:A:118:ILE:HD12	1:A:149[A]:MET:HE3	1.96	0.48
1:E:8:ALA:HB3	1:E:34:SER:HB2	1.96	0.47
1:E:267:GLU:OE2	5:E:501:HOH:O	2.20	0.47
1:D:226:GLN:HE22	4:D:402:PEG:H31	1.79	0.46
1:C:8:ALA:HB3	1:C:34:SER:HB2	1.97	0.46
1:C:33:SER:HA	1:C:53:THR:O	2.14	0.46



Continued from previous page...

A 1	A. 0	Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)	
1:B:6:ILE:HG13	1:B:66:LEU:HD11	1.98	0.45	
1:D:48:MET:HE2	1:D:48:MET:HB3	1.90	0.45	
1:A:93:VAL:CG2	1:A:149[A]:MET:HE1	2.46	0.45	
1:E:185:LEU:HD21	1:E:210:LEU:HD22	1.99	0.45	
1:C:114[A]:ALA:HA	1:C:140:HIS:CG	2.52	0.45	
1:B:222:GLN:NE2	3:B:402:SO4:O2	2.47	0.44	
1:D:33:SER:HA	1:D:53:THR:O	2.17	0.44	
1:E:93:VAL:CG2	1:E:149[A]:MET:HE1	2.47	0.44	
1:E:8:ALA:O	1:E:41:THR:OG1	2.29	0.44	
1:C:2:SER:HB2	1:C:64:ASP:OD2	2.18	0.43	
1:B:78:ILE:O	1:B:82:ILE:HG13	2.19	0.43	
1:D:75:ILE:HG21	1:D:99:VAL:HG11	2.01	0.43	
1:D:8:ALA:HB3	1:D:34:SER:HB2	1.99	0.43	
1:D:78:ILE:O	1:D:82:ILE:HG13	2.19	0.43	
1:C:36:ASP:C	1:C:38:ASP:H	2.26	0.43	
1:A:4:GLY:HA3	1:A:63[A]:SER:OG	2.19	0.42	
1:B:133:THR:O	1:B:159:CYS:HA	2.20	0.42	
1:A:65:VAL:HG11	1:A:149[B]:MET:HE1	2.02	0.41	
1:B:123:ASN:CG	1:B:125:PRO:HD2	2.45	0.41	
1:E:118:ILE:CD1	1:E:149[B]:MET:HG2	2.49	0.41	
1:A:74:ILE:HG22	1:A:78:ILE:HG12	2.03	0.41	
1:B:8:ALA:HB3	1:B:34:SER:HB2	2.02	0.41	
1:B:33:SER:HA	1:B:53:THR:O	2.21	0.40	
1:E:4:GLY:HA3	1:E:63:SER:OG	2.21	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	284/316 (90%)	279 (98%)	5 (2%)	0	100 100



Continued	trom	mromonie	maaa
-	110111	DICULUUS	pauc

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	280/316 (89%)	274 (98%)	6 (2%)	0	100	100
1	\mathbf{C}	278/316 (88%)	272 (98%)	6 (2%)	0	100	100
1	D	277/316 (88%)	273 (99%)	4 (1%)	0	100	100
1	E	274/316 (87%)	269 (98%)	5 (2%)	0	100	100
All	All	1393/1580 (88%)	1367 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	211/251 (84%)	205 (97%)	6 (3%)	38	14
1	В	207/251 (82%)	207 (100%)	0	100	100
1	С	198/251 (79%)	194 (98%)	4 (2%)	50	25
1	D	191/251 (76%)	190 (100%)	1 (0%)	86	78
1	E	195/251 (78%)	194 (100%)	1 (0%)	86	78
All	All	1002/1255 (80%)	990 (99%)	12 (1%)	67	49

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-4	LEU
1	A	142	GLN
1	A	210	LEU
1	A	222	GLN
1	A	232	SER
1	A	271	MET
1	С	2	SER
1	С	29	LYS
1	С	87	GLU
1	С	120	CYS



Continued from previous page...

Mol	Chain	Res	Type
1	D	232	SER
1	Е	62	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	GLN
1	A	226	GLN
1	В	140	HIS
1	В	219	HIS
1	С	73	HIS
1	С	140	HIS
1	С	226	GLN
1	D	226	GLN
1	Ε	140	HIS
1	Ε	226	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Trino	Chain	Dag	T inle	В	ond leng	gths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	С	402	-	4,4,4	0.23	0	6,6,6	0.10	0
2	LHV	D	401	-	7,7,7	1.45	1 (14%)	9,9,9	1.08	0
2	LHV	A	401	-	7,7,7	1.21	1 (14%)	9,9,9	1.20	1 (11%)
2	LHV	В	401	-	7,7,7	1.22	1 (14%)	9,9,9	1.56	3 (33%)
3	SO4	Е	402	-	4,4,4	0.24	0	6,6,6	0.19	0
3	SO4	A	402	-	4,4,4	0.27	0	6,6,6	0.07	0
2	LHV	E	401	-	7,7,7	1.31	1 (14%)	9,9,9	1.25	0
2	LHV	С	401	-	7,7,7	0.92	0	9,9,9	0.92	0
3	SO4	В	402	-	4,4,4	0.31	0	6,6,6	0.11	0
4	PEG	D	402	-	6,6,6	0.18	0	5,5,5	0.03	0
3	SO4	D	403	-	4,4,4	0.24	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LHV	D	401	-	-	2/8/8/8	_
2	LHV	A	401	-	-	4/8/8/8	-
2	LHV	В	401	-	-	0/8/8/8	-
2	LHV	E	401	ı	-	2/8/8/8	-
2	LHV	С	401	-	-	4/8/8/8	-
4	PEG	D	402	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
2	D	401	LHV	CA-C	2.81	1.56	1.52
2	Ε	401	LHV	OE-CA	-2.65	1.37	1.42
2	В	401	LHV	CA-C	2.31	1.56	1.52
2	A	401	LHV	CA-C	-2.07	1.49	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	В	401	LHV	O-C-CA	-2.52	114.91	121.62
2	В	401	LHV	OXT-C-CA	2.51	120.28	113.31
2	A	401	LHV	CB-CA-C	-2.22	107.61	111.60
2	В	401	LHV	OE-CA-CB	-2.08	105.20	110.15



There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	LHV	C-CA-CB-CG1
2	A	401	LHV	OE-CA-CB-CG1
2	A	401	LHV	C-CA-CB-CG2
2	A	401	LHV	OE-CA-CB-CG2
2	С	401	LHV	C-CA-CB-CG1
2	С	401	LHV	OE-CA-CB-CG1
2	С	401	LHV	C-CA-CB-CG2
2	С	401	LHV	OE-CA-CB-CG2
2	D	401	LHV	C-CA-CB-CG1
4	D	402	PEG	O2-C3-C4-O4
2	D	401	LHV	OE-CA-CB-CG1
4	D	402	PEG	C4-C3-O2-C2
2	Е	401	LHV	O-C-CA-CB
2	Е	401	LHV	OXT-C-CA-CB


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	402	SO4	1	0
4	D	402	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	282/316 (89%)	0.02	10 (3%) 47 49	15, 30, 52, 89	4 (1%)
1	В	279/316 (88%)	0.24	12 (4%) 40 41	15, 33, 61, 81	3 (1%)
1	С	275/316 (87%)	0.24	13 (4%) 37 39	16, 35, 65, 83	5 (1%)
1	D	277/316 (87%)	0.48	23 (8%) 19 18	15, 40, 71, 86	2 (0%)
1	E	274/316 (86%)	0.37	15 (5%) 32 33	15, 38, 69, 87	2 (0%)
All	All	1387/1580 (87%)	0.27	73 (5%) 33 34	15, 34, 67, 89	16 (1%)

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-3	TYR	5.1
1	D	25	LEU	4.2
1	D	74	ILE	4.1
1	С	0	SER	4.0
1	Е	74	ILE	3.8
1	A	-4	LEU	3.7
1	A	274	GLN	3.5
1	D	270	SER	3.4
1	Е	0	SER	3.3
1	D	35	PRO	3.3
1	Е	37	MET	3.2
1	A	-6	GLU	3.1
1	A	268	LEU	3.0
1	В	42	VAL	3.0
1	С	42	VAL	3.0
1	D	39	LEU	2.9
1	В	40	ALA	2.9
1	A	-2	PHE	2.9
1	D	45	LEU	2.9
1	В	43	SER	2.8



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	Е	143	VAL	2.8
1	В	-4	LEU	2.8
1	В	143	VAL	2.8
1	D	44	ALA	2.8
1	С	274	GLN	2.8
1	D	24	VAL	2.7
1	В	273	ASP	2.7
1	D	28	HIS	2.7
1	Е	42	VAL	2.7
1	D	13	PHE	2.6
1	D	8	ALA	2.6
1	A	273	ASP	2.6
1	Е	77	PHE	2.5
1	D	-4	LEU	2.5
1	C	24	VAL	2.5
1	D	42	VAL	2.5
1	D	37	MET	2.5
1	В	38	ASP	2.5
1	С	38	ASP	2.5
1	В	268	LEU	2.4
1	D	77	PHE	2.4
1	Е	40	ALA	2.4
1	D	52	LEU	2.4
1	С	36	ASP	2.4
1	С	37	MET	2.4
1	Е	111	PHE	2.3
1	В	72	PRO	2.3
1	D	271	MET	2.3
1	В	74	ILE	2.3
1	A	272	ALA	2.3
1	С	39	LEU	2.3
1	С	143	VAL	2.2
1	В	37	MET	2.2
1	D	38	ASP	2.2
1	D	-1	GLN	2.2
1	Е	69	ALA	2.2
1	С	1	MET	2.2
1	D	72	PRO	2.2
1	Е	35	PRO	2.2
1	Е	72	PRO	2.2
1	D	26	ALA	2.1
1	D	82	ILE	2.1



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	Е	88	ASP	2.1
1	С	41	THR	2.1
1	Е	270	SER	2.1
1	В	272	ALA	2.1
1	С	54	PRO	2.1
1	Е	23	GLY	2.0
1	Е	39	LEU	2.0
1	A	143	VAL	2.0
1	A	231	VAL	2.0
1	С	40	ALA	2.0
1	D	0	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ extbf{B-factors}(extbf{A}^2)$	Q < 0.9
3	SO4	С	402	5/5	0.72	0.11	64,66,71,74	0
4	PEG	D	402	7/7	0.77	0.16	42,44,50,51	0
3	SO4	Е	402	5/5	0.78	0.15	50,59,60,63	5
3	SO4	В	402	5/5	0.79	0.14	43,44,50,50	5
3	SO4	D	403	5/5	0.82	0.12	65,66,67,68	5
3	SO4	A	402	5/5	0.88	0.12	48,50,51,52	5
2	LHV	С	401	8/8	0.88	0.14	30,41,46,46	0
2	LHV	Е	401	8/8	0.89	0.15	28,37,41,42	0
2	LHV	A	401	8/8	0.91	0.12	29,40,42,46	0
2	LHV	D	401	8/8	0.91	0.12	29,38,41,42	0
2	LHV	В	401	8/8	0.91	0.12	29,40,42,45	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















Electron density around LHV D 401:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

