



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 19, 2024 – 01:18 PM EDT

PDB ID : 8TSM  
Title : Crystal structure of chicken Netrin-1 LN LE1-2  
Authors : Heide, F.; Rafiei, F.; Stetefeld, J.  
Deposited on : 2023-08-11  
Resolution : 3.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1002 (4.14-3.66)
Clashscore	141614	1004 (4.12-3.68)
Ramachandran outliers	138981	1021 (4.14-3.66)
Sidechain outliers	138945	1014 (4.14-3.66)
RSRZ outliers	127900	1275 (4.20-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	502	-	-	-	X

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5539 atoms, of which 2693 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Netrin-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	354	5485	1743	2667	522	523	30	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	ALA	-	expression tag	UNP Q90922
A	23	PRO	-	expression tag	UNP Q90922
A	24	LEU	-	expression tag	UNP Q90922
A	25	ALA	-	expression tag	UNP Q90922
A	406	GLY	-	expression tag	UNP Q90922
A	407	SER	-	expression tag	UNP Q90922
A	408	SER	-	expression tag	UNP Q90922
A	409	GLY	-	expression tag	UNP Q90922
A	410	GLY	-	expression tag	UNP Q90922
A	411	GLY	-	expression tag	UNP Q90922
A	412	LEU	-	expression tag	UNP Q90922
A	413	VAL	-	expression tag	UNP Q90922
A	414	PRO	-	expression tag	UNP Q90922

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	H	N			O
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Netrin-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.41Å 112.41Å 242.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.77 – 3.90 37.77 – 3.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (37.77-3.90) 75.8 (37.77-3.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.61 (at 3.87Å)	Xtrriage
Refinement program	PHENIX 1.20.1-4487_4487	Depositor
R, $R_{free}$	0.312 , 0.319 0.315 , 0.321	Depositor DCC
$R_{free}$ test set	571 reflections (7.73%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	118.5	Xtrriage
Anisotropy	0.797	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 111.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	5539	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	219.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2892	0.62	0/3912

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	103	ARG	Sidechain
1	A	157	ARG	Sidechain
1	A	255	ARG	Sidechain
1	A	298	ARG	Sidechain
1	A	300	ARG	Sidechain
1	A	327	ARG	Sidechain
1	A	335	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2818	2667	2666	77	1
2	A	28	26	26	1	0
All	All	2846	2693	2692	78	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:ARG:NH1	1:A:209:HIS:HB3	1.87	0.90
1:A:55:PRO:HG2	1:A:287:CYS:HB3	1.67	0.76
1:A:256:LEU:HD23	1:A:257:HIS:H	1.52	0.74
1:A:133:ASN:HB2	1:A:135:THR:HG23	1.74	0.69
1:A:313:ALA:HB2	1:A:321:LYS:HA	1.74	0.68
1:A:327:ARG:HG2	1:A:328:PRO:HD2	1.75	0.68
1:A:190:PRO:HG2	1:A:193:ALA:HB2	1.77	0.67
1:A:330:GLN:HB2	1:A:338:ASN:HB3	1.76	0.67
1:A:166:SER:O	1:A:247:THR:OG1	2.09	0.66
1:A:223:SER:HB3	1:A:226:ASP:HB2	1.78	0.65
1:A:300:ARG:HH11	1:A:300:ARG:HG3	1.62	0.64
1:A:192:ARG:N	1:A:205:CYS:O	2.30	0.61
1:A:300:ARG:NH1	1:A:301:ASP:OD1	2.34	0.60
1:A:84:GLU:HG2	1:A:89:GLN:HG2	1.84	0.60
1:A:324:HIS:HA	1:A:342:ALA:HA	1.83	0.59
1:A:321:LYS:HG2	1:A:322:PRO:HD2	1.83	0.58
1:A:157:ARG:CZ	1:A:209:HIS:HB3	2.33	0.58
1:A:290:ASN:ND2	1:A:316:GLU:O	2.34	0.57
1:A:321:LYS:CG	1:A:322:PRO:HD2	2.33	0.57
1:A:72:CYS:HB3	1:A:125:ASP:HA	1.87	0.57
1:A:256:LEU:CD2	1:A:257:HIS:H	2.17	0.56
1:A:114:ASN:HB2	1:A:280:ASP:HB3	1.86	0.56
1:A:357:GLU:HB3	1:A:361:LEU:HD12	1.86	0.56
1:A:310:HIS:HB3	1:A:339:GLU:HB2	1.87	0.55
1:A:193:ALA:HB3	1:A:204:ILE:HG22	1.88	0.55
1:A:157:ARG:NH1	1:A:219:LEU:O	2.41	0.54
1:A:330:GLN:HB2	1:A:338:ASN:CB	2.36	0.54
1:A:289:CYS:SG	1:A:293:ALA:HB3	2.48	0.54
1:A:327:ARG:NH2	1:A:339:GLU:O	2.41	0.54
1:A:157:ARG:HD2	1:A:209:HIS:O	2.08	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:317:CYS:O	1:A:329:TRP:NE1	2.44	0.51
1:A:206:THR:OG1	1:A:207:ASP:N	2.43	0.51
1:A:122:TRP:CH2	1:A:124:SER:HB3	2.46	0.51
1:A:98:ALA:HA	1:A:104:ALA:HB2	1.94	0.50
1:A:143:LYS:HD2	1:A:316:GLU:HG2	1.92	0.50
1:A:60:SER:OG	1:A:283:VAL:O	2.27	0.50
1:A:234:PHE:O	1:A:241:GLN:NE2	2.38	0.50
1:A:134:VAL:HB	1:A:253:PHE:HB2	1.94	0.50
1:A:166:SER:OG	1:A:168:ASP:OD1	2.30	0.50
1:A:175:PRO:HB3	1:A:178:PHE:CZ	2.47	0.49
1:A:127:TYR:CE2	1:A:272:ASP:O	2.65	0.49
1:A:328:PRO:HD3	1:A:354:PHE:CZ	2.47	0.49
1:A:76:PRO:HA	1:A:96:CYS:O	2.13	0.48
1:A:195:ILE:HG21	1:A:228:ARG:HD3	1.94	0.48
1:A:178:PHE:HB2	1:A:205:CYS:HA	1.96	0.48
1:A:133:ASN:N	1:A:133:ASN:OD1	2.46	0.48
1:A:256:LEU:HD23	1:A:257:HIS:N	2.26	0.47
1:A:114:ASN:CB	1:A:280:ASP:HB3	2.44	0.47
1:A:122:TRP:O	1:A:277:ALA:HA	2.15	0.47
1:A:166:SER:HB3	1:A:248:ASP:HB2	1.97	0.47
1:A:182:GLN:HG3	1:A:182:GLN:O	2.16	0.46
1:A:311:ASN:HB3	1:A:324:HIS:CD2	2.50	0.46
1:A:377:THR:HG22	1:A:403:CYS:SG	2.56	0.46
1:A:55:PRO:O	1:A:286:ARG:NH1	2.48	0.46
1:A:153:PHE:HB3	1:A:155:SER:O	2.16	0.46
1:A:312:THR:HG22	1:A:340:CYS:SG	2.56	0.45
1:A:105:HIS:HB3	1:A:123:GLN:N	2.32	0.45
1:A:176:PHE:CZ	1:A:224:THR:HG22	2.52	0.45
1:A:156:PRO:HB3	1:A:212:VAL:HG13	1.99	0.45
1:A:207:ASP:O	1:A:210:THR:OG1	2.33	0.45
1:A:45:ASP:OD2	1:A:49:LEU:HB2	2.17	0.44
1:A:105:HIS:CG	1:A:123:GLN:HB3	2.53	0.44
1:A:313:ALA:HB2	1:A:321:LYS:CA	2.47	0.44
1:A:130:TYR:CE2	1:A:131:PRO:HB3	2.53	0.44
1:A:212:VAL:HG22	1:A:257:HIS:CG	2.52	0.43
2:A:502:NAG:O7	2:A:502:NAG:H3	2.17	0.43
1:A:292:HIS:HB2	1:A:317:CYS:SG	2.58	0.43
1:A:106:PRO:HD2	1:A:109:PHE:CE2	2.54	0.42
1:A:318:ASP:OD1	1:A:318:ASP:N	2.52	0.42
1:A:53:CYS:HB2	1:A:294:SER:HB3	2.02	0.42
1:A:120:THR:O	1:A:279:SER:HA	2.20	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:THR:HA	1:A:251:VAL:O	2.20	0.41
1:A:321:LYS:CG	1:A:322:PRO:CD	2.97	0.41
1:A:181:THR:HA	1:A:207:ASP:CB	2.51	0.41
1:A:177:GLN:OE1	1:A:204:ILE:HG13	2.21	0.40
1:A:300:ARG:HA	1:A:300:ARG:HD2	1.87	0.40
1:A:321:LYS:HG3	1:A:322:PRO:HD2	2.02	0.40
1:A:291:GLY:HA3	1:A:332:ALA:HB3	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:SER:OG	1:A:211:ASP:OD2[8_555]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	350/393 (89%)	295 (84%)	52 (15%)	3 (1%)	17 54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	133	ASN
1	A	302	ASP
1	A	55	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	316/343 (92%)	314 (99%)	2 (1%)	86 91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	ASN
1	A	165	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	324	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	501	1	14,14,15	0.66	1 (7%)	17,19,21	0.68	0
2	NAG	A	502	1	14,14,15	0.68	0	17,19,21	1.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	501	1	-	1/6/23/26	0/1/1/1
2	NAG	A	502	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NAG	C1-C2	2.30	1.55	1.52

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	502	NAG	C3-C2-N2-C7
2	A	502	NAG	O5-C5-C6-O6
2	A	501	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	502	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	354/393 (90%)	-0.14	9 (2%) 57 47	123, 215, 260, 283	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	275	PHE	3.7
1	A	259	PHE	3.6
1	A	368	GLY	2.7
1	A	142	LYS	2.7
1	A	258	THR	2.5
1	A	162	ALA	2.4
1	A	351	ARG	2.3
1	A	141	GLY	2.2
1	A	143	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	502	14/15	0.50	0.42	37,37,88,88	27
2	NAG	A	501	14/15	0.72	0.31	37,37,100,100	27

## 6.5 Other polymers [i](#)

There are no such residues in this entry.