



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 29, 2023 – 02:04 AM EST

PDB ID : 8UG3
Title : Crystal structure of KHK-C and compound 23
Authors : Durbin, J.D.; Guo, S.Y.
Deposited on : 2023-10-05
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ketohexokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	2232	1398	398	424	12	0	1	0
1	B	304	2336	1456	421	446	13	0	5	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	MET	-	expression tag	UNP P50053
A	-15	SER	-	expression tag	UNP P50053
A	-14	LEU	-	expression tag	UNP P50053
A	-13	GLY	-	expression tag	UNP P50053
A	-12	SER	-	expression tag	UNP P50053
A	-11	SER	-	expression tag	UNP P50053
A	-10	HIS	-	expression tag	UNP P50053
A	-9	HIS	-	expression tag	UNP P50053
A	-8	HIS	-	expression tag	UNP P50053
A	-7	HIS	-	expression tag	UNP P50053
A	-6	HIS	-	expression tag	UNP P50053
A	-5	HIS	-	expression tag	UNP P50053
A	-4	SER	-	expression tag	UNP P50053
A	-3	SER	-	expression tag	UNP P50053
A	-2	GLY	-	expression tag	UNP P50053
A	-1	LEU	-	expression tag	UNP P50053
A	0	VAL	-	expression tag	UNP P50053
A	1	PRO	-	expression tag	UNP P50053
A	2	ARG	-	expression tag	UNP P50053
A	3	GLY	-	expression tag	UNP P50053
A	4	SER	-	expression tag	UNP P50053
B	-16	MET	-	expression tag	UNP P50053
B	-15	SER	-	expression tag	UNP P50053
B	-14	LEU	-	expression tag	UNP P50053
B	-13	GLY	-	expression tag	UNP P50053

Continued on next page...

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	113	Total O 113 113	0	0
5	B	191	Total O 191 191	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ketohexokinase



- Molecule 1: Ketohexokinase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.21Å 85.80Å 135.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.38 – 2.02 39.35 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.38-2.02) 99.9 (39.35-2.02)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 2.03Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.179 , 0.216 0.183 , 0.218	Depositor DCC
R_{free} test set	3172 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.2	Xtrriage
Anisotropy	0.079	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.025 for k,h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4969	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: WRE, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/2275	1.03	4/3088 (0.1%)
1	B	0.71	1/2381 (0.0%)	1.09	6/3227 (0.2%)
All	All	0.65	1/4656 (0.0%)	1.06	10/6315 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	228	GLU	CD-OE1	5.73	1.31	1.25

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	77	ARG	NE-CZ-NH2	-8.60	116.00	120.30
1	A	279	ARG	NE-CZ-NH2	-7.20	116.70	120.30
1	A	227	GLU	CB-CA-C	-5.83	98.73	110.40
1	A	93	ASP	CB-CA-C	-5.83	98.74	110.40
1	B	272[A]	ARG	NE-CZ-NH1	-5.43	117.58	120.30
1	B	272[B]	ARG	NE-CZ-NH1	-5.43	117.58	120.30
1	A	152	ARG	NE-CZ-NH1	5.37	122.99	120.30
1	B	85	GLN	CB-CA-C	-5.31	99.78	110.40
1	B	79	ARG	NE-CZ-NH2	-5.05	117.78	120.30
1	B	180	PHE	CB-CA-C	5.03	120.46	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2232	0	2153	34	0
1	B	2336	0	2266	18	0
2	A	29	0	0	0	0
2	B	21	0	0	0	0
3	A	15	0	0	0	0
3	B	20	0	0	1	0
4	B	12	0	16	0	0
5	A	113	0	0	3	0
5	B	191	0	0	2	0
All	All	4969	0	4435	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:ASP:HB2	1:B:31:ARG:HD2	1.74	0.69
1:A:31:ARG:NH2	3:B:302:SO4:O3	2.26	0.67
1:A:146:GLN:HA	1:A:149:MET:HG3	1.74	0.67
1:B:202:GLN:HE21	1:B:202:GLN:HA	1.62	0.64
1:A:193:LYS:HE3	1:A:225:TRP:NE1	2.13	0.64
1:A:183:PHE:HD1	1:A:189:VAL:HG11	1.66	0.61
1:B:202:GLN:HA	1:B:202:GLN:NE2	2.16	0.61
1:A:123:THR:O	1:A:126:GLU:HB2	2.01	0.60
1:A:114:ASP:HB2	1:B:31:ARG:CD	2.35	0.56
1:A:150:LEU:HA	1:A:153:ILE:HD12	1.88	0.56
1:B:-1:LEU:C	1:B:-1:LEU:HD23	2.26	0.55
1:A:93:ASP:HB2	1:A:115:THR:HG21	1.89	0.55
1:A:193:LYS:HE3	1:A:225:TRP:CE2	2.42	0.54
1:A:42:ASN:HB2	1:A:139:GLU:CD	2.28	0.54
1:A:125:PHE:CD2	1:A:149:MET:HB3	2.42	0.53
1:B:202:GLN:HE21	1:B:202:GLN:CA	2.22	0.51
1:A:193:LYS:HE2	1:A:197:LYS:HE2	1.91	0.51
1:A:235:PRO:HD2	5:A:462:HOH:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:202:GLN:NE2	1:B:202:GLN:CA	2.75	0.50
1:A:156:HIS:O	1:A:160:GLN:HG2	2.12	0.50
1:A:116:SER:O	1:A:117:LEU:C	2.51	0.49
1:A:205[B]:GLU:CG	5:A:441:HOH:O	2.60	0.49
1:B:123:THR:HG23	5:B:482:HOH:O	2.12	0.49
1:A:93:ASP:HB2	1:A:115:THR:CG2	2.43	0.48
1:B:2:ARG:NH1	1:B:164:GLN:HG2	2.29	0.47
1:B:99[B]:CYS:SG	1:B:108:ARG:NH2	2.87	0.47
1:A:31:ARG:HD3	1:B:115:THR:HG23	1.97	0.47
1:B:-3:SER:HB3	5:B:418:HOH:O	2.14	0.47
1:A:89:GLN:HG3	1:A:118:PRO:HD2	1.98	0.46
1:A:39:ARG:HG3	1:A:72:LEU:HD22	1.97	0.46
1:B:102:ASN:HB3	1:B:105:ASN:ND2	2.31	0.46
1:A:25:LYS:O	1:A:28:SER:OG	2.29	0.45
1:A:205[B]:GLU:HG3	5:A:441:HOH:O	2.17	0.45
1:A:150:LEU:HD13	1:A:185:TYR:HB2	2.00	0.43
1:A:180:PHE:CE2	1:A:199:LEU:HD21	2.53	0.43
1:A:263:SER:HB2	1:A:281:GLY:HA2	2.00	0.43
1:A:108:ARG:O	1:B:27:ASP:HA	2.19	0.43
1:A:116:SER:O	1:A:117:LEU:O	2.37	0.42
1:B:156:HIS:O	1:B:160:GLN:HG2	2.20	0.42
1:A:184:GLY:O	1:A:216:ARG:HB3	2.19	0.42
1:A:201:PHE:CZ	1:A:210:GLY:HA3	2.55	0.42
1:B:18:SER:OG	1:B:35[B]:GLN:HG3	2.19	0.42
1:A:180:PHE:N	1:A:180:PHE:CD1	2.88	0.41
1:B:19:LEU:O	1:B:33:LEU:N	2.48	0.41
1:A:161:PRO:HG2	1:A:164:GLN:HG3	2.02	0.41
1:A:180:PHE:HE2	1:A:199:LEU:HD21	1.86	0.41
1:A:9:VAL:HA	1:A:60:MET:O	2.21	0.40
1:B:154:ASP:O	1:B:158:THR:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/315 (94%)	284 (96%)	12 (4%)	1 (0%)	41	36
1	B	307/315 (98%)	298 (97%)	8 (3%)	1 (0%)	41	36
All	All	604/630 (96%)	582 (96%)	20 (3%)	2 (0%)	41	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	-3	SER
1	A	117	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/262 (89%)	227 (97%)	7 (3%)	41	40
1	B	251/262 (96%)	246 (98%)	5 (2%)	55	57
All	All	485/524 (93%)	473 (98%)	12 (2%)	47	48

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ASP
1	A	51	SER
1	A	108	ARG
1	A	114	ASP
1	A	162	PRO
1	A	231	ASP
1	A	298	VAL
1	B	21	ASP
1	B	90	SER
1	B	105	ASN
1	B	177	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	298	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	105	ASN
1	B	202	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	WRE	A	301	-	30,32,32	1.18	3 (10%)	32,47,47	1.00	2 (6%)
4	GOL	B	307	-	5,5,5	0.31	0	5,5,5	0.66	0
4	GOL	B	306	-	5,5,5	0.19	0	5,5,5	0.40	0
3	SO4	B	302	-	4,4,4	0.42	0	6,6,6	0.27	0
3	SO4	B	305	-	4,4,4	0.19	0	6,6,6	0.27	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	302	-	4,4,4	0.51	0	6,6,6	0.15	0
3	SO4	B	304	-	4,4,4	0.40	0	6,6,6	0.25	0
3	SO4	A	304	-	4,4,4	0.36	0	6,6,6	0.65	0
3	SO4	B	303	-	4,4,4	0.40	0	6,6,6	0.27	0
3	SO4	A	303	-	4,4,4	0.35	0	6,6,6	0.20	0
2	WRE	B	301	-	20,23,32	1.03	2 (10%)	21,35,47	1.08	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	WRE	A	301	-	-	2/17/39/39	0/4/4/4
4	GOL	B	307	-	-	0/4/4/4	-
4	GOL	B	306	-	-	2/4/4/4	-
2	WRE	B	301	-	-	0/10/23/39	0/3/3/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	WRE	F2-C18	-4.30	1.17	1.32
2	A	301	WRE	F3-C18	-2.86	1.22	1.32
2	B	301	WRE	F2-C18	-2.61	1.23	1.32
2	A	301	WRE	C7-N6	2.53	1.41	1.36
2	B	301	WRE	C7-N6	2.10	1.40	1.36

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	WRE	C16-C15-C9	-3.00	111.04	117.39
2	A	301	WRE	C12-C14-N7	2.16	114.21	109.45
2	B	301	WRE	F3-C18-C6	2.06	116.00	112.47
2	A	301	WRE	C17-N4-C3	-2.03	126.61	129.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	WRE	N2-C7-N6-C10

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	B	306	GOL	O1-C1-C2-O2
4	B	306	GOL	O1-C1-C2-C3
2	A	301	WRE	N3-C7-N6-C10


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/315 (94%)	0.43	20 (6%) 17 17	39, 62, 95, 121	0
1	B	304/315 (96%)	0.12	17 (5%) 24 23	36, 49, 81, 114	0
All	All	602/630 (95%)	0.27	37 (6%) 21 20	36, 55, 91, 121	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	ILE	4.5
1	B	-5	HIS	4.4
1	B	-4	SER	4.0
1	A	161	PRO	3.9
1	A	158	THR	3.5
1	A	162	PRO	3.5
1	B	23	TYR	3.4
1	A	164	GLN	3.3
1	A	1	PRO	3.3
1	B	58	ALA	3.2
1	A	160	GLN	3.1
1	A	2	ARG	3.1
1	A	198	HIS	2.9
1	A	92	GLY	2.8
1	B	105	ASN	2.8
1	B	107	ASN	2.7
1	B	7	LEU	2.7
1	A	123	THR	2.7
1	A	131	THR	2.7
1	A	91	LYS	2.7
1	A	159	ARG	2.5
1	A	218	GLY	2.4
1	B	104	SER	2.4
1	A	166	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	8	CYS	2.4
1	B	50	LEU	2.4
1	A	153	ILE	2.3
1	B	57	CYS	2.2
1	A	163	GLU	2.2
1	B	106	GLY	2.2
1	B	133	PHE	2.1
1	B	47	CYS	2.1
1	B	102	ASN	2.1
1	A	13	VAL	2.1
1	B	81	VAL	2.1
1	B	90	SER	2.0
1	A	282	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	304	5/5	0.67	0.32	77,78,136,136	0
3	SO4	A	303	5/5	0.80	0.18	65,73,83,91	5
3	SO4	B	303	5/5	0.84	0.16	70,73,78,80	5
4	GOL	B	307	6/6	0.86	0.15	59,74,85,94	0
3	SO4	B	304	5/5	0.87	0.37	59,63,69,74	5
3	SO4	A	302	5/5	0.92	0.16	55,56,67,69	5
3	SO4	B	305	5/5	0.93	0.13	61,79,94,104	0
4	GOL	B	306	6/6	0.94	0.08	65,94,99,99	0
2	WRE	B	301	21/29	0.94	0.11	38,47,58,65	0
3	SO4	B	302	5/5	0.96	0.23	49,51,56,57	5


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	WRE	A	301	29/29	0.97	0.12	43,55,92,107	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.