

Full wwPDB EM Validation Report (i)

Nov 16, 2024 – 06:53 AM EST

PDB ID : 8VWJ

EMDB ID : EMD-43589

Title: The base complex of the AcMNPV baculovirus nucleocapsid (Class 2, localised

reconstruction)

Authors : Johnstone, B.A.; Koszalka, P.; Ha, J.; Venugopal, H.; Coulibaly, F.

Deposited on : 2024-02-01

Resolution : 4.78 Å(reported)

Based on initial model : .

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113

MolProbity : 4.02b-467

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : 1.9.13

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 4.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality	of chain
1	A	347	29% 79%	21%
1	В	347	37%	• 20%
1	С	347	10%	· 18%
1	D	347	8%	· 18%
1	Е	347	80%	• 18%
1	F	347	82%	• 18%
1	V	347	26% 79%	21%
1	W	347	28%	20%
1	X	347	12%	• 18%



 $Continued\ from\ previous\ page...$

Mol	Chain	Length	Quality of chain										
1	Y	347	82%	18%									
1	Z	347	81%	• 18%									
1	a	347	12%	• 18%									
2	G	691	28% 72%										
2	Н	691	30% 70%										
2	I	691	32% • 68°	V ₆									
2	b	691	28% 71%										
2	c	691	30% 70%										
2	d	691	32% 68%										
3	J	290	6%	28%									
3	M	290	84%	16%									
3	Q	290	63%	36%									
3	S	290	76%	23%									
3	e	290	7%	28%									
3	h	290	•										
	K		83% 5%	• 16%									
4		361	64%	35%									
4	L	361	64%	36%									
4	R	361	6 0% ∙	39%									
4	T	361	61% 7%	39%									
4	f	361	65%	35%									
4	g	361	62%	37%									
5	N	390	93%	• 7%									
5	i	390	92%	• 7%									
6	О	477	98%										
6	j	477	98%										



 $Continued\ from\ previous\ page...$

	<i>J</i>	I	1 3	
\mathbf{Mol}	Chain	Length	Quality of chain	
	_		17%	
7	Р	320	95%	1
_	1	220	17%	
7	k	320	94%	,



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 156295 atoms, of which 77632 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Major capsid protein.

Mol	Chain	Residues			Atom	s			AltConf	Trace
1	A	274	Total	С	Н	N	О	S	0	0
1	A	214	4371	1396	2157	387	419	12	0	U
1	В	279	Total	С	Н	N	О	S	0	0
1	Б	219	4449	1417	2195	396	429	12	0	U
1	С	285	Total	С	Н	N	О	S	0	0
1		200	4564	1459	2252	406	435	12	0	U
1	D	284	Total	С	Н	N	О	S	0	0
1	D	204	4529	1442	2235	403	437	12		0
1	Е	283	Total	С	Н	N	О	S	0	0
1	12	200	4509	1443	2223	398	432	13		0
1	F	286	Total	С	Н	N	О	S	0	0
1	I.	280	4552	1457	2242	403	437	13	U	U
1	V	274	Total	С	Н	N	О	S	0	0
1	V	214	4371	1396	2157	387	419	12	0	U
1	W	279	Total	С	Н	N	Ο	S	0	0
1	VV	219	4449	1417	2195	396	429	12	0	U
1	X	285	Total	С	Η	N	Ο	S	0	0
1	Λ	200	4564	1459	2252	406	435	12	0	U
1	Y	285	Total	С	Η	N	Ο	S	0	0
1	1	200	4554	1455	2244	404	439	12	0	U
1	Z	283	Total	С	Η	N	O	S	0	0
1	L	200	4509	1443	2223	398	432	13	U	U
1	2	286	Total	С	Н	N	О	S	1 ()	0
	a	200	4552	1457	2242	403	437	13	0	U

• Molecule 2 is a protein called Capsid-associated protein VP80.

Mol	Chain	Residues				AltConf	Trace			
2	G	194	Total	С	Н	N	О	S	0	0
2	G	194	3225	1049	1609	265	290	12		U
2	Н	210	Total	С	Н	N	О	S	0	0
2	11	210	3513	1133	1760	293	314	13	0	U
2	Ţ	224	Total	С	Н	N	О	S	0	0
	1	224	3740	1209	1862	315	340	14	U	U



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues				AltConf	Trace			
2	h	197	Total	С	Н	N	О	S	0	0
	b	197	3286	1066	1644	269	295	12	0	U
2		206	Total	С	Н	N	О	S	0	0
	С	200	3449	1113	1730	289	304	13	0	U
9	d	224	Total	С	Н	N	О	S	0	0
2	a	ZZ4	3748	1209	1870	315	340	14	0	U

• Molecule 3 is a protein called Occlusion-derived virus envelope protein E27.

Mol	Chain	Residues			Atom	\mathbf{s}			AltConf	Trace
3	J	210	Total	С	Н	N	О	S	0	0
3	J	210	3389	1095	1690	270	325	9		0
3	M	244	Total	С	Н	N	О	S	0	0
3	101	244	3956	1269	1979	317	381	10	0	U
3	0	186	Total	С	Η	N	O	S	0	0
3	Q	100	3004	976	1494	238	288	8	0	U
3	S	223	Total	С	Н	N	О	S	0	0
	S	229	3628	1170	1814	286	350	8	U	U
3	e	210	Total	С	Н	N	О	S	0	0
	C	210	3384	1095	1685	270	325	9	U	U
3	h	243	Total	С	Η	N	О	S	0	0
	11	240	3933	1263	1965	315	380	10		

• Molecule 4 is a protein called Protein C42.

Mol	Chain	Residues			Atom	.S			AltConf	Trace
4	K	234	Total	С	Н	N	О	S	0	0
4	K	234	3842	1217	1908	332	369	16	0	U
4	L	231	Total	С	Н	N	О	S	0	0
4	L	231	3809	1203	1903	327	360	16	0	U
4	R	220	Total	С	Н	N	О	S	0	0
4	π	220	3625	1150	1807	307	345	16	0	<u> </u>
4	Т	222	Total	С	Н	N	О	S	0	0
4	1	222	3670	1161	1834	315	344	16	0	U
4	f	234	Total	С	Н	N	О	S	0	0
4	1	234	3841	1217	1907	332	369	16	0	U
4	ď	229	Total	С	Н	N	О	S	0	0
4	g	229	3782	1195	1890	324	357	16		U

• Molecule 5 is a protein called Protein AC109.



Mol	Chain	Residues				AltConf	Trace			
5	N	364	Total	С	Н	N	О	S	0	0
	11	304	5917	1913	2953	494	541	16	U	U
5	;	364	Total	С	Н	N	О	S	0	0
	1	304	5917	1913	2953	494	541	16		U

 \bullet Molecule 6 is a protein called Protein AC142.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	0	469	Total	С	Н	N	О	S	0	0
0		409	7709	2507	3852	634	693	23	0	0
6	:	460	Total	С	Н	N	О	S	0	0
0	J	469	7709	2507	3852	634	693	23	U	U

• Molecule 7 is a protein called 38K (AC98).

Mol	Chain	Residues				AltConf	Trace			
7	D	307	Total	С	Н	N	О	S	0	0
'	1	307	5118	1682	2528	424	470	14	U	
7	1,	307	Total	С	Н	N	О	S	0	0
'	K	307	5116	1682	2526	424	470	14	U	

• Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
8	A	1	Total Zn 1 1	0
8	В	1	Total Zn 1 1	0
8	С	1	Total Zn 1 1	0
8	D	1	Total Zn 1 1	0
8	Е	1	Total Zn 1 1	0
8	F	1	Total Zn 1 1	0
8	V	1	Total Zn 1 1	0
8	W	1	Total Zn 1 1	0
8	X	1	Total Zn 1 1	0



Continued from previous page...


Mol	Chain	Residues	Atoms	AltConf
8	Y	1	Total Zn 1 1	0
8	Z	1	Total Zn 1 1	0
8	a	1	Total Zn 1 1	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Major capsid protein







• Molecule 1: Major capsid protein







- Major capsia protoin

Chain Y: 82% 18%











• Molecule 2: Capsid-associated protein VP80






R672
N687
Y688
ASN
VAL
ILE

• Molecule 2: Capsid-associated protein VP80












SEER ASPER A













• Molecule 4: Protein C42











4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	53750	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1400	Depositor
Magnification	64000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.615	Depositor
Minimum map value	-0.816	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.079	Depositor
Recommended contour level	0.294	Depositor
Map size (Å)	541.44, 541.44, 541.44	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.41, 1.41, 1.41	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	langles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.25	0/2258	0.52	0/3058
1	В	0.25	0/2298	0.52	0/3111
1	С	0.27	0/2358	0.54	0/3193
1	D	0.26	0/2338	0.54	0/3166
1	Е	0.28	0/2332	0.54	0/3159
1	F	0.28	0/2357	0.55	0/3194
1	V	0.26	0/2258	0.52	0/3058
1	W	0.25	0/2298	0.51	0/3111
1	X	0.27	0/2358	0.53	0/3193
1	Y	0.26	0/2356	0.54	0/3190
1	Z	0.28	0/2332	0.54	0/3159
1	a	0.27	0/2357	0.54	0/3194
2	G	0.27	0/1650	0.46	0/2221
2	Н	0.26	0/1788	0.46	0/2404
2	I	0.28	0/1914	0.49	0/2572
2	b	0.27	0/1676	0.50	0/2255
2	С	0.26	0/1754	0.48	0/2358
2	d	0.27	0/1914	0.48	0/2572
3	J	0.27	0/1732	0.48	0/2344
3	M	0.29	0/2012	0.52	0/2719
3	Q	0.28	0/1540	0.50	0/2084
3	S	0.28	0/1847	0.50	0/2497
3	е	0.27	0/1732	0.48	0/2344
3	h	0.29	0/2003	0.52	0/2708
4	K	0.29	0/1968	0.55	0/2657
4	L	0.30	0/1941	0.54	0/2621
4	R	0.27	0/1851	0.54	0/2501
4	Т	0.28	0/1870	0.52	0/2525
4	f	0.30	0/1968	0.56	0/2657
4	g	0.29	0/1927	0.54	0/2601
5	N	0.29	0/3040	0.50	0/4132
5	i	0.30	0/3040	0.52	0/4132



Mol	Chain	Bond	Bond lengths		Bond angles	
WIOI		RMSZ	# Z > 5	RMSZ	# Z >5	
6	О	0.30	0/3956	0.51	0/5357	
6	j	0.30	0/3956	0.52	0/5357	
7	Р	0.27	0/2662	0.50	0/3603	
7	k	0.26	0/2662	0.49	0/3603	
All	All	0.28	0/80303	0.52	0/108610	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	I	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	I	594	ARG	Sidechain

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	268/347 (77%)	246 (92%)	22 (8%)	0	100 100



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	В	$273/347\ (79\%)$	255 (93%)	18 (7%)	0	100	100
1	C	$279/347\ (80\%)$	250 (90%)	28 (10%)	1 (0%)	30	68
1	D	$278/347\ (80\%)$	254 (91%)	24 (9%)	0	100	100
1	E	$279/347\ (80\%)$	256 (92%)	23 (8%)	0	100	100
1	F	282/347 (81%)	248 (88%)	33 (12%)	1 (0%)	30	68
1	V	268/347 (77%)	252 (94%)	16 (6%)	0	100	100
1	W	273/347 (79%)	262 (96%)	11 (4%)	0	100	100
1	X	279/347 (80%)	257 (92%)	22 (8%)	0	100	100
1	Y	279/347 (80%)	251 (90%)	28 (10%)	0	100	100
1	Z	279/347 (80%)	246 (88%)	32 (12%)	1 (0%)	30	68
1	a	282/347 (81%)	246 (87%)	36 (13%)	0	100	100
2	G	192/691 (28%)	165 (86%)	27 (14%)	0	100	100
2	Н	208/691 (30%)	178 (86%)	30 (14%)	0	100	100
2	I	220/691 (32%)	205 (93%)	15 (7%)	0	100	100
2	b	195/691 (28%)	176 (90%)	19 (10%)	0	100	100
2	С	204/691 (30%)	171 (84%)	33 (16%)	0	100	100
2	d	220/691 (32%)	203 (92%)	17 (8%)	0	100	100
3	J	206/290 (71%)	194 (94%)	12 (6%)	0	100	100
3	M	240/290 (83%)	213 (89%)	27 (11%)	0	100	100
3	Q	182/290 (63%)	163 (90%)	19 (10%)	0	100	100
3	S	219/290 (76%)	196 (90%)	22 (10%)	1 (0%)	25	64
3	е	206/290 (71%)	193 (94%)	13 (6%)	0	100	100
3	h	239/290 (82%)	207 (87%)	32 (13%)	0	100	100
4	K	232/361 (64%)	206 (89%)	25 (11%)	1 (0%)	30	68
4	L	229/361 (63%)	200 (87%)	29 (13%)	0	100	100
4	R	218/361 (60%)	201 (92%)	17 (8%)	0	100	100
4	Т	220/361 (61%)	190 (86%)	30 (14%)	0	100	100
4	f	232/361 (64%)	202 (87%)	30 (13%)	0	100	100
4	g	227/361 (63%)	197 (87%)	29 (13%)	1 (0%)	30	68
5	N	360/390 (92%)	326 (91%)	34 (9%)	0	100	100
5	i	360/390 (92%)	318 (88%)	41 (11%)	1 (0%)	37	72



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
6	О	467/477 (98%)	434 (93%)	33 (7%)	0	100	100
6	j	467/477 (98%)	436 (93%)	30 (6%)	1 (0%)	44	78
7	Р	303/320~(95%)	272 (90%)	31 (10%)	0	100	100
7	k	303/320~(95%)	272 (90%)	31 (10%)	0	100	100
All	All	9468/14590 (65%)	8541 (90%)	919 (10%)	8 (0%)	50	83

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	K	339	SER
6	j	25	ASP
1	Z	165	PRO
1	С	312	ARG
4	g	318	ILE
1	F	192	ALA
5	i	352	ASP
3	S	147	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers Percenti		ntiles
1	A	247/299~(83%)	246 (100%)	1 (0%)	89	91
1	В	$252/299\ (84\%)$	250 (99%)	2 (1%)	79	85
1	С	$257/299\ (86\%)$	255 (99%)	2 (1%)	79	85
1	D	$257/299\ (86\%)$	253 (98%)	4 (2%)	58	74
1	E	$255/299\ (85\%)$	250 (98%)	5 (2%)	50	69
1	F	$258/299\ (86\%)$	256 (99%)	2 (1%)	79	85
1	V	247/299~(83%)	246 (100%)	1 (0%)	89	91
1	W	$252/299\ (84\%)$	252 (100%)	0	100	100
1	X	$257/299\ (86\%)$	255 (99%)	2 (1%)	79	85



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Rotameric	Outliers	Perce	$\overline{\text{ntiles}}$
1	Y	$258/299\ (86\%)$	257 (100%)	1 (0%)	89	91
1	Z	$255/299\ (85\%)$	253 (99%)	2 (1%)	79	85
1	a	258/299~(86%)	256 (99%)	2 (1%)	79	85
2	G	184/634 (29%)	183 (100%)	1 (0%)	86	90
2	Н	200/634 (32%)	199 (100%)	1 (0%)	86	90
2	I	213/634 (34%)	210 (99%)	3 (1%)	62	76
2	b	187/634 (30%)	186 (100%)	1 (0%)	86	90
2	С	196/634 (31%)	196 (100%)	0	100	100
2	d	213/634 (34%)	212 (100%)	1 (0%)	86	90
3	J	195/273 (71%)	195 (100%)	0	100	100
3	M	228/273 (84%)	227 (100%)	1 (0%)	89	91
3	Q	175/273 (64%)	173 (99%)	2 (1%)	70	80
3	S	210/273 (77%)	208 (99%)	2 (1%)	73	82
3	е	195/273 (71%)	194 (100%)	1 (0%)	86	90
3	h	227/273 (83%)	224 (99%)	3 (1%)	65	77
4	K	222/326 (68%)	221 (100%)	1 (0%)	86	90
4	L	218/326 (67%)	217 (100%)	1 (0%)	86	90
4	R	208/326 (64%)	206 (99%)	2 (1%)	73	82
4	Т	210/326 (64%)	209 (100%)	1 (0%)	86	90
4	f	222/326~(68%)	221 (100%)	1 (0%)	86	90
4	g	217/326 (67%)	212 (98%)	5 (2%)	45	64
5	N	343/366 (94%)	341 (99%)	2 (1%)	84	88
5	i	343/366 (94%)	339 (99%)	4 (1%)	67	79
6	О	435/440 (99%)	433 (100%)	2 (0%)	86	90
6	j	435/440 (99%)	435 (100%)	0	100	100
7	Р	293/304 (96%)	290 (99%)	3 (1%)	73	82
7	k	293/304 (96%)	286 (98%)	7 (2%)	44	63
All	All	8915/13206 (68%)	8846 (99%)	69 (1%)	77	85

All (69) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	CYS



Continued from previous page...

Conti	$nued\ fron$	n previo	ous page
Mol	Chain	Res	Type
1	В	110	MET
1	В	266	LEU
1	B C C D	47	PHE
1	С	231	THR
1	D	57	THR PHE
1	D	67	PHE
1	D	171	ARG
1	D	226	PHE THR
1	Е	77	THR
1	Е	78	ILE
1	Е	166	ASN
1	Е	254	ARG
1	E	301	TYR
1	F	31	THR
1	F	273	LYS
2	G	604	LYS
2	Н	505	MET
2	I	521	LEU
2 2 4	I	592	TYR
2	I	595	GLU
1	K	262	TYR
4	L	157	LYS
3	M	162	CYS ASP
5	N	45	ASP
5	N	216	ARG LYS
6	О	250	
6	О	458	ASP
7	Р	33	LYS
7	Р	66	TYR
7	Р	211	ARG
3	Q	69	GLU
3	Q R	144	MET
4	R	174	CYS CYS
4	R	216	
3	S	10	ARG
3	S	147	ARG
4	Т	141	LYS
1	V	169	CYS
1	X	49	CYS
1	X	132	CYS
1	Y	176	GLU
1	Z	163	THR



Continued from previous page...

Mol	Chain	Res	$oxed{ ext{Type}}$
1	Z	164	ASN
1	a	44	ASP
1	a	100	ASN
2	b	480	LYS
2	d	544	MET
3	е	234	TYR
4	f	171	ASP
4	g	117	ARG
4	g	141	LYS
4	g	158	TYR
4	g	223	ARG
4	g	309	TYR
3	h	156	ASN
3	h	157	PHE
3	h	165	ARG
5	i	45	ASP
5	i	185	LEU
5	i	220	ASN
5	i	364	PHE
7	k	33	LYS
7	k	165	GLU
7	k	181	ARG
7	k	198	ASP
7	k	211	ARG
7	k	230	ARG
7	k	258	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (70) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	ASN
1	A	218	GLN
1	В	164	ASN
1	В	262	GLN
1	С	81	HIS
1	С	126	ASN
1	D	129	ASN
1	Ε	119	HIS
1	F	51	ASN
2	Н	516	ASN
2	Н	570	HIS
2	Н	617	ASN



Continued from previous page...

Continued from previous page				
Mol	Chain	Res	Type	
2	Н	659	ASN	
2	I	513	ASN	
2		659	ASN	
4	L	128	HIS	
4	L	159	HIS	
4	L	258	GLN	
4	L	336	ASN	
3	M	93	ASN	
3	M	218	HIS	
5	N	220	ASN	
5	N	254	ASN	
6	О	14	HIS	
6	О	169	ASN	
6	0	198	HIS	
6	О	203	ASN	
6	О	263	ASN	
6	О	281	GLN	
6	О	351	HIS	
6	O O O P P	406	GLN	
7	Р	65	ASN	
7	Р	248	ASN	
3	Q	92	HIS	
4	R	260	GLN	
3	S	73	ASN	
4	Т	133	ASN	
1	V	144	ASN	
1	V	218	GLN	
1	W	42	HIS	
1	W	164	ASN	
1	W	262	GLN	
1	X	112	GLN	
1	X Y	205	GLN	
1	Y	124	ASN	
1	Y	264	ASN	
1	Y	308	ASN	
1	Z	146	GLN	
1	a	184	ASN	
1	a	251	ASN	
2	b	617	ASN	
2	b	624	ASN	
2	С	535	ASN	
2	d	617	ASN	



Continued from previous page					
	($\ 'ontinue a$	t trom	previous	page

Mol	Chain	Res	Type
2	d	661	GLN
2	d	678	GLN
3	е	156	ASN
3	е	241	HIS
4	f	129	HIS
4	f	258	GLN
4	f	299	ASN
4	f	308	ASN
3	h	96	HIS
5	i	249	ASN
5	i	302	ASN
6	j	14	HIS
6	j	169	ASN
6	j j j	181	ASN
6	j	226	ASN
7	k	266	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-43589. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)


6.2.1 Primary map








Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192

The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)


6.3.1 Primary map








Y Index: 215




Z Index: 168


6.3.2 Raw map



X Index: 3



Y Index: 3




Z Index: 160

The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.294. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_43589_msk_1.map (i)






7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $1916~\mathrm{nm^3}$; this corresponds to an approximate mass of $1731~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.209 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



^{*}Reported resolution corresponds to spatial frequency of 0.209 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estim	ation o	criterion (FSC cut-off)
rtesolution estimate (A)	0.143	0.5	Half-bit
Reported by author	4.78	-	-
Author-provided FSC curve	4.78	6.78	4.98
Unmasked-calculated*	8.25	10.49	8.47


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.25 differs from the reported value 4.78 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-43589 and PDB model 8VWJ. Per-residue inclusion information can be found in section 3 on page 9.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.294 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.294).



9.4 Atom inclusion (i)



At the recommended contour level, 69% of all backbone atoms, 68% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.294) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6830	0.2430
A	0.4870	0.2150
В	0.4540	0.2090
С	0.6940	0.2290
D	0.7220	0.2420
E	0.7760	0.2510
F	0.7540	0.2490
G	0.4180	0.2260
Н	0.5000	0.2250
I	0.7290	0.2320
J	0.6760	0.2520
K	0.7280	0.2600
L	0.7540	0.2520
M	0.7460	0.2780
N	0.8160	0.2950
O	0.8560	0.2900
P	0.5880	0.2400
Q	0.7720	0.2690
R	0.8260	0.2570
S	0.7400	0.2780
Т	0.8020	0.2620
V	0.5130	0.1940
W	0.5030	0.1780
X	0.6660	0.2190
Y	0.7400	0.2290
Z	0.7740	0.2330
a	0.6780	0.2270
b	0.4070	0.2100
С	0.5210	0.2020
d	0.7930	0.2090
e	0.6490	0.2340
f	0.7100	0.2310
g	0.7480	0.2430
h	0.7460	0.2710
i	0.8510	0.2690





 $Continued\ from\ previous\ page...$

Chain	Atom inclusion	Q-score
j	0.8590	0.2740
k	0.5910	0.2330

