



Full wwPDB X-ray Structure Validation Report i

Sep 24, 2024 – 04:08 PM JST

PDB ID : 8W7R
Title : H. walsbyi bacteriorhodopsin mutant - W94F
Authors : Li, G.Y.; Chen, J.C.; Yang, C.S.
Deposited on : 2023-08-31
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.002 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 3743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bacteriorhodopsin-I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	229	Total	C 1766	N 1176	O 271	S 312	7	0	1	0
1	B	229	Total	C 1760	N 1173	O 270	S 310	7	0	0	0

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q18DH8
A	2	ALA	-	expression tag	UNP Q18DH8
A	94	PHE	TRP	engineered mutation	UNP Q18DH8
A	255	LEU	-	expression tag	UNP Q18DH8
A	256	GLU	-	expression tag	UNP Q18DH8
A	257	HIS	-	expression tag	UNP Q18DH8
A	258	HIS	-	expression tag	UNP Q18DH8
A	259	HIS	-	expression tag	UNP Q18DH8
A	260	HIS	-	expression tag	UNP Q18DH8
A	261	HIS	-	expression tag	UNP Q18DH8
A	262	HIS	-	expression tag	UNP Q18DH8
B	1	MET	-	initiating methionine	UNP Q18DH8
B	2	ALA	-	expression tag	UNP Q18DH8
B	94	PHE	TRP	engineered mutation	UNP Q18DH8
B	255	LEU	-	expression tag	UNP Q18DH8
B	256	GLU	-	expression tag	UNP Q18DH8
B	257	HIS	-	expression tag	UNP Q18DH8
B	258	HIS	-	expression tag	UNP Q18DH8
B	259	HIS	-	expression tag	UNP Q18DH8
B	260	HIS	-	expression tag	UNP Q18DH8
B	261	HIS	-	expression tag	UNP Q18DH8
B	262	HIS	-	expression tag	UNP Q18DH8

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 16 10 6	0	0
3	B	1	Total C O 16 10 6	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 16 10 6	0	0
3	B	1	Total C O 16 10 6	0	0

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 25 21 4	0	0
4	B	1	Total C O 25 21 4	0	0
4	B	1	Total C O 25 21 4	0	0

- Molecule 5 is RETINAL (three-letter code: RET) (formula: C₂₀H₂₈O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C 20 20	0	0
5	B	1	Total C 20 20	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	12	Total O 12 12	0	0
6	B	14	Total O 14 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacteriorhodopsin-I



- Molecule 1: Bacteriorhodopsin-I



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.50Å 126.75Å 55.04Å 90.00° 111.44° 90.00°	Depositor
Resolution (Å)	26.96 – 2.50 26.96 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (26.96-2.50) 99.5 (26.96-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.09 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R , R_{free}	0.213 , 0.257 0.213 , 0.257	Depositor DCC
R_{free} test set	1088 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.5	Xtriage
Anisotropy	0.233	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 58.3	EDS
L-test for twinning ²	$< L > = 0.51$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3743	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OLC, 1PE, RET

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.41	0/1804	0.76	0/2460
1	B	0.40	0/1798	0.77	0/2452
All	All	0.41	0/3602	0.76	0/4912

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	233	ARG	Sidechain
1	B	235	ARG	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1766	0	1813	14	0
1	B	1760	0	1809	18	0
2	A	6	0	8	0	0
2	B	6	0	8	0	0
3	A	16	0	22	1	0
3	B	48	0	66	1	0
4	A	25	0	40	0	0
4	B	50	0	80	0	0
5	A	20	0	27	2	0
5	B	20	0	27	5	0
6	A	12	0	0	2	0
6	B	14	0	0	0	0
All	All	3743	0	3900	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:410:HOH:O	1:B:11:LEU:HD12	1.64	0.97
1:B:11:LEU:HB3	1:B:17:GLY:HA3	1.70	0.71
1:B:235:ARG:NH2	1:B:235:ARG:H	1.89	0.71
5:B:307:RET:H161	5:B:307:RET:H8	1.73	0.70
1:B:113:GLN:HE21	3:B:304:1PE:H152	1.66	0.60
5:A:304:RET:H161	5:A:304:RET:H8	1.86	0.55
1:A:90:ARG:NH1	6:A:401:HOH:O	2.30	0.53
1:B:94:PHE:HB3	1:B:127:ILE:HD12	1.89	0.53
1:A:74:THR:HG22	1:A:75:GLU:N	2.25	0.52
1:A:94:PHE:HB3	1:A:127:ILE:HD12	1.89	0.52
1:B:194:PRO:HB3	5:B:307:RET:H183	1.91	0.51
1:A:147:THR:O	1:A:151:ILE:HG12	2.11	0.51
1:A:90:ARG:HH11	1:A:90:ARG:HG3	1.76	0.50
1:A:54:ILE:O	1:A:57:PRO:HD2	2.13	0.49
1:B:74:THR:HG22	1:B:75:GLU:N	2.29	0.48
5:B:307:RET:H161	5:B:307:RET:C8	2.39	0.47
5:A:304:RET:H161	5:A:304:RET:C8	2.45	0.47
1:B:90:ARG:HG3	1:B:90:ARG:HH11	1.79	0.47
1:B:136:THR:HG22	1:B:138:VAL:H	1.80	0.46
1:B:136:THR:HG21	1:B:141:ALA:HB3	1.97	0.46
1:B:174:ASP:OD1	1:B:235:ARG:NH2	2.48	0.46
1:A:74:THR:CG2	1:A:75:GLU:N	2.78	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54:ILE:O	1:B:57:PRO:HD2	2.17	0.45
1:B:147:THR:O	1:B:151:ILE:HG12	2.15	0.45
5:B:307:RET:H7	5:B:307:RET:H181	1.71	0.45
1:B:56:ILE:HB	1:B:57:PRO:HD3	1.99	0.45
1:A:19:TRP:HB2	1:A:213:THR:HG22	1.98	0.44
1:A:56:ILE:HB	1:A:57:PRO:HD3	1.98	0.44
1:B:74:THR:CG2	1:B:75:GLU:N	2.81	0.43
3:A:302:1PE:H221	3:A:302:1PE:HI31	1.55	0.43
1:A:237:ILE:O	1:A:238:MET:HB2	2.18	0.43
1:B:146:TRP:CD1	1:B:197:TRP:CE3	3.07	0.42
1:B:193:TYR:HD1	1:B:216:PHE:HE1	1.68	0.41
1:A:43:ASP:HA	1:A:44:PRO:HD3	1.94	0.41
1:A:146:TRP:CD1	1:A:197:TRP:CE3	3.09	0.41
1:B:98:THR:HG22	5:B:307:RET:H202	2.02	0.41
1:A:193:TYR:N	1:A:194:PRO:HD2	2.36	0.41
1:A:161:VAL:HG22	1:A:183:ARG:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	228/262 (87%)	225 (99%)	2 (1%)	1 (0%)	30 49
1	B	227/262 (87%)	225 (99%)	2 (1%)	0	100 100
All	All	455/524 (87%)	450 (99%)	4 (1%)	1 (0%)	44 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	238	MET

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	182/207 (88%)	178 (98%)	4 (2%)	47 73
1	B	181/207 (87%)	175 (97%)	6 (3%)	33 59
All	All	363/414 (88%)	353 (97%)	10 (3%)	38 65

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LEU
1	A	47	LYS
1	A	104	ASP
1	A	160	LEU
1	B	47	LYS
1	B	104	ASP
1	B	140	VAL
1	B	160	LEU
1	B	173	GLU
1	B	235	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	113	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	OLC	A	303	-	24,24,24	0.23	0	25,25,25	0.33	0
3	1PE	B	304	-	15,15,15	0.34	0	14,14,14	0.29	0
5	RET	B	307	1	20,20,21	2.77	5 (25%)	27,27,28	1.45	4 (14%)
4	OLC	B	301	-	24,24,24	0.33	0	25,25,25	0.29	0
2	GOL	B	306	-	5,5,5	0.09	0	5,5,5	0.34	0
4	OLC	B	302	-	24,24,24	0.24	0	25,25,25	0.43	0
3	1PE	B	303	-	15,15,15	0.18	0	14,14,14	0.13	0
3	1PE	A	302	-	15,15,15	0.26	0	14,14,14	0.12	0
2	GOL	A	301	-	5,5,5	0.12	0	5,5,5	0.51	0
5	RET	A	304	1	20,20,21	2.39	3 (15%)	27,27,28	1.44	5 (18%)
3	1PE	B	305	-	15,15,15	0.26	0	14,14,14	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	A	303	-	-	10/24/24/24	-
3	1PE	B	304	-	-	9/13/13/13	-
5	RET	B	307	1	-	0/13/30/31	0/1/1/1
4	OLC	B	301	-	-	15/24/24/24	-
2	GOL	B	306	-	-	2/4/4/4	-
4	OLC	B	302	-	-	10/24/24/24	-
3	1PE	B	303	-	-	6/13/13/13	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1PE	A	302	-	-	2/13/13/13	-
2	GOL	A	301	-	-	4/4/4/4	-
5	RET	A	304	1	-	0/13/30/31	0/1/1/1
3	1PE	B	305	-	-	8/13/13/13	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	307	RET	C14-C13	9.94	1.41	1.33
5	A	304	RET	C14-C13	8.90	1.40	1.33
5	B	307	RET	C10-C9	5.29	1.42	1.35
5	A	304	RET	C10-C9	3.41	1.40	1.35
5	B	307	RET	C8-C9	-3.34	1.38	1.45
5	A	304	RET	C8-C9	-2.33	1.40	1.45
5	B	307	RET	C12-C13	-2.20	1.41	1.45
5	B	307	RET	C11-C12	2.05	1.39	1.34

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	304	RET	C19-C9-C10	-4.07	117.23	122.92
5	B	307	RET	C2-C1-C6	3.24	115.48	110.48
5	B	307	RET	C11-C10-C9	3.18	131.85	127.31
5	B	307	RET	C19-C9-C8	3.16	123.06	118.08
5	A	304	RET	C19-C9-C8	2.93	122.69	118.08
5	B	307	RET	C19-C9-C10	-2.85	118.93	122.92
5	A	304	RET	C2-C1-C6	2.73	114.69	110.48
5	A	304	RET	C11-C12-C13	2.18	132.53	126.42
5	A	304	RET	C11-C10-C9	2.02	130.19	127.31

There are no chirality outliers.

All (66) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	GOL	O1-C1-C2-O2
2	A	301	GOL	O1-C1-C2-C3
2	A	301	GOL	C1-C2-C3-O3
2	B	306	GOL	O1-C1-C2-C3
4	B	301	OLC	C21-C22-C24-O25
4	B	302	OLC	O20-C21-C22-C24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	302	1PE	C13-C23-OH3-C22
4	B	302	OLC	O20-C21-C22-O23
3	B	303	1PE	OH5-C14-C24-OH4
3	B	303	1PE	OH6-C15-C25-OH5
3	B	304	1PE	OH5-C14-C24-OH4
3	B	305	1PE	OH4-C13-C23-OH3
4	A	303	OLC	C14-C15-C16-C17
4	A	303	OLC	C13-C14-C15-C16
4	B	301	OLC	C6-C7-C8-C9
3	B	304	1PE	OH7-C16-C26-OH6
4	A	303	OLC	C5-C6-C7-C8
4	B	301	OLC	C4-C5-C6-C7
4	A	303	OLC	C11-C12-C13-C14
4	B	301	OLC	C13-C14-C15-C16
4	B	302	OLC	C5-C6-C7-C8
2	A	301	GOL	O2-C2-C3-O3
2	B	306	GOL	O1-C1-C2-O2
4	B	301	OLC	O23-C22-C24-O25
4	B	301	OLC	C10-C11-C12-C13
4	A	303	OLC	O20-C21-C22-C24
4	A	303	OLC	C4-C5-C6-C7
4	B	302	OLC	C2-C3-C4-C5
3	B	304	1PE	OH2-C12-C22-OH3
3	B	305	1PE	OH6-C15-C25-OH5
4	B	302	OLC	C14-C15-C16-C17
4	B	301	OLC	C15-C16-C17-C18
4	B	301	OLC	O20-C21-C22-C24
4	B	301	OLC	C2-C3-C4-C5
3	B	304	1PE	OH6-C15-C25-OH5
4	B	302	OLC	C3-C4-C5-C6
3	B	303	1PE	C13-C23-OH3-C22
3	B	305	1PE	C13-C23-OH3-C22
4	B	301	OLC	C3-C4-C5-C6
3	B	305	1PE	C14-C24-OH4-C13
4	B	301	OLC	C1-C2-C3-C4
4	B	302	OLC	C4-C5-C6-C7
3	A	302	1PE	OH7-C16-C26-OH6
4	B	302	OLC	C7-C8-C9-C10
3	B	304	1PE	C13-C23-OH3-C22
3	B	304	1PE	C12-C22-OH3-C23
3	B	305	1PE	C16-C26-OH6-C15
3	B	305	1PE	C23-C13-OH4-C24

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	304	1PE	C23-C13-OH4-C24
4	B	301	OLC	O20-C21-C22-O23
3	B	305	1PE	C25-C15-OH6-C26
3	B	303	1PE	OH7-C16-C26-OH6
3	B	303	1PE	OH4-C13-C23-OH3
3	B	303	1PE	C15-C25-OH5-C14
4	B	301	OLC	C11-C12-C13-C14
4	B	302	OLC	C10-C11-C12-C13
4	B	301	OLC	C5-C6-C7-C8
4	A	303	OLC	O20-C21-C22-O23
4	B	302	OLC	C15-C16-C17-C18
4	B	301	OLC	C7-C8-C9-C10
3	B	304	1PE	C16-C26-OH6-C15
4	A	303	OLC	C9-C10-C11-C12
3	B	304	1PE	C14-C24-OH4-C13
4	A	303	OLC	C2-C3-C4-C5
3	B	305	1PE	C24-C14-OH5-C25
4	A	303	OLC	C22-C21-O20-C1


There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	304	1PE	1	0
5	B	307	RET	5	0
3	A	302	1PE	1	0
5	A	304	RET	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	229/262 (87%)	-0.06	4 (1%)	69	65	18, 38, 67, 94 1 (0%)
1	B	229/262 (87%)	-0.04	4 (1%)	69	65	28, 38, 69, 96 0
All	All	458/524 (87%)	-0.05	8 (1%)	69	65	18, 38, 68, 96 1 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	GLY	4.8
1	B	11	LEU	3.9
1	A	238	MET	3.5
1	A	11	LEU	3.2
1	B	239	GLY	2.6
1	B	238	MET	2.3
1	A	13	VAL	2.2
1	B	74	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,


median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	OLC	B	302	25/25	0.74	0.19	50,108,130,142	0
4	OLC	A	303	25/25	0.77	0.21	58,82,101,104	0
3	1PE	B	305	16/16	0.78	0.17	69,74,85,87	0
3	1PE	A	302	16/16	0.81	0.22	72,82,95,99	0
2	GOL	A	301	6/6	0.83	0.15	57,60,75,81	0
4	OLC	B	301	25/25	0.90	0.16	40,48,99,101	0
2	GOL	B	306	6/6	0.91	0.10	69,76,79,84	0
3	1PE	B	304	16/16	0.91	0.13	50,60,78,79	0
3	1PE	B	303	16/16	0.92	0.13	55,67,77,77	0
5	RET	B	307	20/21	0.92	0.10	31,38,43,43	0
5	RET	A	304	20/21	0.94	0.08	31,34,39,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.