

Full wwPDB X-ray Structure Validation Report (i)

Mar 24, 2025 – 02:17 PM JST

PDB ID : 8YVH

Title: NADPH complex of Imine Reductase Mutant(M6) from Pochonia chlamy-

dosporia 170

Authors: Shi, M.; zheng, G.W.

Deposited on : 2024-03-28

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.21

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.004 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.41.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	293	88%	11%	-
1	В	293	84%	13%	



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4782 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Oxidoreductase.

\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	293	Total 2133	C 1336	N 360	O 424	S 13	0	0	0
1	В	287	Total 2090	C 1312	N 352	O 414	S 12	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	GLU	SER	conflict	UNP A0A179FZ20
A	47	PRO	ALA	conflict	UNP A0A179FZ20
A	50	ALA	VAL	conflict	UNP A0A179FZ20
A	72	ASN	SER	conflict	UNP A0A179FZ20
A	73	GLN	TYR	conflict	UNP A0A179FZ20
A	76	VAL	THR	conflict	UNP A0A179FZ20
A	124	THR	VAL	conflict	UNP A0A179FZ20
A	125	THR	PRO	conflict	UNP A0A179FZ20
A	127	GLU	ILE	conflict	UNP A0A179FZ20
A	128	TYR	LEU	conflict	UNP A0A179FZ20
A	178	VAL	ILE	conflict	UNP A0A179FZ20
A	180	SER	GLY	conflict	UNP A0A179FZ20
A	195	ALA	LYS	conflict	UNP A0A179FZ20
A	210	LEU	PHE	conflict	UNP A0A179FZ20
A	212	ALA	LYS	conflict	UNP A0A179FZ20
A	217	GLN	TRP	conflict	UNP A0A179FZ20
A	220	VAL	ARG	conflict	UNP A0A179FZ20
A	224	ARG	HIS	conflict	UNP A0A179FZ20
A	236	ILE	LEU	conflict	UNP A0A179FZ20
A	239	GLY	GLN	conflict	UNP A0A179FZ20
В	40	GLU	SER	conflict	UNP A0A179FZ20
В	47	PRO	ALA	conflict	UNP A0A179FZ20
В	50	ALA	VAL	conflict	UNP A0A179FZ20
В	72	ASN	SER	conflict	UNP A0A179FZ20
В	73	GLN	TYR	conflict	UNP A0A179FZ20


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	76	VAL	THR	conflict	UNP A0A179FZ20
В	124	THR	VAL	conflict	UNP A0A179FZ20
В	125	THR	PRO	conflict	UNP A0A179FZ20
В	127	GLU	ILE	conflict	UNP A0A179FZ20
В	128	TYR	LEU	conflict	UNP A0A179FZ20
В	178	VAL	ILE	conflict	UNP A0A179FZ20
В	180	SER	GLY	conflict	UNP A0A179FZ20
В	195	ALA	LYS	conflict	UNP A0A179FZ20
В	210	LEU	PHE	conflict	UNP A0A179FZ20
В	212	ALA	LYS	conflict	UNP A0A179FZ20
В	217	GLN	TRP	conflict	UNP A0A179FZ20
В	220	VAL	ARG	conflict	UNP A0A179FZ20
В	224	ARG	HIS	conflict	UNP A0A179FZ20
В	236	ILE	LEU	conflict	UNP A0A179FZ20
В	239	GLY	GLN	conflict	UNP A0A179FZ20


• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	Λ	1	Total	С	N	О	Р	0	0
2	2 A	1	48	21	7	17	3	U	
9	D	1	Total	С	N	О	Р	0	0
	Б	1	48	21	7	17	3	U	U

• Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 8 4 1 3	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	254	Total O 254 254	0	0
4	В	201	Total O 201 201	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Oxidoreductase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	37.40Å 117.95Å 63.53Å	Donositor
a, b, c, α , β , γ	90.00° 100.89° 90.00°	Depositor
Resolution (Å)	35.09 - 2.10	Depositor
Resolution (A)	35.09 - 2.10	EDS
% Data completeness	99.3 (35.09-2.10)	Depositor
(in resolution range)	99.4 (35.09-2.10)	EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	6.91 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
D D.	0.167 , 0.241	Depositor
R, R_{free}	0.175 , 0.247	DCC
R_{free} test set	1602 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 42.0	EDS
L-test for twinning ²	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4782	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond	angles
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.74	0/2159	0.80	0/2931
1	В	0.73	0/2116	0.80	0/2874
All	All	0.74	0/4275	0.80	0/5805

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2133	0	2199	34	0
1	В	2090	0	2154	29	0
2	A	48	0	25	0	0
2	В	48	0	25	2	0
3	A	8	0	12	2	0
4	A	254	0	0	6	0
4	В	201	0	0	8	0
All	All	4782	0	4415	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.



All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:73:GLN:HE22	1:B:93:ASN:HD21	1.29	0.78
1:A:179:MET:HE2	1:B:214:MET:SD	2.23	0.78
1:A:198:GLU:OE1	1:B:167:LYS:HE3	1.91	0.70
1:A:238:MET:HE1	4:B:549:HOH:O	1.92	0.69
1:A:1:MET:N	4:A:401:HOH:O	2.25	0.68
1:A:210:LEU:HD11	4:B:565:HOH:O	1.97	0.64
1:A:214:MET:HG2	1:B:182:VAL:HG11	1.82	0.61
1:B:224:ARG:HD2	4:B:475:HOH:O	2.03	0.59
1:B:224:ARG:NH2	4:B:406:HOH:O	2.35	0.58
1:B:135:VAL:HA	1:B:157:THR:O	2.04	0.58
1:B:73:GLN:NE2	1:B:93:ASN:HD21	2.00	0.57
1:B:73:GLN:HE21	1:B:73:GLN:HA	1.70	0.55
1:A:115:GLU:HG2	4:A:625:HOH:O	2.07	0.55
1:B:224:ARG:CD	4:B:475:HOH:O	2.55	0.54
1:B:73:GLN:HE22	1:B:93:ASN:ND2	2.02	0.53
1:A:182:VAL:HG22	1:B:182:VAL:HG22	1.90	0.53
1:A:238:MET:HE2	2:B:301:NAP:C2N	2.39	0.53
1:A:135:VAL:HG22	1:A:157:THR:HG23	1.93	0.51
3:A:302:TRS:H22	4:A:586:HOH:O	2.11	0.50
1:A:132:PRO:O	1:A:157:THR:CG2	2.59	0.50
1:A:211:VAL:HG11	1:B:218:LEU:CD1	2.43	0.48
1:A:132:PRO:O	1:A:157:THR:HG22	2.13	0.48
1:B:46:VAL:HB	1:B:47:PRO:HD3	1.95	0.47
1:B:152:LEU:HB3	1:B:158:THR:HG21	1.97	0.47
1:A:238:MET:CE	2:B:301:NAP:C2N	2.92	0.47
1:A:139:SER:HB2	1:A:164:ASP:O	2.14	0.47
1:A:108:LEU:O	1:A:112:ARG:HG2	2.13	0.46
1:A:238:MET:CE	4:B:549:HOH:O	2.56	0.46
1:A:227:GLU:HG3	4:A:607:HOH:O	2.15	0.46
1:B:193:VAL:HG13	1:B:198:GLU:HB2	1.98	0.46
1:B:84:ILE:HD12	1:B:85:ASP:N	2.31	0.46
1:A:142:LYS:O	1:A:146:GLU:HG2	2.16	0.45
1:A:152:LEU:HB3	1:A:158:THR:HG21	1.97	0.45
3:A:302:TRS:H31	4:A:586:HOH:O	2.17	0.45
1:B:209:PRO:O	1:B:212:ALA:HB3	2.17	0.45
1:A:120:GLY:HA2	1:A:137:LEU:O	2.16	0.45
1:B:73:GLN:O	1:B:77:GLN:HG3	2.16	0.45
1:B:19:LEU:HD23	1:B:19:LEU:HA	1.90	0.44
1:A:210:LEU:CG	4:B:565:HOH:O	2.64	0.44
1:A:210:LEU:CD1	4:B:565:HOH:O	2.62	0.43

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:A:179:MET:CE	1:A:179:MET:HA	2.48	0.43
1:B:120:GLY:HA2	1:B:137:LEU:O	2.19	0.43
1:B:10:SER:HA	1:B:33:THR:O	2.18	0.43
1:A:224:ARG:HD3	4:A:429:HOH:O	2.19	0.42
1:A:135:VAL:HG22	1:A:157:THR:CG2	2.49	0.42
1:A:179:MET:CE	1:B:214:MET:CG	2.98	0.42
1:A:286:LEU:HD22	1:B:187:VAL:HG12	2.01	0.41
1:A:78:GLN:O	1:A:81:ASP:HB2	2.20	0.41
1:A:239:GLY:HA3	1:A:266:MET:HE1	2.02	0.41
1:B:188:GLN:OE1	1:B:257:VAL:HA	2.20	0.41
1:A:184:GLU:OE2	1:B:184:GLU:OE2	2.38	0.41
1:B:218:LEU:N	1:B:219:PRO:CD	2.84	0.41
1:A:1:MET:HG2	1:A:28:SER:O	2.21	0.40
1:A:135:VAL:HA	1:A:157:THR:O	2.21	0.40
1:B:214:MET:SD	1:B:214:MET:N	2.95	0.40
1:A:179:MET:HE2	1:B:214:MET:CE	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	291/293 (99%)	287 (99%)	4 (1%)	0	100	100
1	В	285/293 (97%)	278 (98%)	7 (2%)	0	100	100
All	All	576/586 (98%)	565 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	235/235 (100%)	229 (97%)	6 (3%)	41	46
1	В	229/235 (97%)	216 (94%)	13 (6%)	17	15
All	All	464/470 (99%)	445 (96%)	19 (4%)	26	27

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	ASP
1	A	110	GLN
1	A	163	ASP
1	A	214	MET
1	A	228	LYS
1	A	279	ILE
1	В	73	GLN
1	В	74	ASP
1	В	80	LEU
1	В	163	ASP
1	В	206	LEU
1	В	210	LEU
1	В	213	SER
1	В	214	MET
1	В	217	GLN
1	В	238	MET
1	В	241	GLU
1	В	257	VAL
1	В	279	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	96	ASN
1	A	110	GLN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	В	8	HIS
1	В	16	ASN
1	В	73	GLN
1	В	131	GLN
1	В	217	GLN
1	В	245	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Chain Res Link Bond lengths		В	ond ang	gles			
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	В	301	-	45,52,52	1.20	3 (6%)	56,80,80	1.38	10 (17%)
2	NAP	A	301	-	45,52,52	1.12	4 (8%)	56,80,80	1.28	8 (14%)
3	TRS	A	302	-	7,7,7	0.19	0	9,9,9	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



, ,		, 1.	C 1	. 1 . 1		· 1 / · C 1
- means	no	outliers	of tha	t kind	were	identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	В	301	-	-	5/31/67/67	0/5/5/5
2	NAP	A	301	-	-	5/31/67/67	0/5/5/5
3	TRS	A	302	-	-	5/9/9/9	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	В	301	NAP	O4B-C1B	4.41	1.47	1.41
2	A	301	NAP	O4B-C1B	2.95	1.45	1.41
2	В	301	NAP	C2N-N1N	2.22	1.37	1.35
2	A	301	NAP	P2B-O2B	2.21	1.63	1.59
2	A	301	NAP	C7N-N7N	-2.19	1.28	1.33
2	A	301	NAP	C5A-C4A	2.13	1.46	1.40
2	В	301	NAP	P2B-O2B	2.05	1.63	1.59

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	301	NAP	N3A-C2A-N1A	-4.16	122.17	128.68
2	A	301	NAP	N3A-C2A-N1A	-3.79	122.75	128.68
2	A	301	NAP	O7N-C7N-N7N	-3.06	118.22	122.58
2	В	301	NAP	O2B-P2B-O1X	-2.76	98.75	109.39
2	В	301	NAP	PN-O3-PA	-2.75	123.39	132.83
2	A	301	NAP	O7N-C7N-C3N	2.70	122.86	119.63
2	В	301	NAP	N6A-C6A-N1A	2.50	123.76	118.57
2	A	301	NAP	O2A-PA-O1A	2.48	124.50	112.24
2	A	301	NAP	C2A-N1A-C6A	2.39	122.85	118.75
2	A	301	NAP	C3B-C2B-C1B	-2.38	98.42	102.89
2	В	301	NAP	C3N-C2N-N1N	2.38	122.75	120.43
2	В	301	NAP	O2N-PN-O1N	2.35	123.87	112.24
2	В	301	NAP	O2A-PA-O1A	2.35	123.84	112.24
2	A	301	NAP	C1B-N9A-C4A	-2.32	122.57	126.64
2	В	301	NAP	O2X-P2B-O1X	2.29	119.64	110.68
2	A	301	NAP	PN-O3-PA	-2.07	125.73	132.83
2	В	301	NAP	C5A-C6A-N6A	-2.05	117.24	120.35
2	В	301	NAP	C6N-N1N-C2N	-2.04	120.11	121.97

There are no chirality outliers.

All (15) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	301	NAP	C5B-O5B-PA-O2A
2	A	301	NAP	O4D-C1D-N1N-C2N
2	В	301	NAP	C2B-O2B-P2B-O1X
2	В	301	NAP	C2B-O2B-P2B-O2X
2	В	301	NAP	O4D-C1D-N1N-C2N
2	В	301	NAP	O4D-C1D-N1N-C6N
3	A	302	TRS	C2-C-C1-O1
3	A	302	TRS	C3-C-C1-O1
2	В	301	NAP	O4B-C4B-C5B-O5B
3	A	302	TRS	C1-C-C2-O2
2	A	301	NAP	C1B-C2B-O2B-P2B
2	A	301	NAP	C3B-C2B-O2B-P2B
2	A	301	NAP	O4B-C4B-C5B-O5B
3	A	302	TRS	N-C-C1-O1
3	A	302	TRS	N-C-C2-O2


There are no ring outliers.

2 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	301	NAP	2	0
3	A	302	TRS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	293/293 (100%)	-0.35	1 (0%) 90 91	12, 20, 36, 53	0
1	В	287/293 (97%)	-0.19	1 (0%) 90 91	11, 23, 40, 56	0
All	All	580/586 (98%)	-0.27	2 (0%) 90 91	11, 22, 40, 56	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	209	PRO	2.5
1	В	110	GLN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	TRS	A	302	8/8	0.84	0.12	26,29,32,33	0
2	NAP	В	301	48/48	0.96	0.06	14,20,26,27	0
2	NAP	A	301	48/48	0.98	0.05	14,17,23,24	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

