



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 20, 2025 – 05:54 PM JST

PDB ID : 8ZEB
Title : Crystal structure of BCL-XL bound by cp-B6X-4
Authors : Sun, X.N.; Liu, J.X.; Wu, D.L.
Deposited on : 2024-05-05
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> % 93% 6% . </div> </div>
1	B	141	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 1%, green 96%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> 2% 87% 11% . </div> </div>
2	C	15	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 87%, yellow 7%, orange 7%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> 87% 7% 7% </div> </div>
2	D	15	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 93%, yellow 7%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> 93% 7% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2661 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bcl-2-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	0	0
			1128	723	190	211	4			
1	B	139	Total	C	N	O	S	0	1	0
			1131	725	190	212	4			

There are 110 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	PHE	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	ASP	deletion	UNP Q07817
A	?	-	VAL	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	ASN	deletion	UNP Q07817
A	?	-	ARG	deletion	UNP Q07817
A	?	-	THR	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	ALA	deletion	UNP Q07817
A	?	-	PRO	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	GLY	deletion	UNP Q07817
A	?	-	THR	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	MET	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	THR	deletion	UNP Q07817
A	?	-	PRO	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	ALA	deletion	UNP Q07817
A	?	-	ILE	deletion	UNP Q07817

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASN	deletion	UNP Q07817
A	?	-	GLY	deletion	UNP Q07817
A	?	-	ASN	deletion	UNP Q07817
A	?	-	PRO	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	TRP	deletion	UNP Q07817
A	?	-	HIS	deletion	UNP Q07817
A	?	-	LEU	deletion	UNP Q07817
A	?	-	ALA	deletion	UNP Q07817
A	?	-	ASP	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	PRO	deletion	UNP Q07817
A	?	-	ALA	deletion	UNP Q07817
A	?	-	VAL	deletion	UNP Q07817
A	?	-	ASN	deletion	UNP Q07817
A	?	-	GLY	deletion	UNP Q07817
A	?	-	ALA	deletion	UNP Q07817
A	?	-	THR	deletion	UNP Q07817
A	?	-	GLY	deletion	UNP Q07817
A	?	-	HIS	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	SER	deletion	UNP Q07817
A	?	-	LEU	deletion	UNP Q07817
A	?	-	ASP	deletion	UNP Q07817
A	?	-	ALA	deletion	UNP Q07817
A	?	-	ARG	deletion	UNP Q07817
A	?	-	GLU	deletion	UNP Q07817
A	?	-	VAL	deletion	UNP Q07817
A	?	-	ILE	deletion	UNP Q07817
B	?	-	PHE	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	ASP	deletion	UNP Q07817
B	?	-	VAL	deletion	UNP Q07817
B	?	-	GLU	deletion	UNP Q07817
B	?	-	GLU	deletion	UNP Q07817
B	?	-	ASN	deletion	UNP Q07817
B	?	-	ARG	deletion	UNP Q07817
B	?	-	THR	deletion	UNP Q07817
B	?	-	GLU	deletion	UNP Q07817
B	?	-	ALA	deletion	UNP Q07817
B	?	-	PRO	deletion	UNP Q07817

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	GLU	deletion	UNP Q07817
B	?	-	GLY	deletion	UNP Q07817
B	?	-	THR	deletion	UNP Q07817
B	?	-	GLU	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	GLU	deletion	UNP Q07817
B	?	-	MET	deletion	UNP Q07817
B	?	-	GLU	deletion	UNP Q07817
B	?	-	THR	deletion	UNP Q07817
B	?	-	PRO	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	ALA	deletion	UNP Q07817
B	?	-	ILE	deletion	UNP Q07817
B	?	-	ASN	deletion	UNP Q07817
B	?	-	GLY	deletion	UNP Q07817
B	?	-	ASN	deletion	UNP Q07817
B	?	-	PRO	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	TRP	deletion	UNP Q07817
B	?	-	HIS	deletion	UNP Q07817
B	?	-	LEU	deletion	UNP Q07817
B	?	-	ALA	deletion	UNP Q07817
B	?	-	ASP	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	PRO	deletion	UNP Q07817
B	?	-	ALA	deletion	UNP Q07817
B	?	-	VAL	deletion	UNP Q07817
B	?	-	ASN	deletion	UNP Q07817
B	?	-	GLY	deletion	UNP Q07817
B	?	-	ALA	deletion	UNP Q07817
B	?	-	THR	deletion	UNP Q07817
B	?	-	GLY	deletion	UNP Q07817
B	?	-	HIS	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	SER	deletion	UNP Q07817
B	?	-	LEU	deletion	UNP Q07817
B	?	-	ASP	deletion	UNP Q07817
B	?	-	ALA	deletion	UNP Q07817
B	?	-	ARG	deletion	UNP Q07817
B	?	-	GLU	deletion	UNP Q07817
B	?	-	VAL	deletion	UNP Q07817

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ILE	deletion	UNP Q07817

- Molecule 2 is a protein called THR-PRO-GLU-TRP-ALA-SER-THR-LEU-ARG-PRO-GLU-(CCS)-(EDN)-(Z7Z)-(PRS).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	15	Total	C	N	O	S	0	0	0
			124	81	19	22	2			
2	C	15	Total	C	N	O	S	0	0	0
			124	81	19	22	2			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	88	Total	O	0	0
			88	88		
3	B	57	Total	O	0	0
			57	57		
3	D	5	Total	O	0	0
			5	5		
3	C	4	Total	O	0	0
			4	4		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bcl-2-like protein 1

Chain A: 



- Molecule 1: Bcl-2-like protein 1

Chain B: 




- Molecule 2: THR-PRO-GLU-TRP-ALA-SER-THR-LEU-ARG-PRO-GLU-(CCS)-(EDN)-(Z7Z)-(PRS)

Chain D: 



- Molecule 2: THR-PRO-GLU-TRP-ALA-SER-THR-LEU-ARG-PRO-GLU-(CCS)-(EDN)-(Z7Z)-(PRS)

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	64.02Å 97.61Å 93.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.80 – 1.95 48.80 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.2 (48.80-1.95) 99.3 (48.80-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.13_2998, PHENIX 1.13_2998	Depositor
R, R_{free}	0.181 , 0.226 0.183 , 0.226	Depositor DCC
R_{free} test set	1073 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.544	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2661	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Z7Z, EDN, CCS, PRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1157	0.57	0/1568
1	B	0.44	0/1163	0.53	0/1576
2	C	0.42	0/93	0.55	0/128
2	D	0.34	0/93	0.44	0/128
All	All	0.45	0/2506	0.55	0/3400

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1128	0	1073	9	0
1	B	1131	0	1078	14	0
2	C	124	0	97	2	0
2	D	124	0	97	0	0
3	A	88	0	0	0	0
3	B	57	0	0	1	0
3	C	4	0	0	0	0
3	D	5	0	0	0	0
All	All	2661	0	2345	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:9:ARG:HE	2:C:9:ARG:HA	1.41	0.85
1:A:12:PHE:CZ	1:A:16:LYS:HE3	2.24	0.72
1:B:3:GLN:HA	3:B:201:HOH:O	1.95	0.66
1:A:96:GLU:OE2	1:A:100:ARG:HD3	1.96	0.66
1:B:10:VAL:HG12	1:B:26:GLN:HG2	1.82	0.62
2:C:9:ARG:HA	2:C:9:ARG:NE	2.14	0.60
1:A:83:MET:SD	1:B:3:GLN:HB3	2.41	0.60
1:A:6:ARG:NH2	1:B:7:GLU:OE2	2.35	0.59
1:B:114:ILE:O	1:B:114:ILE:HD12	2.02	0.59
1:B:129:GLU:OE1	1:B:132:ARG:HD2	2.05	0.56
1:A:3:GLN:NE2	1:B:189:ASP:OD1	2.44	0.47
1:B:96:GLU:OE1	1:B:100:ARG:NH1	2.47	0.46
1:A:3:GLN:HG2	1:B:83:MET:HG3	1.98	0.45
1:B:4:SER:HB3	1:B:7:GLU:OE1	2.18	0.44
1:A:3:GLN:HG2	1:B:83:MET:CG	2.48	0.44
1:B:179:GLU:OE2	1:B:183:GLN:HG2	2.17	0.43
1:A:8:LEU:HD11	1:B:86:VAL:HG12	2.02	0.42
1:A:6:ARG:HH22	1:B:7:GLU:CD	2.24	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/141 (97%)	135 (98%)	2 (2%)	0	100	100
1	B	138/141 (98%)	136 (99%)	1 (1%)	1 (1%)	19	10
2	C	10/15 (67%)	10 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	10/15 (67%)	10 (100%)	0	0	100	100
All	All	295/312 (95%)	291 (99%)	3 (1%)	1 (0%)	37	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	25	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/120 (98%)	118 (100%)	0	100	100
1	B	119/120 (99%)	119 (100%)	0	100	100
2	C	10/10 (100%)	9 (90%)	1 (10%)	6	1
2	D	10/10 (100%)	10 (100%)	0	100	100
All	All	257/260 (99%)	256 (100%)	1 (0%)	89	89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	9	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CCS	C	12	2	7,8,10	0.68	0	3,8,12	0.59	0
2	PRS	D	15	2	5,7,8	3.50	2 (40%)	3,8,10	6.26	2 (66%)
2	PRS	C	15	2	5,7,8	3.42	2 (40%)	3,8,10	6.17	2 (66%)
2	CCS	D	12	2	7,8,10	0.72	0	3,8,12	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CCS	C	12	2	-	0/4/7/10	-
2	PRS	D	15	2	-	0/0/9/11	0/1/1/1
2	PRS	C	15	2	-	0/0/9/11	0/1/1/1
2	CCS	D	12	2	-	0/4/7/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	15	PRS	CD-N	-7.02	1.27	1.45
2	C	15	PRS	CD-N	-6.97	1.28	1.45
2	D	15	PRS	CD-SG	-3.46	1.76	1.81
2	C	15	PRS	CD-SG	-3.14	1.76	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	15	PRS	SG-CD-N	10.47	116.53	105.72
2	C	15	PRS	SG-CD-N	10.46	116.52	105.72
2	D	15	PRS	O-C-CA	-2.36	118.59	124.78
2	C	15	PRS	O-C-CA	-2.00	119.53	124.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	139/141 (98%)	-0.07	1 (0%) 84 87	16, 26, 41, 56	0
1	B	139/141 (98%)	0.21	3 (2%) 62 68	17, 30, 46, 53	1 (0%)
2	C	11/15 (73%)	0.14	0 100 100	26, 30, 53, 55	0
2	D	11/15 (73%)	0.37	0 100 100	29, 35, 54, 57	0
All	All	300/312 (96%)	0.08	4 (1%) 74 79	16, 28, 47, 57	1 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	GLY	2.3
1	B	196	GLY	2.2
1	B	114	ILE	2.2
1	B	3	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CCS	D	12	9/11	0.82	0.13	43,47,54,56	0
2	CCS	C	12	9/11	0.92	0.09	32,39,45,48	0
2	PRS	D	15	7/8	0.96	0.06	30,31,34,36	0
2	PRS	C	15	7/8	0.97	0.06	27,29,32,36	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.