



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2025 – 02:40 PM JST

PDB ID : 8ZML / pdb_00008zml
Title : Structure of TNIK with inhibitor
Authors : Fan, Y.
Deposited on : 2024-05-23
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7106 (2.50-2.46)
Clashscore	180529	7991 (2.50-2.46)
Ramachandran outliers	177936	7888 (2.50-2.46)
Sidechain outliers	177891	7890 (2.50-2.46)
RSRZ outliers	164620	7106 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	303	7% 96% ..
1	B	303	11% 93% 5% ..

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRAF2 and NCK-interacting protein kinase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	301	Total 2457	C 1553	N 434	O 451	P 2	S 17	0	4	0
1	B	300	Total 2461	C 1555	N 434	O 453	P 2	S 17	0	6	0

- Molecule 2 is Rentosertib (CCD ID: A1D8H) (formula: $C_{27}H_{30}FN_7O$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 36	C 27	F 1	N 7	O 1	0	0
2	B	1	Total 36	C 27	F 1	N 7	O 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	77	Total	O	0	0
			77	77		
5	B	67	Total	O	0	0
			67	67		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TRAF2 and NCK-interacting protein kinase



- Molecule 1: TRAF2 and NCK-interacting protein kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.89Å 89.59Å 97.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	66.00 – 2.49 66.00 – 2.49	Depositor EDS
% Data completeness (in resolution range)	99.9 (66.00-2.49) 99.9 (66.00-2.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.256 , 0.326 0.257 , 0.323	Depositor DCC
R_{free} test set	1315 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	58.5	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5144	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1D8H, EDO, TPO, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	0/2485	1.45	0/3346
1	B	1.01	0/2492	1.46	2/3358 (0.1%)
All	All	1.01	0/4977	1.46	2/6704 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	CYS	CA-C-N	5.18	125.84	119.99
1	B	108	CYS	C-N-CA	5.18	125.84	119.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2457	0	2460	5	0
1	B	2461	0	2457	5	0
2	A	36	0	0	0	0
2	B	36	0	0	0	0
3	A	8	0	12	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	77	0	0	3	0
5	B	67	0	0	0	0
All	All	5144	0	4929	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:ARG:O	5:A:501:HOH:O	2.11	0.68
1:A:244:LEU:N	5:A:501:HOH:O	2.25	0.59
1:B:145:HIS:CD2	1:B:212:PHE:HB2	2.43	0.53
1:B:292:GLN:N	1:B:293:PRO:CD	2.74	0.51
1:A:188:PHE:CE1	1:A:195:MET:HE1	2.51	0.44
1:A:245:ILE:HB	1:A:246:PRO:HD3	2.00	0.44
1:B:289:ILE:O	1:B:290:ARG:C	2.62	0.42
1:A:17:ALA:N	5:A:502:HOH:O	2.54	0.41
1:B:160:LEU:HD22	1:B:170:VAL:HG11	2.02	0.41
1:B:284[A]:MET:HE2	1:B:284[A]:MET:HB2	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/303 (99%)	284 (95%)	13 (4%)	2 (1%)	19	32
1	B	300/303 (99%)	282 (94%)	16 (5%)	2 (1%)	19	32
All	All	599/606 (99%)	566 (94%)	29 (5%)	4 (1%)	19	32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	290	ARG
1	A	35	THR
1	B	274	HIS
1	A	297	GLN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/263 (101%)	262 (99%)	3 (1%)	70	86
1	B	266/263 (101%)	259 (97%)	7 (3%)	41	65
All	All	531/526 (101%)	521 (98%)	10 (2%)	52	74

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	36	TYR
1	A	297	GLN
1	B	10	LEU
1	B	46	LYS
1	B	65	GLU
1	B	256	LYS
1	B	257	LYS
1	B	292	GLN
1	B	299	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	GLN
1	B	143	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	B	181	1	8,10,11	0.74	0	10,14,16	0.84	0
1	TPO	A	187	1	8,10,11	0.70	0	10,14,16	0.79	0
1	TPO	A	181	1	8,10,11	0.74	0	10,14,16	0.81	0
1	TPO	B	187	1	8,10,11	0.67	0	10,14,16	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	B	181	1	-	1/9/11/13	-
1	TPO	A	187	1	-	2/9/11/13	-
1	TPO	A	181	1	-	1/9/11/13	-
1	TPO	B	187	1	-	0/9/11/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	181	TPO	O-C-CA-CB
1	A	187	TPO	CB-OG1-P-O3P
1	B	181	TPO	O-C-CA-CB

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	187	TPO	CB-OG1-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	A1D8H	A	401	-	36,40,40	2.37	10 (27%)	43,57,57	1.28	5 (11%)
2	A1D8H	B	401	-	36,40,40	2.36	10 (27%)	43,57,57	1.18	3 (6%)
3	EDO	A	402	-	3,3,3	0.06	0	2,2,2	0.18	0
3	EDO	A	403	-	3,3,3	0.07	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1D8H	A	401	-	-	1/16/34/34	0/5/5/5
2	A1D8H	B	401	-	-	4/16/34/34	0/5/5/5
3	EDO	A	402	-	-	1/1/1/1	-
3	EDO	A	403	-	-	1/1/1/1	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	A1D8H	C12-N11	-7.23	1.30	1.46
2	A	401	A1D8H	C12-N11	-7.09	1.30	1.46
2	A	401	A1D8H	C29-C28	5.49	1.55	1.49
2	B	401	A1D8H	C29-C28	5.19	1.55	1.49
2	A	401	A1D8H	C02-N03	5.11	1.49	1.35
2	B	401	A1D8H	C02-N03	5.05	1.49	1.35
2	B	401	A1D8H	C04-N03	4.05	1.49	1.41
2	A	401	A1D8H	C04-N03	3.97	1.49	1.41
2	A	401	A1D8H	C07-N08	3.76	1.49	1.38
2	A	401	A1D8H	C13-N11	-3.74	1.38	1.46
2	B	401	A1D8H	C13-N11	-3.72	1.38	1.46
2	B	401	A1D8H	C10-N11	-3.68	1.38	1.46
2	A	401	A1D8H	C10-N11	-3.67	1.38	1.46
2	B	401	A1D8H	C07-N08	3.53	1.48	1.38
2	B	401	A1D8H	C21-C28	-2.65	1.37	1.43
2	A	401	A1D8H	C21-C28	-2.62	1.37	1.43
2	B	401	A1D8H	C17-C02	2.36	1.56	1.50
2	A	401	A1D8H	O01-C02	-2.18	1.18	1.23
2	A	401	A1D8H	C17-C02	2.17	1.55	1.50
2	B	401	A1D8H	O01-C02	-2.14	1.19	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	A1D8H	C26-N27-C28	2.87	108.47	102.99
2	B	401	A1D8H	C13-N11-C10	2.84	113.49	109.52
2	B	401	A1D8H	C26-N27-C28	2.83	108.38	102.99
2	A	401	A1D8H	C12-N11-C13	2.82	114.88	110.66
2	A	401	A1D8H	C09-C10-N11	2.62	113.77	110.80
2	A	401	A1D8H	C12-N11-C10	2.62	114.58	110.66
2	A	401	A1D8H	C13-N11-C10	2.17	112.55	109.52
2	B	401	A1D8H	C34-C32-C31	-2.01	120.15	122.83

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	EDO	O1-C1-C2-O2
3	A	403	EDO	O1-C1-C2-O2
2	B	401	A1D8H	C06-C07-N08-C14
2	B	401	A1D8H	C15-C07-N08-C14
2	B	401	A1D8H	N27-C28-C29-C30

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	401	A1D8H	N27-C28-C29-C35
2	B	401	A1D8H	N27-C28-C29-C35

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	299/303 (98%)	0.75	21 (7%)	24	22	22, 63, 106, 133	4 (1%)
1	B	298/303 (98%)	0.94	33 (11%)	12	11	24, 68, 113, 131	6 (2%)
All	All	597/606 (98%)	0.84	54 (9%)	17	16	22, 66, 109, 133	10 (1%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	298	VAL	6.1
1	B	293	PRO	4.5
1	B	36	TYR	4.3
1	A	295	GLU	4.3
1	B	146[A]	GLN	4.3
1	A	36	TYR	4.2
1	B	289	ILE	4.0
1	A	284[A]	MET	4.0
1	B	298	VAL	3.7
1	A	300	ILE	3.7
1	B	137	ILE	3.4
1	A	302	LEU	3.3
1	B	284[A]	MET	3.2
1	B	306	ILE	3.1
1	B	291	ASP	3.0
1	B	300	ILE	3.0
1	B	60	GLY	2.9
1	B	133	ILE	2.9
1	A	301	GLN	2.9
1	B	258	TRP	2.9
1	B	130	ILE	2.8
1	A	299	ARG	2.7
1	A	293	PRO	2.7
1	B	283	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	125	LEU	2.7
1	B	294	ASN	2.7
1	B	223	ALA	2.6
1	B	287	PRO	2.5
1	B	288	PHE	2.5
1	B	227	ALA	2.5
1	A	309	THR	2.4
1	B	134	CYS	2.4
1	B	55	VAL	2.4
1	A	94	PRO	2.4
1	B	253	LEU	2.3
1	A	95	PRO	2.3
1	A	60	GLY	2.3
1	A	182	VAL	2.3
1	B	305	HIS	2.3
1	B	131	ALA	2.3
1	A	306	ILE	2.2
1	A	305	HIS	2.2
1	B	61	ASP	2.2
1	A	220	GLY	2.2
1	B	37	GLY	2.2
1	B	239[A]	MET	2.2
1	B	290	ARG	2.2
1	A	157	GLN	2.1
1	B	302	LEU	2.1
1	A	129	TRP	2.1
1	A	296	ARG	2.1
1	B	296	ARG	2.1
1	B	265	PHE	2.1
1	A	15	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	B	181	11/12	0.78	0.12	86,87,91,91	0
1	TPO	A	181	11/12	0.83	0.10	80,82,84,85	0
1	TPO	B	187	11/12	0.86	0.12	61,61,62,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	TPO	A	187	11/12	0.94	0.08	50,50,51,52	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	403	4/4	0.83	0.16	71,71,71,71	0
3	EDO	A	402	4/4	0.85	0.13	69,69,69,70	0
2	A1D8H	B	401	36/36	0.90	0.12	42,47,67,68	0
2	A1D8H	A	401	36/36	0.93	0.09	42,46,58,59	0
4	MG	B	402	1/1	0.93	0.07	54,54,54,54	0
4	MG	A	404	1/1	0.98	0.05	56,56,56,56	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1D8H B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1D8H A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.