



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2024 – 12:26 PM EDT

PDB ID : 9BLG  
Title : Crystal structure of non-receptor protein tyrosine phosphatase SHP2 in complex with PF-07284892  
Authors : Bester, S.M.; Wu, W.-I.; Mou, T.-C.  
Deposited on : 2024-04-30  
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	539	 7% 85% 7% • 7%
1	B	539	 8% 85% 8% • 7%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 8645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	499	4047	2546	720	763	18	0	3	0
1	B	503	4058	2548	722	770	18	0	0	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP Q06124
A	-12	HIS	-	expression tag	UNP Q06124
A	-11	HIS	-	expression tag	UNP Q06124
A	-10	HIS	-	expression tag	UNP Q06124
A	-9	HIS	-	expression tag	UNP Q06124
A	-8	HIS	-	expression tag	UNP Q06124
A	-7	HIS	-	expression tag	UNP Q06124
A	-6	GLU	-	expression tag	UNP Q06124
A	-5	ASN	-	expression tag	UNP Q06124
A	-4	LEU	-	expression tag	UNP Q06124
A	-3	TYR	-	expression tag	UNP Q06124
A	-2	PHE	-	expression tag	UNP Q06124
A	-1	GLN	-	expression tag	UNP Q06124
A	0	SER	-	expression tag	UNP Q06124
B	-13	MET	-	initiating methionine	UNP Q06124
B	-12	HIS	-	expression tag	UNP Q06124
B	-11	HIS	-	expression tag	UNP Q06124
B	-10	HIS	-	expression tag	UNP Q06124
B	-9	HIS	-	expression tag	UNP Q06124
B	-8	HIS	-	expression tag	UNP Q06124
B	-7	HIS	-	expression tag	UNP Q06124
B	-6	GLU	-	expression tag	UNP Q06124
B	-5	ASN	-	expression tag	UNP Q06124
B	-4	LEU	-	expression tag	UNP Q06124
B	-3	TYR	-	expression tag	UNP Q06124

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	PHE	-	expression tag	UNP Q06124
B	-1	GLN	-	expression tag	UNP Q06124
B	0	SER	-	expression tag	UNP Q06124

- Molecule 2 is (1S)-1'-{6-[(2-amino-3-chloropyridin-4-yl)sulfanyl]-1,2,4-triazin-3-yl}-1,3-dihydrospiro[indene-2,4'-piperidin]-1-amine (three-letter code: A1AQ1) (formula: C<sub>21</sub>H<sub>22</sub>ClN<sub>7</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			S
2	A	1	Total	C	Cl	N	S	0	0
			30	21	1	7	1		
2	B	1	Total	C	Cl	N	S	0	0
			30	21	1	7	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	213	Total	O	0	0
			213	213		
3	B	267	Total	O	0	0
			267	267		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.27Å 214.27Å 55.81Å 90.00° 96.06° 90.00°	Depositor
Resolution (Å)	28.27 – 2.06 28.27 – 2.06	Depositor EDS
% Data completeness (in resolution range)	94.3 (28.27-2.06) 94.4 (28.27-2.06)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.85 (at 2.06Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, $R_{free}$	0.188 , 0.238 0.187 , 0.238	Depositor DCC
$R_{free}$ test set	3054 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.5	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8645	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1AQ1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/4140	0.49	0/5584
1	B	0.24	0/4141	0.48	0/5584
All	All	0.25	0/8281	0.49	0/11168

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4047	0	3981	20	0
1	B	4058	0	3979	22	0
2	A	30	0	0	0	0
2	B	30	0	0	0	0
3	A	213	0	0	2	0
3	B	267	0	0	1	0
All	All	8645	0	7960	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:ARG:HD2	1:A:96:ILE:HD13	1.44	1.00
1:A:226:ILE:HD11	1:A:488:ILE:HD11	1.69	0.73
1:A:221:ILE:HD11	1:A:226:ILE:HD13	1.80	0.62
1:A:274:LYS:NZ	3:A:704:HOH:O	2.30	0.61
1:B:388:VAL:HG21	1:B:402:LYS:HE2	1.85	0.58
1:A:107:PRO:HG3	1:A:190:LEU:HD12	1.85	0.58
1:A:2:THR:N	3:A:709:HOH:O	2.38	0.57
1:B:107:PRO:HG3	1:B:190:LEU:HD12	1.87	0.57
1:B:133:GLY:HA3	1:B:213:LYS:HB2	1.87	0.56
1:A:79:GLN:O	1:A:83:GLU:HG2	2.06	0.54
1:B:290:VAL:HG11	1:B:344:MET:HG3	1.93	0.51
1:B:203:VAL:HG22	1:B:209:VAL:HG22	1.94	0.50
1:A:88:LEU:HG	1:A:96:ILE:CG2	2.42	0.49
1:B:65:LEU:HD23	1:B:68:GLY:HA3	1.96	0.48
1:B:274:LYS:HE2	1:B:274:LYS:HB3	1.56	0.48
1:A:290:VAL:HG11	1:A:344:MET:HG3	1.97	0.47
1:B:4:ARG:HD3	1:B:256:GLN:HA	1.96	0.47
1:A:96:ILE:HG13	1:A:97:GLU:N	2.30	0.47
1:B:272:GLU:HG3	1:B:301:VAL:HG21	1.97	0.47
1:A:228:SER:O	1:A:232:GLU:HG3	2.15	0.46
1:B:55:LYS:HD3	1:B:66:TYR:CE2	2.51	0.46
1:B:497:VAL:HG12	1:B:504:MET:HG3	1.96	0.46
1:A:418:TYR:HB3	1:A:438:PHE:CE1	2.51	0.45
1:B:20:LEU:HD23	1:B:102:LEU:HD12	1.97	0.45
1:A:28:SER:HA	1:A:100:TYR:O	2.18	0.44
1:A:46:ARG:C	1:A:47:ARG:HG3	2.37	0.44
1:B:287:HIS:CD2	1:B:288:THR:HG23	2.53	0.44
1:A:133:GLY:HA3	1:A:213:LYS:HB2	2.00	0.44
1:B:226:ILE:HD11	1:B:515:TYR:HD1	1.83	0.43
1:B:79:GLN:O	1:B:83:GLU:HG3	2.18	0.43
1:B:28:SER:HA	1:B:100:TYR:O	2.19	0.43
1:A:125:LEU:HB3	1:A:216:LEU:HD21	2.00	0.43
1:B:146:ASP:HB3	3:B:701:HOH:O	2.19	0.43
1:B:418:TYR:HB3	1:B:438:PHE:CE1	2.54	0.42
1:B:525:LEU:HD12	1:B:525:LEU:HA	1.88	0.42
1:A:125:LEU:O	1:A:129:LYS:HG2	2.20	0.42
1:A:65:LEU:HD22	1:A:68:GLY:HA3	2.02	0.41
1:B:125:LEU:HD22	1:B:129:LYS:HE3	2.03	0.41
1:A:278:ARG:NH2	1:A:333:CYS:O	2.54	0.40
1:A:388:VAL:HG21	1:A:402:LYS:HE3	2.03	0.40
1:B:94:ASP:OD1	1:B:94:ASP:N	2.54	0.40

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:439:LEU:HD23	1:B:439:LEU:HA	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/539 (91%)	480 (98%)	11 (2%)	1 (0%)	47	39
1	B	495/539 (92%)	482 (97%)	12 (2%)	1 (0%)	47	39
All	All	987/1078 (92%)	962 (98%)	23 (2%)	2 (0%)	47	39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	505	VAL
1	B	505	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	446/481 (93%)	435 (98%)	11 (2%)	47	41
1	B	445/481 (92%)	437 (98%)	8 (2%)	59	55
All	All	891/962 (93%)	872 (98%)	19 (2%)	53	48

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	47	ARG
1	A	129	LYS
1	A	164	LYS
1	A	199	LYS
1	A	211	GLN
1	A	235	LYS
1	A	244	LYS
1	A	364	LYS
1	A	413	ARG
1	A	481	GLU
1	B	5	ARG
1	B	19	LEU
1	B	94	ASP
1	B	160	SER
1	B	205	THR
1	B	272	GLU
1	B	274	LYS
1	B	413	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	B	103	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1AQ1	A	601	-	32,34,34	1.09	2 (6%)	38,50,50	1.71	10 (26%)
2	A1AQ1	B	601	-	32,34,34	1.07	2 (6%)	38,50,50	1.73	9 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1AQ1	A	601	-	-	2/8/35/35	0/5/5/5
2	A1AQ1	B	601	-	-	2/8/35/35	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	A1AQ1	C27-N29	2.44	1.40	1.34
2	A	601	A1AQ1	C5-N8	2.42	1.40	1.35
2	B	601	A1AQ1	C27-N29	2.37	1.40	1.34
2	B	601	A1AQ1	C5-N8	2.29	1.40	1.35

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	A1AQ1	C27-C28-CL1	4.04	120.89	118.92
2	B	601	A1AQ1	C28-C27-N29	-3.98	120.61	122.94
2	A	601	A1AQ1	C5-N4-N3	3.92	119.08	117.09
2	A	601	A1AQ1	C28-C27-N29	-3.91	120.65	122.94
2	B	601	A1AQ1	C28-C27-N26	3.72	122.41	119.30
2	A	601	A1AQ1	C28-C27-N26	3.71	122.41	119.30
2	B	601	A1AQ1	C5-N4-N3	3.67	118.96	117.09

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	A1AQ1	C27-C28-CL1	3.66	120.70	118.92
2	B	601	A1AQ1	C15-C16-C17	2.97	113.17	110.62
2	A	601	A1AQ1	C15-C16-C17	2.89	113.11	110.62
2	B	601	A1AQ1	C14-C15-C16	2.67	113.28	110.61
2	A	601	A1AQ1	C14-C15-C16	2.67	113.28	110.61
2	A	601	A1AQ1	C21-C22-C16	-2.44	117.91	121.01
2	B	601	A1AQ1	C21-C22-C16	-2.41	117.95	121.01
2	A	601	A1AQ1	C1-C2-N3	-2.37	121.03	122.20
2	B	601	A1AQ1	C14-C15-C19	-2.27	125.41	129.72
2	A	601	A1AQ1	C22-C16-C17	-2.23	125.51	129.45
2	B	601	A1AQ1	C22-C16-C17	-2.18	125.60	129.45
2	A	601	A1AQ1	C14-C15-C19	-2.17	125.60	129.72

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	A1AQ1	C1-C2-S7-C23
2	B	601	A1AQ1	C1-C2-S7-C23
2	B	601	A1AQ1	N3-C2-S7-C23
2	A	601	A1AQ1	N3-C2-S7-C23

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	499/539 (92%)	0.31	39 (7%) <b>13</b>   <b>13</b>	11, 25, 48, 71	0
1	B	503/539 (93%)	0.28	41 (8%) <b>11</b>   <b>12</b>	11, 24, 48, 68	0
All	All	1002/1078 (92%)	0.30	80 (7%) <b>12</b>   <b>12</b>	11, 24, 48, 71	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	158	GLY	5.0
1	A	85	HIS	4.7
1	B	159	GLU	4.7
1	A	409	GLY	4.4
1	B	157	LYS	4.4
1	A	2	THR	4.4
1	A	300	PRO	4.4
1	A	116	HIS	4.3
1	A	298	ASN	4.0
1	B	298	ASN	4.0
1	B	116	HIS	3.9
1	A	93	GLY	3.9
1	A	294	ASP	3.8
1	B	91	LYS	3.6
1	B	160	SER	3.5
1	A	144	PRO	3.5
1	B	92	ASN	3.4
1	B	457	VAL	3.4
1	A	297	PRO	3.4
1	B	93	GLY	3.3
1	A	296	ASP	3.3
1	B	85	HIS	3.1
1	B	235	LYS	3.1
1	A	48	ASN	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	141	GLN	2.9
1	A	523	GLU	2.9
1	A	205	THR	2.9
1	B	409	GLY	2.9
1	B	525	LEU	2.8
1	A	142	SER	2.8
1	A	177	LEU	2.7
1	A	38	PRO	2.7
1	B	299	GLU	2.7
1	B	199	LYS	2.7
1	A	364	LYS	2.7
1	A	299	GLU	2.6
1	A	301	VAL	2.6
1	A	363	GLY	2.6
1	B	86	GLY	2.6
1	B	300	PRO	2.6
1	A	235	LYS	2.6
1	A	161	ASN	2.5
1	A	244	LYS	2.5
1	B	38	PRO	2.5
1	B	313	GLU	2.5
1	B	470	ILE	2.5
1	A	143	HIS	2.5
1	A	18	ASN	2.5
1	B	143	HIS	2.5
1	A	47	ARG	2.5
1	B	177	LEU	2.5
1	B	205	THR	2.5
1	B	95	VAL	2.4
1	A	86	GLY	2.4
1	A	362	ARG	2.4
1	A	37	ASN	2.3
1	A	89	LYS	2.2
1	A	199	LYS	2.2
1	B	14	VAL	2.2
1	B	244	LYS	2.2
1	B	324	LYS	2.2
1	B	481	GLU	2.2
1	B	524	THR	2.2
1	B	469	PHE	2.2
1	B	505	VAL	2.2
1	B	144	PRO	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	471	VAL	2.2
1	A	83	GLU	2.1
1	B	94	ASP	2.1
1	B	141	GLN	2.1
1	A	451	ASP	2.1
1	A	36	SER	2.1
1	B	466	THR	2.1
1	A	295	GLY	2.1
1	B	494	ILE	2.1
1	B	296	ASP	2.1
1	B	254	LEU	2.0
1	B	354	VAL	2.0
1	A	426	HIS	2.0
1	A	35	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	A1AQ1	A	601	30/30	0.95	0.11	18,22,29,32	0
2	A1AQ1	B	601	30/30	0.95	0.14	17,21,24,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around A1AQ1 A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A1AQ1 B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.