



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 26, 2024 – 10:54 pm BST

PDB ID : 9F6C  
Title : Cardiac myosin motor domain in the pre-powerstroke state co-crystallized with the inhibitor aficamten  
Authors : Robert-Paganin, J.; Hartman, J.J.; Morgan, B.P.; Malik, F.I.; Houdusse, A.  
Deposited on : 2024-05-01  
Resolution : 2.33 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2747 (2.36-2.32)
Clashscore	180529	2936 (2.36-2.32)
Ramachandran outliers	177936	2912 (2.36-2.32)
Sidechain outliers	177891	2912 (2.36-2.32)
RSRZ outliers	164620	2747 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	782	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 81%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey;"></div> </div>
1	B	782	<div style="display: flex; align-items: center;"> <div style="width: 15%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 81%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	VO4	A	1005	-	-	X	-
6	VO4	B	804	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 11944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	717	Total 5794	C 3708	N 984	O 1072	S 30	1	2	0
1	B	706	Total 5699	C 3650	N 962	O 1056	S 31	1	1	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	1	Total 1	Mg 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is aficanten (three-letter code: 6I6) (formula:  $C_{18}H_{19}N_5O_2$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			25	18	5	2		
4	B	1	Total	C	N	O	0	0
			25	18	5	2		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
5	A	1	27	10	5	10	2	0	0
5	B	1	27	10	5	10	2	0	0

- Molecule 6 is VANADATE ION (three-letter code: VO4) (formula:  $O_4V$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O V		
6	A	1	5	4 1	0	0
6	B	1	5	4 1	0	0


- Molecule 7 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
7	A	196	Total 196	O 196	0	0
7	B	134	Total 134	O 134	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Myosin-7



- Molecule 1: Myosin-7







## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.42Å 128.33Å 103.53Å 90.00° 91.47° 90.00°	Depositor
Resolution (Å)	46.60 – 2.33 46.60 – 2.33	Depositor EDS
% Data completeness (in resolution range)	67.2 (46.60-2.33) 67.3 (46.60-2.33)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.34Å)	Xtrriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.205 , 0.244 0.197 , 0.234	Depositor DCC
$R_{free}$ test set	2562 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.3	Xtrriage
Anisotropy	0.020	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 60.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.056 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	11944	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, M3L, ADP, 6I6, VO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/5890	0.70	0/7939
1	B	0.50	0/5793	0.69	0/7807
All	All	0.51	0/11683	0.69	0/15746

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5794	0	5778	38	0
1	B	5699	0	5671	34	0
2	A	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	25	0	0	2	0
4	B	25	0	0	1	0
5	A	27	0	12	0	0
5	B	27	0	12	0	0
6	A	5	0	0	2	0
6	B	5	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	196	0	0	2	0
7	B	134	0	0	0	0
All	All	11944	0	11473	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:504:GLU:HG2	1:B:764:PHE:HE1	1.49	0.78
1:B:504:GLU:HG2	1:B:764:PHE:CE1	2.31	0.66
1:B:339:LEU:O	1:B:446:THR:HG21	1.97	0.64
6:B:804:VO4:O1	6:B:804:VO4:V	1.56	0.62
6:B:804:VO4:V	6:B:804:VO4:O2	1.57	0.60
6:A:1005:VO4:O2	6:A:1005:VO4:V	1.59	0.59
1:B:544:THR:HG22	1:B:547:THR:HG23	1.83	0.59
1:B:435:MET:HG3	1:B:616:MET:HE2	1.85	0.58
1:B:356:ILE:HD11	1:B:431:VAL:HG22	1.85	0.58
1:A:730:ILE:HD11	1:A:745:LEU:HD12	1.85	0.58
6:B:804:VO4:V	6:B:804:VO4:O4	1.59	0.57
6:A:1005:VO4:V	6:A:1005:VO4:O4	1.59	0.57
1:A:356:ILE:HD11	1:A:431:VAL:HG22	1.88	0.56
1:B:245:GLY:HA3	4:B:802:6I6:N1	2.21	0.56
1:A:559:LYS:HD3	1:B:55:GLU:CG	2.36	0.55
1:B:656:ASN:O	1:B:660:THR:HG23	2.06	0.54
1:B:47:VAL:HG22	1:B:64:GLU:HG2	1.88	0.54
1:B:84:PHE:HD1	1:B:87:ILE:HD12	1.72	0.54
1:B:301:LEU:HB2	1:B:303:ILE:HG12	1.89	0.54
1:A:84:PHE:HD1	1:A:87:ILE:HD12	1.73	0.53
1:A:301:LEU:HD22	1:A:351:LYS:HA	1.91	0.53
1:B:724:ILE:HG22	1:B:780:ARG:HG2	1.91	0.53
1:B:301:LEU:HD22	1:B:351:LYS:HA	1.91	0.53
1:A:403:ARG:HG2	1:A:410:TYR:CD1	2.44	0.52
1:B:108:ARG:NH1	1:B:113:MET:HB3	2.25	0.52
1:A:301:LEU:HB2	1:A:303:ILE:HG12	1.90	0.52
1:A:656:ASN:O	1:A:660:THR:HG23	2.08	0.52
1:A:192:ILE:HD11	1:A:248:ILE:HD13	1.92	0.52
1:A:245:GLY:HA3	4:A:1003:6I6:N1	2.24	0.52
1:B:403:ARG:HG2	1:B:410:TYR:CD1	2.45	0.52
1:A:765:PHE:HD1	1:A:769:LEU:HD23	1.74	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:GLU:HG2	1:A:108:ARG:HH21	1.76	0.51
1:B:88:GLU:HG2	1:B:108:ARG:HH21	1.75	0.51
1:B:39:VAL:HG11	1:B:69:VAL:HG11	1.93	0.50
1:B:765:PHE:HD1	1:B:769:LEU:HD23	1.77	0.50
1:B:39:VAL:HG13	1:B:40:PRO:HD2	1.94	0.49
1:A:97:HIS:CE1	1:A:100:ALA:HB2	2.48	0.48
1:A:443:ILE:O	1:A:446:THR:HG22	2.14	0.48
1:A:559:LYS:HD3	1:B:55:GLU:HG2	1.95	0.48
1:B:97:HIS:CE1	1:B:100:ALA:HB2	2.49	0.47
1:B:81:PRO:HD2	1:B:84:PHE:HD2	1.79	0.47
1:A:108:ARG:HG2	1:A:113:MET:HE2	1.97	0.47
1:A:564[B]:GLN:HG3	1:A:578:SER:OG	2.14	0.46
1:A:684:ILE:HG22	1:A:686:ASN:HD22	1.80	0.46
1:A:39:VAL:HG11	1:A:69:VAL:HG11	1.98	0.46
1:B:467:ILE:HG22	1:B:586:VAL:HG22	1.98	0.46
1:A:730:ILE:CD1	1:A:745:LEU:HD12	2.46	0.45
1:A:33:LEU:HD12	1:A:34:LYS:H	1.82	0.45
1:A:753:HIS:HA	1:A:756:TYR:CE2	2.51	0.45
1:A:81:PRO:HD2	1:A:84:PHE:HD2	1.82	0.45
1:B:251:HIS:HB3	1:B:453:ARG:HG2	1.98	0.45
1:A:474:GLU:HA	4:A:1003:6I6:O1	2.18	0.45
1:B:243:ARG:HD3	1:B:269:GLU:OE2	2.16	0.44
1:A:251:HIS:HB3	1:A:453:ARG:HG2	2.00	0.44
1:A:544:THR:HG22	1:A:547:THR:HG23	2.00	0.44
1:B:684:ILE:HG22	1:B:686:ASN:HD22	1.82	0.43
1:A:417:VAL:O	1:A:421:VAL:HG13	2.18	0.43
1:A:92:MET:HG2	7:A:1287:HOH:O	2.17	0.43
1:A:624:TYR:HD2	7:A:1290:HOH:O	2.01	0.43
1:A:136:ALA:HA	1:A:139:VAL:HG13	2.00	0.43
1:A:108:ARG:NH1	1:A:113:MET:HB3	2.34	0.43
1:A:723:ARG:HG3	1:A:730:ILE:HD12	2.00	0.43
1:B:228:GLU:O	1:B:232:ASN:HB2	2.20	0.42
1:B:81:PRO:HD2	1:B:84:PHE:CD2	2.54	0.42
1:A:435:MET:HG3	1:A:616:MET:HE2	2.02	0.42
1:A:39:VAL:HG13	1:A:40:PRO:HD2	2.02	0.41
1:A:249:ARG:HD3	1:A:262:ASP:OD1	2.21	0.41
1:B:185:THR:HG22	1:B:189:LYS:HE2	2.02	0.41
1:B:136:ALA:HA	1:B:139:VAL:HG13	2.02	0.41
1:B:249:ARG:HD3	1:B:262:ASP:OD1	2.21	0.41
1:A:616:MET:O	1:A:620:LEU:HG	2.21	0.41
1:B:361:ASN:HB3	1:B:380:GLU:HG2	2.02	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:GLU:O	1:B:220:ILE:HG22	2.21	0.40
1:A:228:GLU:O	1:A:232:ASN:HB2	2.21	0.40
1:A:114:ILE:O	1:A:124:THR:HA	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	707/782 (90%)	686 (97%)	20 (3%)	1 (0%)	48 58
1	B	693/782 (89%)	677 (98%)	15 (2%)	1 (0%)	48 58
All	All	1400/1564 (90%)	1363 (97%)	35 (2%)	2 (0%)	48 58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	267	LEU
1	B	267	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	624/672 (93%)	601 (96%)	23 (4%)	29 37

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	614/672 (91%)	597 (97%)	17 (3%)	38 48
All	All	1238/1344 (92%)	1198 (97%)	40 (3%)	34 42

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	VAL
1	A	108	ARG
1	A	211	THR
1	A	255	THR
1	A	257	LYS
1	A	359	PHE
1	A	379	GLU
1	A	405	LYS
1	A	406	VAL
1	A	412	THR
1	A	418	GLN
1	A	451	GLN
1	A	453	ARG
1	A	454	GLN
1	A	455	TYR
1	A	484	LYS
1	A	510	PHE
1	A	569	ILE
1	A	605	VAL
1	A	666	HIS
1	A	671	ARG
1	A	721	ARG
1	A	736	ILE
1	B	34	LYS
1	B	69	VAL
1	B	108	ARG
1	B	255	THR
1	B	359	PHE
1	B	379	GLU
1	B	406	VAL
1	B	412	THR
1	B	418	GLN
1	B	453	ARG
1	B	454	GLN
1	B	455	TYR
1	B	666	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	671	ARG
1	B	699	LEU
1	B	738	SER
1	B	780	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	163	GLN
1	A	219	GLN
1	A	726	ASN
1	B	163	GLN
1	B	562	ASN
1	B	726	ASN
1	B	760	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	M3L	B	549	1	10,11,12	0.64	0	9,14,16	0.43	0
1	M3L	B	129	1	10,11,12	0.69	0	9,14,16	0.49	0
1	M3L	A	129	1	10,11,12	0.93	1 (10%)	9,14,16	0.46	0
1	M3L	A	549	1	10,11,12	0.82	1 (10%)	9,14,16	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	M3L	B	549	1	-	2/9/10/12	-
1	M3L	B	129	1	-	3/9/10/12	-
1	M3L	A	129	1	-	3/9/10/12	-
1	M3L	A	549	1	-	2/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	129	M3L	CB-CA	2.68	1.57	1.53
1	A	549	M3L	CB-CA	2.17	1.56	1.53

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	549	M3L	CA-CB-CG-CD
1	B	549	M3L	CA-CB-CG-CD
1	B	549	M3L	CE-CD-CG-CB
1	A	549	M3L	CE-CD-CG-CB
1	A	129	M3L	CD-CE-NZ-CM3
1	B	129	M3L	CD-CE-NZ-CM3
1	A	129	M3L	CD-CE-NZ-CM2
1	B	129	M3L	CD-CE-NZ-CM2
1	A	129	M3L	CD-CE-NZ-CM1
1	B	129	M3L	CD-CE-NZ-CM1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	ADP	B	803	6,3	24,29,29	0.84	1 (4%)	29,45,45	0.90	1 (3%)
2	SO4	A	1001	-	4,4,4	0.30	0	6,6,6	0.13	0
6	VO4	B	804	5,3	1,4,4	1.11	0	-		
4	6I6	A	1003	-	21,28,28	0.69	1 (4%)	27,40,40	1.30	2 (7%)
5	ADP	A	1004	6,3	24,29,29	0.74	0	29,45,45	0.78	1 (3%)
6	VO4	A	1005	5,3	1,4,4	0.29	0	-		
4	6I6	B	802	-	21,28,28	0.73	1 (4%)	27,40,40	1.12	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	6I6	A	1003	-	-	0/12/23/23	0/4/4/4
5	ADP	B	803	6,3	-	3/12/32/32	0/3/3/3
5	ADP	A	1004	6,3	-	3/12/32/32	0/3/3/3
4	6I6	B	802	-	-	0/12/23/23	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	802	6I6	C3-N	2.81	1.39	1.35
4	A	1003	6I6	C3-N	2.69	1.39	1.35
5	B	803	ADP	PB-O3B	-2.45	1.45	1.54

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003	6I6	C-C1-C2	5.60	120.34	112.59
4	B	802	6I6	C-C1-C2	4.28	118.50	112.59
4	B	802	6I6	C4-C3-N1	2.64	122.56	119.12
4	A	1003	6I6	C4-C3-N1	2.61	122.52	119.12

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	803	ADP	C5-C6-N6	2.37	123.95	120.35
5	A	1004	ADP	C5-C6-N6	2.26	123.79	120.35

There are no chirality outliers.

All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	1004	ADP	PA-O3A-PB-O2B
5	B	803	ADP	PA-O3A-PB-O2B
5	B	803	ADP	PA-O3A-PB-O1B
5	A	1004	ADP	PA-O3A-PB-O3B
5	B	803	ADP	PA-O3A-PB-O3B
5	A	1004	ADP	PA-O3A-PB-O1B



There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	804	VO4	3	0
4	A	1003	6I6	2	0
6	A	1005	VO4	2	0
4	B	802	6I6	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	715/782 (91%)	0.54	41 (5%) 30 37	33, 64, 102, 142	3 (0%)
1	B	704/782 (90%)	1.14	117 (16%) 5 7	33, 78, 122, 169	2 (0%)
All	All	1419/1564 (90%)	0.84	158 (11%) 12 15	33, 70, 114, 169	5 (0%)

All (158) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	130	TRP	8.0
1	B	729	ALA	7.1
1	A	130	TRP	5.6
1	B	336	PHE	5.0
1	A	663[A]	ARG	5.0
1	B	202	GLY	4.9
1	B	39	VAL	4.7
1	B	580	ILE	4.5
1	A	625	ALA	4.3
1	B	33	LEU	4.2
1	A	736	ILE	4.2
1	B	724	ILE	4.2
1	B	467	ILE	4.1
1	B	557	LEU	4.0
1	B	781	LEU	4.0
1	B	70	THR	3.8
1	B	61	ALA	3.7
1	B	279	ALA	3.7
1	B	718	PHE	3.7
1	B	709	PHE	3.7
1	B	533	ILE	3.6
1	B	764	PHE	3.6
1	B	558	GLY	3.5
1	B	69	VAL	3.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	570	LYS	3.5
1	B	767	ALA	3.5
1	A	563	PHE	3.4
1	B	49	ALA	3.4
1	B	574	GLU	3.4
1	A	378	THR	3.4
1	A	213	LYS	3.4
1	B	297	LEU	3.3
1	A	63	THR	3.3
1	B	341	PHE	3.3
1	B	510	PHE	3.3
1	B	57	GLY	3.3
1	A	569	ILE	3.3
1	A	565	LYS	3.2
1	B	367	LYS	3.2
1	B	294	LYS	3.2
1	B	722	TYR	3.2
1	B	744	LYS	3.1
1	B	569	ILE	3.1
1	B	112	TRP	3.1
1	B	214	GLY	3.0
1	B	68	THR	3.0
1	B	52	LEU	3.0
1	B	35	LYS	3.0
1	B	295	PRO	3.0
1	B	624	TYR	3.0
1	A	201	ILE	3.0
1	A	450	LYS	3.0
1	B	556	HIS	2.9
1	B	289	ILE	2.9
1	B	577	PHE	2.9
1	B	452	PRO	2.9
1	B	278	LYS	2.9
1	B	303	ILE	2.9
1	B	565	LYS	2.9
1	A	577	PHE	2.8
1	B	94	THR	2.8
1	A	408	ASN	2.8
1	A	33	LEU	2.8
1	B	72	LYS	2.8
1	B	536	GLU	2.8
1	B	501	TYR	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	59	VAL	2.7
1	B	354	GLY	2.7
1	B	378	THR	2.7
1	B	298	LEU	2.7
1	A	510	PHE	2.7
1	B	386	TYR	2.7
1	B	406	VAL	2.7
1	B	645	GLN	2.7
1	B	330	MET	2.7
1	B	506	ILE	2.7
1	A	211	THR	2.6
1	A	220	ILE	2.6
1	B	377	GLY	2.6
1	B	567	ARG	2.6
1	B	92[A]	MET	2.6
1	B	34	LYS	2.6
1	B	540	PHE	2.5
1	B	773	LEU	2.5
1	A	56	GLY	2.5
1	B	446	THR	2.5
1	A	624	TYR	2.5
1	B	563	PHE	2.5
1	B	765	PHE	2.5
1	B	290	LEU	2.5
1	A	367	LYS	2.4
1	B	383	LYS	2.4
1	A	767	ALA	2.4
1	A	552	LEU	2.4
1	A	369	ARG	2.4
1	B	56	GLY	2.4
1	B	37	VAL	2.4
1	A	761	THR	2.4
1	A	782	SER	2.4
1	B	578	SER	2.4
1	A	570	LYS	2.4
1	A	214	GLY	2.4
1	B	579	LEU	2.4
1	B	622	ALA	2.4
1	B	410	TYR	2.4
1	B	253	GLY	2.4
1	A	412	THR	2.3
1	B	708	GLY	2.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	38	PHE	2.3
1	B	751	ILE	2.3
1	B	758	PHE	2.3
1	B	762	LYS	2.3
1	B	299	ASP	2.3
1	B	530	ILE	2.3
1	A	536	GLU	2.2
1	B	67	LYS	2.2
1	B	598	LYS	2.2
1	B	539	MET	2.2
1	A	731	PRO	2.2
1	B	326	ALA	2.2
1	B	304	THR	2.2
1	B	531	MET	2.2
1	B	769	LEU	2.2
1	A	544	THR	2.2
1	A	417	VAL	2.2
1	B	431	VAL	2.2
1	B	58	LYS	2.2
1	B	75	GLN	2.2
1	B	740	LYS	2.2
1	B	716	GLY	2.1
1	B	541	PRO	2.1
1	B	350	TYR	2.1
1	B	412	THR	2.1
1	B	455	TYR	2.1
1	A	421	VAL	2.1
1	A	336	PHE	2.1
1	B	62	GLU	2.1
1	B	220	ILE	2.1
1	B	329	LEU	2.1
1	A	279	ALA	2.1
1	B	514	GLY	2.1
1	A	72	LYS	2.1
1	A	557	LEU	2.1
1	B	449	THR	2.1
1	A	480	PHE	2.1
1	B	451	GLN	2.1
1	A	573	PRO	2.1
1	B	133	VAL	2.0
1	B	542	LYS	2.0
1	B	339	LEU	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	375	PRO	2.0
1	B	254	ALA	2.0
1	B	438	TRP	2.0
1	B	757	LYS	2.0
1	A	52	LEU	2.0
1	B	573	PRO	2.0
1	B	617	LEU	2.0
1	B	725	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	M3L	A	549	12/13	0.82	0.20	71,79,96,96	0
1	M3L	B	129	12/13	0.84	0.17	73,77,85,86	0
1	M3L	A	129	12/13	0.92	0.12	57,60,68,69	0
1	M3L	B	549	12/13	0.92	0.17	80,89,107,109	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	1001	5/5	0.95	0.08	102,102,104,105	0
4	6I6	A	1003	25/25	0.96	0.07	47,55,62,64	0
4	6I6	B	802	25/25	0.97	0.07	45,49,57,57	0
5	ADP	B	803	27/27	0.97	0.07	47,62,67,72	0
6	VO4	B	804	5/5	0.97	0.08	50,52,58,65	0
3	MG	B	801	1/1	0.98	0.04	53,53,53,53	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	VO4	A	1005	5/5	0.98	0.07	43,48,53,61	0
5	ADP	A	1004	27/27	0.98	0.06	40,52,56,65	0
3	MG	A	1002	1/1	0.99	0.07	53,53,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around 6I6 B 802:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP B 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.