

Full wwPDB X-ray Structure Validation Report (i)

Sep 9, 2025 – 01:06 pm BST

PDB ID : 9HPS / pdb 00009hps

Title: Human BclxLdeltaLT-VDAC1-N fusion protein complex structure

Authors: Janowski, R.; Niessing, D.

Deposited on : 2024-12-16

Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 2.0rc1 EDS : 3.0

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.45.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	164625	3187 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 1437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called BclxLdeltaLT.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	146	Total C N O S 1206 767 204 231 4	0	4	0
1	В	18	Total C N O 136 88 23 25	0	0	0

• Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0

• Molecule 3 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	74	Total O 74 74	0	0
3	В	6	Total O 6 6	0	0

 $\operatorname{MolProbity}$ failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	35.12Å 98.75Å 103.91Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.64 - 1.95	Depositor
resolution (A)	44.64 - 1.95	EDS
% Data completeness	98.7 (44.64-1.95)	Depositor
(in resolution range)	98.7 (44.64-1.95)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.81 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
P.P.	0.170 , 0.201	Depositor
R, R_{free}	0.183 , 0.205	DCC
R_{free} test set	677 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtriage
Anisotropy	0.084	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38, 46.8	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1437	wwPDB-VP
Average B, all atoms $(Å^2)$	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.11% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

4.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Type	Type Chain Dec		Chain	Chain	Chain Res L	Link	B	ond leng	${ m gths}$	В	ond ang	gles
MIOI		туре	Туре	Chain	LIIIK		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
2	SO4	A	303	-	4,4,4	0.29	0	6,6,6	0.23	0			
2	SO4	A	302	-	4,4,4	0.39	0	6,6,6	0.15	0			
2	SO4	A	301	-	4,4,4	0.41	0	6,6,6	0.14	0			

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	146/207 (70%)	0.28	11 (7%) 22 26	10, 27, 78, 103	5 (3%)
1	В	18/207 (8%)	1.67	6 (33%) 1 1	35, 54, 71, 73	0
All	All	164/414 (39%)	0.43	17 (10%) 13 16	10, 29, 77, 103	5 (3%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	24	TRP	5.1
1	В	587	LEU	4.6
1	A	102	ARG	4.2
1	В	583	TYR	4.2
1	A	104	ALA	4.1
1	A	27	PHE	3.3
1	A	23	SER	3.0
1	A	3	GLN	3.0
1	В	585	PHE	2.8
1	В	580	THR	2.7
1	A	81	THR	2.6
1	A	107	ASP	2.6
1	A	25	SER	2.2
1	A	103	ARG	2.2
1	В	584	GLY	2.2
1	В	582	GLY	2.1
1	A	195	TYR	2.0

5.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	SO4	A	303	5/5	0.69	0.15	55,64,72,73	5
2	SO4	A	302	5/5	0.88	0.12	34,36,43,47	5
2	SO4	A	301	5/5	0.93	0.14	37,40,45,49	5

5.5 Other polymers (i)

There are no such residues in this entry.

