

Full wwPDB X-ray Structure Validation Report (i)

Aug 11, 2025 – 02:04 pm BST

PDB ID : 9IBE / pdb 00009ibe

Title : D-2-hydroxyacid dehydrogenase (D2HDH) from Haloferax mediterranei in

complex with potassium, 2-ketohexanoic acid, NADP+ and chloride

Authors: Baker, P.J.; Barrett, J.R.; Dakhil, A.A.A.B.; Domenech, J.; Bisson, C.; Pra-

manpol, N.; Ferrer, J.; Rice, D.W.

Deposited on : 2025-02-12

Resolution : 1.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 2.0rc1 EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.45.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	164625	1447 (1.28-1.24)
Clashscore	180529	1571 (1.28-1.24)
Ramachandran outliers	177936	1538 (1.28-1.24)
Sidechain outliers	177891	1537 (1.28-1.24)
RSRZ outliers	164620	1447 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	308	97%	•
1	В	308	94%	6%
1	С	308	96%	•
1	D	308	94%	6% •



2 Entry composition (i)


There are 7 unique types of molecules in this entry. The entry contains 11390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called D-2-hydroxyacid dehydrogenase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	Λ	308	Total	С	N	О	S	0	7		
1	A	300	2382	1492	407	475	8	U	1		
1	В	308	Total	С	N	О	S	0	8	0	
1	Ъ	300	2403	1502	416	477	8	0		0	
1	С	308	Total	С	N	О	S	0	8	0	
1		300	2391	1496	408	479	8	0	0	"	
1	D	308	Total	С	N	О	S	0	4	0	
	308	2370	1483	405	474	8	0	4	U		

• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	Λ	1	Total	С	N	О	Р	0	0
2	A	1	48	21	7	17	3	U	
2	D	1	Total	С	N	О	Р	0	0
2	Б	1	48	21	7	17	3	U	0



Continued from previous page...

	Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
	9	C	1	Total	С	N	О	Р	0	0
	2	C	1	48	21	7	17	3	U	
ĺ	2	D	1	Total	С	N	N O P	0		
	$\mathcal{L} \qquad \mathcal{D}$	$D \mid 1 \mid$		21	7	17	3	U	U	

• Molecule 3 is 2-Ketohexanoic acid (CCD ID: 7N5) (formula: C₆H₁₀O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 9 6 3	0	0
3	В	1	Total C O 9 6 3	0	0
3	С	1	Total C O 9 6 3	0	0
3	D	1	Total C O 9 6 3	0	0

• Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	7	Total K 7 7	0	0
4	В	5	Total K 5 5	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	7	Total K 7 7	0	0
4	D	4	Total K 5 5	0	1

• Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total Mg 2 2	0	0
5	В	4	Total Mg 4 4	0	0
5	С	3	Total Mg 3 3	0	0
5	D	3	Total Mg 3 3	0	0

• Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Cl 2 2	0	0
6	В	1	Total Cl 1 1	0	0
6	С	1	Total Cl 1 1	0	0
6	D	1	Total Cl 1 1	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	383	Total O 385 385	0	2
7	В	435	Total O 436 436	0	1
7	С	413	Total O 416 416	0	3
7	D	338	Total O 338 338	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: D-2-hydroxyacid dehydrogenase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	66.63Å 75.34Å 78.23Å	Donositon
a, b, c, α , β , γ	108.84° 108.08° 95.53°	Depositor
Resolution (Å)	52.28 - 1.26	Depositor
Resolution (A)	52.28 - 1.26	EDS
% Data completeness	92.2 (52.28-1.26)	Depositor
(in resolution range)	89.6 (52.28-1.26)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.47 (at 1.26Å)	Xtriage
Refinement program	REFMAC 5.8.0430 (refmacat 0.4.88)	Depositor
D D.	0.152 , 0.188	Depositor
R, R_{free}	0.160 , 0.192	DCC
R_{free} test set	16700 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	10.6	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 34.8	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	11390	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 66.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1838e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 7N5, NAP, K, CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.73	0/2457	1.10	1/3351~(0.0%)
1	В	0.74	0/2474	1.09	$3/3372 \ (0.1\%)$
1	С	0.72	0/2467	1.08	$2/3365 \ (0.1\%)$
1	D	0.73	0/2432	1.09	6/3317~(0.2%)
All	All	0.73	0/9830	1.09	12/13405 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	В	0	2
1	С	0	1
1	D	0	4
All	All	0	9

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$\operatorname{Ideal}(^{o})$
1	A	201	ASP	CA-CB-CG	7.67	120.27	112.60
1	D	201	ASP	CA-CB-CG	7.11	119.71	112.60
1	D	165	ARG	NE-CZ-NH1	-6.33	115.17	121.50
1	В	165	ARG	NE-CZ-NH1	-6.28	115.22	121.50
1	D	260	ASP	CA-CB-CG	5.99	118.59	112.60
1	С	41	PHE	CA-CB-CG	-5.87	107.93	113.80
1	D	202	GLU	CB-CG-CD	5.69	122.27	112.60
1	В	165	ARG	NE-CZ-NH2	5.42	124.08	119.20



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	В	296	LYS	CB-CG-CD	5.41	123.74	111.30
1	D	165	ARG	NE-CZ-NH2	5.36	124.02	119.20
1	С	166	ARG	CG-CD-NE	-5.30	100.35	112.00
1	D	259	GLU	CB-CA-C	5.11	120.08	110.63

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	20	ARG	Sidechain
1	A	306	ARG	Sidechain
1	В	137[A]	ARG	Sidechain
1	В	137[B]	ARG	Sidechain
1	С	182	ARG	Sidechain
1	D	126	ARG	Sidechain
1	D	165	ARG	Sidechain
1	D	166	ARG	Sidechain
1	D	306	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2382	0	2268	3	0
1	В	2403	0	2297	8	1
1	С	2391	0	2266	4	0
1	D	2370	0	2256	8	0
2	A	48	0	25	0	0
2	В	48	0	25	0	0
2	С	48	0	25	0	0
2	D	48	0	25	0	0
3	A	9	0	0	0	0
3	В	9	0	0	0	0
3	С	9	0	0	0	0
3	D	9	0	0	0	0
4	A	7	0	0	0	0
4	В	5	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	С	7	0	0	0	0
4	D	5	0	0	0	0
5	A	2	0	0	0	0
5	В	4	0	0	0	0
5	С	3	0	0	0	0
5	D	3	0	0	0	0
6	A	2	0	0	0	0
6	В	1	0	0	0	0
6	С	1	0	0	0	0
6	D	1	0	0	0	0
7	A	385	0	0	0	0
7	В	436	0	0	1	0
7	С	416	0	0	0	1
7	D	338	0	0	2	0
All	All	11390	0	9187	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance } (\text{\AA}) \end{array}$	Clash overlap (Å)
1:B:291[A]:ARG:HD3	7:B:839[A]:HOH:O	1.91	0.70
1:A:137:ARG:NH2	1:A:160:GLU:OE1	2.41	0.53
1:D:137:ARG:NH2	1:D:160:GLU:OE1	2.42	0.52
1:D:126:ARG:HD3	7:D:560:HOH:O	2.10	0.50
1:D:126:ARG:CD	7:D:560:HOH:O	2.60	0.50
1:B:1:MET:HE1	1:B:83:TYR:CD2	2.49	0.48
1:B:255:GLU:HA	1:B:256:PRO:C	2.38	0.48
1:D:255:GLU:HA	1:D:256:PRO:C	2.38	0.48
1:D:128[B]:GLU:H	1:D:128[B]:GLU:CD	2.24	0.46
1:A:247:ALA:O	1:A:269:VAL:HA	2.17	0.45
1:B:1:MET:HE2	1:B:297:ILE:HG21	2.00	0.44
1:A:255:GLU:HA	1:A:256:PRO:C	2.43	0.42
1:B:86[B]:ASN:OD1	1:B:88:THR:HG22	2.19	0.42
1:C:247:ALA:O	1:C:269:VAL:HA	2.19	0.42
1:D:247:ALA:O	1:D:269:VAL:HA	2.19	0.42
1:B:247:ALA:O	1:B:269:VAL:HA	2.20	0.41
1:C:137:ARG:NH2	1:C:160:GLU:OE1	2.53	0.41
1:D:221:LEU:O	1:D:247:ALA:HA	2.20	0.41
1:C:255:GLU:HA	1:C:256:PRO:C	2.45	0.41



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:227:GLY:HA3	1:C:252:PHE:C	2.46	0.41
1:B:86[A]:ASN:OD1	1:B:305:ASN:HB3	2.21	0.41
1:D:227:GLY:HA3	1:D:252:PHE:C	2.47	0.40
1:B:227:GLY:HA3	1:B:252:PHE:C	2.46	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:201[A]:ASP:OD2	7:C:624:HOH:O[1_554]	2.11	0.09

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	313/308 (102%)	309 (99%)	4 (1%)	0	100	100
1	В	314/308 (102%)	308 (98%)	6 (2%)	0	100	100
1	С	314/308 (102%)	308 (98%)	6 (2%)	0	100	100
1	D	310/308 (101%)	304 (98%)	6 (2%)	0	100	100
All	All	1251/1232 (102%)	1229 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	A	252/245 (103%)	252 (100%)	0	100	100
1	В	253/245 (103%)	252 (100%)	1 (0%)	89	76
1	С	253/245 (103%)	252 (100%)	1 (0%)	89	76
1	D	249/245 (102%)	247 (99%)	2 (1%)	79	52
All	All	1007/980 (103%)	1003 (100%)	4 (0%)	89	76

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	279	THR
1	С	279	THR
1	D	295	GLU
1	D	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	2	HIS
1	D	19	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 49 ligands modelled in this entry, 41 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	NAP	В	401	4	45,52,52	0.93	1 (2%)	56,80,80	1.00	3 (5%)
3	7N5	В	402	-	8,8,8	2.37	3 (37%)	9,9,9	1.40	2 (22%)
3	7N5	D	402	-	8,8,8	1.74	1 (12%)	9,9,9	1.11	1 (11%)
2	NAP	A	401	4	45,52,52	0.67	0	56,80,80	0.90	3 (5%)
3	7N5	A	402	-	8,8,8	1.42	2 (25%)	9,9,9	1.92	2 (22%)
3	7N5	С	402	-	8,8,8	2.08	3 (37%)	9,9,9	1.56	1 (11%)
2	NAP	С	401	4	45,52,52	0.81	1 (2%)	56,80,80	0.95	2 (3%)
2	NAP	D	401	4	45,52,52	0.87	2 (4%)	56,80,80	0.87	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	В	401	4	-	4/31/67/67	0/5/5/5
3	7N5	В	402	-	-	3/8/8/8	-
3	7N5	D	402	-	-	3/8/8/8	-
2	NAP	A	401	4	-	3/31/67/67	0/5/5/5
3	7N5	A	402	-	-	4/8/8/8	-
3	7N5	С	402	-	-	2/8/8/8	-
2	NAP	С	401	4	-	4/31/67/67	0/5/5/5
2	NAP	D	401	4	-	3/31/67/67	0/5/5/5

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(\mathring{A})$	Ideal(A)
3	В	402	7N5	C2-C1	-5.63	1.46	1.53
3	С	402	7N5	C2-C1	-4.35	1.47	1.53
2	В	401	NAP	P2B-O2B	4.26	1.67	1.59
3	D	402	7N5	C2-C1	-3.73	1.48	1.53
2	D	401	NAP	P2B-O2B	3.32	1.65	1.59
2	С	401	NAP	C2N-N1N	3.19	1.38	1.35



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	A	402	7N5	C2-C1	-2.84	1.49	1.53
3	С	402	7N5	O1B-C1	-2.46	1.23	1.30
2	D	401	NAP	C2N-N1N	2.31	1.37	1.35
3	С	402	7N5	O1A-C1	2.23	1.28	1.22
3	В	402	7N5	O1A-C1	2.22	1.28	1.22
3	В	402	7N5	C4-C3	2.13	1.60	1.52
3	A	402	7N5	O1A-C1	2.08	1.28	1.22

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	A	402	7N5	C3-C2-C1	3.99	123.38	115.97
3	A	402	7N5	C4-C3-C2	3.44	120.71	113.63
3	С	402	7N5	O2-C2-C3	3.26	128.42	121.20
3	В	402	7N5	C4-C3-C2	2.66	119.11	113.63
2	С	401	NAP	O2B-P2B-O1X	-2.56	99.53	109.39
2	В	401	NAP	C3B-C2B-C1B	-2.55	98.09	102.89
2	В	401	NAP	O3X-P2B-O2X	2.54	117.34	107.64
2	A	401	NAP	C5A-C6A-N6A	2.32	123.89	120.35
2	D	401	NAP	C3B-C2B-C1B	-2.29	98.58	102.89
3	В	402	7N5	O2-C2-C3	2.25	126.19	121.20
2	С	401	NAP	C3B-C2B-C1B	-2.23	98.70	102.89
2	A	401	NAP	O7N-C7N-C3N	-2.19	117.01	119.63
2	В	401	NAP	C1B-N9A-C4A	-2.15	122.86	126.64
2	A	401	NAP	C3B-C2B-C1B	-2.08	98.98	102.89
2	D	401	NAP	O3X-P2B-O1X	2.06	118.75	110.68
2	D	401	NAP	C4A-C5A-N7A	2.03	111.52	109.40
3	D	402	7N5	O2-C2-C3	2.02	125.68	121.20

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	401	NAP	C3B-C2B-O2B-P2B
2	В	401	NAP	O4D-C1D-N1N-C6N
2	С	401	NAP	O4D-C1D-N1N-C6N
2	D	401	NAP	C3B-C2B-O2B-P2B
3	A	402	7N5	O2-C2-C3-C4
3	A	402	7N5	O1B-C1-C2-C3
3	В	402	7N5	O1B-C1-C2-C3
3	С	402	7N5	O1B-C1-C2-C3
3	D	402	7N5	O1B-C1-C2-C3



Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	С	401	NAP	C3B-C2B-O2B-P2B
2	A	401	NAP	C3B-C2B-O2B-P2B
2	A	401	NAP	C1B-C2B-O2B-P2B
3	A	402	7N5	C3-C4-C5-C6
3	A	402	7N5	C1-C2-C3-C4
3	D	402	7N5	O2-C2-C3-C4
3	В	402	7N5	O2-C2-C3-C4
3	С	402	7N5	O2-C2-C3-C4
2	С	401	NAP	C1B-C2B-O2B-P2B
2	D	401	NAP	C1B-C2B-O2B-P2B
2	В	401	NAP	C1B-C2B-O2B-P2B
3	В	402	7N5	C1-C2-C3-C4
3	D	402	7N5	C1-C2-C3-C4
2	A	401	NAP	O4B-C4B-C5B-O5B
2	В	401	NAP	O4B-C4B-C5B-O5B
2	С	401	NAP	O4B-C4B-C5B-O5B
2	D	401	NAP	O4B-C4B-C5B-O5B

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





















5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	308/308 (100%)	-0.08	8 (2%) 57 57	7, 13, 37, 77	7 (2%)
1	В	308/308 (100%)	-0.21	5 (1%) 70 69	7, 13, 28, 90	8 (2%)
1	С	308/308 (100%)	-0.19	7 (2%) 61 61	7, 13, 37, 69	8 (2%)
1	D	308/308 (100%)	0.09	14 (4%) 39 38	8, 15, 49, 73	4 (1%)
All	All	1232/1232 (100%)	-0.10	34 (2%) 55 55	7, 13, 40, 90	27 (2%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	1	MET	5.0
1	D	1	MET	3.5
1	В	2	HIS	3.5
1	В	27	ASP	3.4
1	С	1	MET	3.3
1	С	298	ALA	3.2
1	D	43	PRO	3.1
1	С	297	ILE	3.0
1	A	1	MET	3.0
1	В	76	VAL	3.0
1	A	34	PHE	2.9
1	С	76	VAL	2.7
1	D	28	LEU	2.7
1	A	297	ILE	2.5
1	A	76	VAL	2.5
1	A	2	HIS	2.4
1	В	26	SER	2.4
1	С	299	THR	2.4
1	D	294	ILE	2.4
1	D	297	ILE	2.4
1	D	32	VAL	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	75	GLY	2.3
1	A	308	VAL	2.3
1	D	76	VAL	2.3
1	D	27	ASP	2.3
1	D	2	HIS	2.2
1	С	302	GLU	2.1
1	D	74	VAL	2.1
1	D	30	VAL	2.1
1	С	300	GLY	2.1
1	D	300	GLY	2.1
1	D	31	PRO	2.0
1	A	74	VAL	2.0
1	D	298	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathrm{A}}^2)$	Q<0.9
3	7N5	С	402	9/9	0.96	0.09	14,16,29,35	0
4	K	A	408	1/1	0.96	0.08	29,29,29,29	1
4	K	С	408	1/1	0.96	0.10	27,27,27,27	1
6	CL	A	413	1/1	0.96	0.12	35,35,35,35	0
3	7N5	В	402	9/9	0.97	0.07	13,14,23,27	0
3	7N5	A	402	9/9	0.97	0.08	15,16,32,34	0
5	MG	В	411	1/1	0.97	0.18	32,32,32,32	0
3	7N5	D	402	9/9	0.97	0.07	15,16,26,27	0
6	CL	D	410	1/1	0.97	0.10	33,33,33,33	0
4	K	D	406[B]	1/1	0.98	0.05	39,39,39,39	1



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$oxed{ \mathbf{B\text{-}factors}(\mathbf{\mathring{A}}^2) }$	Q<0.9
4	K	В	407	1/1	0.98	0.17	33,33,33,33	0
5	MG	D	409	1/1	0.98	0.10	34,34,34,34	0
2	NAP	С	401	48/48	0.98	0.04	9,12,19,24	0
4	K	D	406[A]	1/1	0.98	0.05	33,33,33,33	1
2	NAP	В	401	48/48	0.99	0.03	7,10,14,16	0
4	K	D	404	1/1	0.99	0.12	26,26,26,26	0
4	K	A	406	1/1	0.99	0.05	36,36,36,36	0
2	NAP	A	401	48/48	0.99	0.04	8,11,16,22	0
5	MG	A	411	1/1	0.99	0.07	23,23,23,23	0
5	MG	В	409	1/1	0.99	0.03	16,16,16,16	0
5	MG	В	410	1/1	0.99	0.08	19,19,19,19	0
4	K	В	404	1/1	0.99	0.11	25,25,25,25	0
5	MG	С	412	1/1	0.99	0.14	25,25,25,25	0
2	NAP	D	401	48/48	0.99	0.03	9,11,14,18	0
6	CL	A	412	1/1	0.99	0.13	24,24,24,24	0
4	K	С	404	1/1	0.99	0.09	25,25,25,25	0
6	CL	В	412	1/1	0.99	0.04	17,17,17,17	0
4	K	С	406	1/1	0.99	0.03	31,31,31,31	0
4	K	В	403	1/1	1.00	0.02	13,13,13,13	0
4	K	D	405	1/1	1.00	0.06	18,18,18,18	0
4	K	A	405	1/1	1.00	0.07	19,19,19,19	0
4	K	В	405	1/1	1.00	0.07	15,15,15,15	0
5	MG	A	410	1/1	1.00	0.03	10,10,10,10	0
4	K	В	406	1/1	1.00	0.06	25,25,25,25	0
5	MG	В	408	1/1	1.00	0.01	9,9,9,9	0
4	K	A	403	1/1	1.00	0.02	13,13,13,13	0
4	K	С	403	1/1	1.00	0.02	13,13,13,13	0
4	K	A	407	1/1	1.00	0.07	15,15,15,15	0
5	MG	С	410	1/1	1.00	0.01	9,9,9,9	0
5	MG	С	411	1/1	1.00	0.07	17,17,17,17	0
4	K	С	405	1/1	1.00	0.09	17,17,17,17	0
5	MG	D	407	1/1	1.00	0.01	11,11,11,11	0
5	MG	D	408	1/1	1.00	0.04	14,14,14,14	0
4	K	A	404	1/1	1.00	0.09	23,23,23,23	0
4	K	С	407	1/1	1.00	0.05	19,19,19,19	0
4	K	A	409	1/1	1.00	0.06	17,17,17,17	0
4	K	С	409	1/1	1.00	0.08	21,21,21,21	0
6	CL	С	413	1/1	1.00	0.05	17,17,17,17	0
4	K	D	403	1/1	1.00	0.01	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different




orientation to approximate a three-dimensional view.


Electron density around 7N5 C 402: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around K A 408:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around K C 408:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around CL A 413:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around 7N5 B 402:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 7N5 A 402:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















Electron density around CL D 410:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around K D 406 (B):

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around K B 407:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















Electron density around K D 406 (A):


 $2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)
















Electron density around K D 404:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around K A 406: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around K B 404:
















Electron density around CL A 412:












Electron density around K C 404:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around CL B 412:












Electron density around K C 406:













Electron density around K B 403:









Electron density around K D 405:












Electron density around K A 405:












Electron density around K B 405:












Electron density around K B 406:












Electron density around K A 403:












Electron density around K C 403:












Electron density around K A 407:












Electron density around K C 405:












Electron density around K A 404:












Electron density around K C 407:












Electron density around K A 409:












Electron density around K C 409:













Electron density around CL C 413:











6.5 Other polymers (i)

There are no such residues in this entry.

