

Full wwPDB EM Validation Report (i)

Jul 15, 2025 – 11:29 PM JST

PDB ID : 9ITB / pdb 00009itb

EMDB ID : EMD-60857

Title: LPA-bound LPAR6 in complex with miniGq

Authors: He, Y.; Duan, Y.

Deposited on : 2024-07-19

Resolution : 2.89 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : FAILED

Mogul : 1.8.5 (274361), CSD as 541 be (2020)

MolProbity : 4-5-2 with Phenix2.0rc1

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ: FAILED

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain						
1	A	362	61%	37%					
2	В	345	93%						
3	N	161	75%	• 20%					
4	R	294	86%	5% 8%					
5	G	71	79%	21%					



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 7980 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called engineered miniGaq.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	228	Total 1818		N 319	O 341	S 7	0	0

• Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	В	334	Total 2514	C 1559	N 439	O 495	S 21	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-4	MET	-	initiating methionine	UNP P62873
В	-3	GLY	-	expression tag	UNP P62873
В	-2	SER	-	expression tag	UNP P62873
В	-1	LEU	-	expression tag	UNP P62873
В	0	LEU	-	expression tag	UNP P62873
В	1	GLN	-	expression tag	UNP P62873

• Molecule 3 is a protein called Nb35.

Mo	Chain	Residues		Atoms				AltConf	Trace
3	N	128	Total 963	C 600	N 167	O 190	S 6	0	0

• Molecule 4 is a protein called Lysophosphatidic acid receptor 6.

Mol	Chain	Residues		Atoms				AltConf	Trace
4	R	271	Total 2147	C 1441	N 327	O 355	S 24	0	0

There are 2 discrepancies between the modelled and reference sequences:




Chain	Residue	Modelled	Actual	Comment	Reference
R	43	ALA	ILE	conflict	UNP P43657
R	166	ARG	GLU	conflict	UNP P43657

• Molecule 5 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms			AltConf	Trace		
5	G	56	Total	С	N	О	S	0	0
	G	50	407	258	68	78	3		


• Molecule 6 is (2R)-2-hydroxy-3-(phosphonooxy) propyl (9E)-octadec-9-enoate (CCD ID: NKP) (formula: $C_{21}H_{41}O_7P$).



Mol	Chain	Residues	${f Atoms}$				AltConf
6	R	1	Total 29	C 21	O 7	P 1	0


 \bullet Molecule 7 is CHOLESTEROL (CCD ID: CLR) (formula: $\mathrm{C_{27}H_{46}O}).$





Mol	Chain	Residues	Atoms	AltConf
7	R.	1	Total C O	0
'	16	1	28 27 1	0
7	R.	1	Total C O	0
'	16	1	28 27 1	0
7	B	1	Total C O	0
'	10	1	28 27 1	

 $\bullet\,$ Molecule 8 is PALMITIC ACID (CCD ID: PLM) (formula: $\mathrm{C}_{16}\mathrm{H}_{32}\mathrm{O}_2).$




	Mol	Chain	Residues	Atoms			AltConf
Ī	8	R	1	Total	С	О	0
	O	10	1	18	16	2	



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: engineered miniGaq



• Molecule 5: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2



Chain G: 79% 21%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	211505	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TECNAI F30	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	60	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NKP, PLM, CLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bo	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.20	0/1850	0.41	0/2500
2	В	0.22	0/2560	0.50	0/3479
3	N	0.19	0/983	0.46	0/1333
4	R	0.20	0/2203	0.46	1/3000~(0.0%)
5	G	0.14	0/413	0.28	0/561
All	All	0.21	0/8009	0.46	1/10873~(0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
4	R	82	TRP	N-CA-C	6.44	120.10	109.79

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1818	0	1753	7	0
2	В	2514	0	2383	8	0
3	N	963	0	927	4	0
4	R	2147	0	2181	14	0
5	G	407	0	402	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	R	29	0	39	1	0
7	R	84	0	138	4	0
8	R	18	0	31	1	0
All	All	7980	0	7854	33	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
7:R:402:CLR:H211	7:R:403:CLR:H14	1.73	0.70
4:R:179:THR:O	4:R:183:ARG:NH1	2.27	0.67
2:B:115:GLY:HA3	2:B:146:LEU:HD23	1.77	0.67
4:R:116:LEU:HD21	7:R:402:CLR:H112	1.77	0.66
4:R:82:TRP:CD1	4:R:82:TRP:O	2.52	0.63
4:R:50:ASN:O	4:R:53:THR:OG1	2.15	0.61
1:A:391:TYR:OH	4:R:128:ARG:NH2	2.34	0.61
4:R:288:PRO:HA	4:R:292:TYR:HD2	1.67	0.59
3:N:52:SER:O	3:N:72:ARG:NH1	2.35	0.59
4:R:46:LEU:HD12	4:R:48:VAL:H	1.69	0.57
4:R:82:TRP:O	4:R:82:TRP:CG	2.56	0.56
3:N:94:TYR:O	3:N:121:GLY:HA2	2.05	0.56
1:A:295:ASP:N	1:A:295:ASP:OD1	2.40	0.55
4:R:288:PRO:HA	4:R:292:TYR:CD2	2.41	0.55
4:R:82:TRP:HZ2	4:R:88:LEU:HB3	1.70	0.55
4:R:136:VAL:HG21	7:R:403:CLR:H191	1.89	0.54
1:A:280:ARG:HA	1:A:283:ARG:HH21	1.74	0.53
1:A:393:LEU:HD22	4:R:227:VAL:HG23	1.91	0.52
3:N:33:LYS:HG2	3:N:52:SER:HA	1.92	0.51
1:A:27:GLU:OE2	2:B:89:LYS:NZ	2.31	0.50
2:B:81:ILE:HB	2:B:91:HIS:HB2	1.94	0.50
1:A:280:ARG:HA	1:A:283:ARG:HE	1.76	0.49
2:B:166:CYS:HB2	2:B:180:PHE:HB2	1.93	0.49
4:R:86:ASP:OD1	4:R:86:ASP:N	2.44	0.48
2:B:309:ALA:O	2:B:339:TRP:HH2	1.96	0.48
2:B:325:MET:O	2:B:340:ASN:ND2	2.46	0.48
8:R:405:PLM:HD1	8:R:405:PLM:HA2	1.70	0.47
3:N:47:TRP:HZ2	3:N:50:ASP:HB2	1.82	0.44
2:B:291:ASP:O	2:B:292:PHE:CD1	2.70	0.44
4:R:41:ILE:HG23	4:R:45:VAL:HG22	1.99	0.44
		Continue	ed on nert nage



Continued from previous page...

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} (\mathring{\rm A})$	overlap (Å)
7:R:402:CLR:H241	7:R:403:CLR:H161	1.98	0.44
6:R:401:NKP:HARA	6:R:401:NKP:HAU	1.64	0.43
1:A:228:ARG:NH1	2:B:186:ASP:OD1	2.53	0.42

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$222/362\ (61\%)$	217 (98%)	5 (2%)	0	100	100
2	В	330/345~(96%)	316 (96%)	14 (4%)	0	100	100
3	N	$126/161\ (78\%)$	121 (96%)	5 (4%)	0	100	100
4	R	$265/294\ (90\%)$	253 (96%)	12 (4%)	0	100	100
5	G	54/71~(76%)	52 (96%)	2 (4%)	0	100	100
All	All	997/1233 (81%)	959 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
1	A	192/317 (61%)	192 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
2	В	267/287 (93%)	267 (100%)	0	100	100	
3	N	104/128 (81%)	104 (100%)	0	100	100	
4	R	240/272~(88%)	240 (100%)	0	100	100	
5	G	40/58~(69%)	40 (100%)	0	100	100	
All	All	843/1062 (79%)	843 (100%)	0	100	100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	В	220	GLN
3	N	5	GLN
3	N	74	ASN
3	N	84	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res		Link	Bo	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	CLR	R	403	-	31,31,31	0.84	1 (3%)	48,48,48	1.29	4 (8%)
7	CLR	R	402	-	31,31,31	0.70	0	48,48,48	1.46	8 (16%)
6	NKP	R	401	-	28,28,28	1.51	2 (7%)	31,32,32	1.25	3 (9%)
8	PLM	R	405	-	17,17,17	0.55	0	17,17,17	1.16	1 (5%)
7	CLR	R	404	-	31,31,31	0.58	0	48,48,48	1.13	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	CLR	R	403	-	-	2/10/68/68	0/4/4/4
7	CLR	R	402	-	-	2/10/68/68	0/4/4/4
6	NKP	R	401	-	-	7/28/28/28	-
8	PLM	R	405	-	-	6/15/15/15	-
7	CLR	R	404	-	-	6/10/68/68	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\mathring{\mathrm{A}})$	Ideal(A)
6	R	401	NKP	PAC-OAF	6.84	1.82	1.60
6	R	401	NKP	OAF-CAG	-2.62	1.34	1.44
7	R	403	CLR	C10-C9	-2.35	1.52	1.56

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
7	R	402	CLR	C14-C8-C9	-4.54	103.01	109.09
7	R	403	CLR	C10-C9-C8	-3.70	107.18	112.73
6	R	401	NKP	OAF-PAC-OAD	-3.53	96.59	106.47
7	R	403	CLR	C11-C9-C10	-2.99	109.14	113.08
6	R	401	NKP	OAB-PAC-OAA	2.83	118.44	107.64
7	R	402	CLR	C2-C3-C4	-2.76	106.52	110.31
7	R	402	CLR	C13-C17-C20	-2.70	115.26	119.49
7	R	404	CLR	C13-C17-C20	-2.57	115.46	119.49
7	R	402	CLR	C7-C8-C9	2.53	112.77	109.71



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
7	R	402	CLR	C13-C14-C8	-2.48	110.70	114.38
7	R	403	CLR	C19-C10-C9	-2.37	108.85	111.68
7	R	403	CLR	C13-C14-C8	-2.35	110.90	114.38
7	R	402	CLR	C8-C7-C6	-2.29	109.45	112.73
7	R	402	CLR	C16-C15-C14	-2.25	100.68	105.13
7	R	404	CLR	C11-C12-C13	-2.19	109.03	112.78
6	R	401	NKP	OAB-PAC-OAF	-2.17	100.97	106.73
7	R	404	CLR	C10-C9-C8	-2.12	109.56	112.73
8	R	405	PLM	C3-C2-C1	-2.08	109.22	114.47
7	R	402	CLR	C11-C9-C8	-2.00	108.87	111.75

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	R	401	NKP	OAF-CAG-CAH-CAI
6	R	401	NKP	OAF-CAG-CAH-OBC
7	R	402	CLR	C17-C20-C22-C23
7	R	402	CLR	C21-C20-C22-C23
7	R	404	CLR	C21-C20-C22-C23
7	R	404	CLR	C22-C23-C24-C25
6	R	401	NKP	OBC-CAH-CAI-OAJ
8	R	405	PLM	C4-C5-C6-C7
8	R	405	PLM	C2-C3-C4-C5
7	R	403	CLR	C21-C20-C22-C23
7	R	403	CLR	C17-C20-C22-C23
6	R	401	NKP	CAH-CAG-OAF-PAC
6	R	401	NKP	CAT-CAU-CAV-CAW
6	R	401	NKP	CAG-CAH-CAI-OAJ
8	R	405	PLM	C3-C4-C5-C6
7	R	404	CLR	C13-C17-C20-C21
6	R	401	NKP	CAV-CAW-CAX-CAY
8	R	405	PLM	O2-C1-C2-C3
7	R	404	CLR	C23-C24-C25-C27
8	R	405	PLM	O1-C1-C2-C3
8	R	405	PLM	CA-CB-CC-CD
7	R	404	CLR	C23-C24-C25-C26
7	R	404	CLR	C13-C17-C20-C22


There are no ring outliers.

4 monomers are involved in 6 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	R	403	CLR	3	0
7	R	402	CLR	3	0
6	R	401	NKP	1	0
8	R	405	PLM	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

