



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 12, 2025 – 02:24 PM JST

PDB ID : 9KGJ / pdb_00009kgj
Title : Discovery of an orally bioavailable reversible covalent SARS-CoV-2 Mpro inhibitor with pan-coronavirus activity
Authors : Baburajendran, N.
Deposited on : 2024-11-08
Resolution : 1.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3869 (1.40-1.36)
Clashscore	180529	4183 (1.40-1.36)
Ramachandran outliers	177936	4116 (1.40-1.36)
Sidechain outliers	177891	4115 (1.40-1.36)
RSRZ outliers	164620	3867 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	28% 88% 7% .
1	B	311	22% 89% 7% .
2	C	4	25% 50% 25%
2	D	4	25% 75%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase nsp5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	0	0
			2308	1460	393	433	22			
1	B	299	Total	C	N	O	S	0	0	0
			2298	1459	384	433	22			


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P0DTD1
A	-3	PRO	-	expression tag	UNP P0DTD1
A	-2	LEU	-	expression tag	UNP P0DTD1
A	-1	GLY	-	expression tag	UNP P0DTD1
A	0	SER	-	expression tag	UNP P0DTD1
B	-4	GLY	-	expression tag	UNP P0DTD1
B	-3	PRO	-	expression tag	UNP P0DTD1
B	-2	LEU	-	expression tag	UNP P0DTD1
B	-1	GLY	-	expression tag	UNP P0DTD1
B	0	SER	-	expression tag	UNP P0DTD1

- Molecule 2 is a protein called (S)-3-((S)-3-cyclobutyl-2-((R)-2-cyclopentyl-N-methyl-2-pival amidoacetamido)propanamido)-N-cyclopropyl-2-oxo-4-((S)-2-oxopiperidin-3-yl)butanamide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	4	Total	C	N	O	0	0	0
			37	28	4	5			
2	D	4	Total	C	N	O	0	0	0
			37	28	4	5			

- Molecule 3 is cyclopropylcarbamic acid (CCD ID: A1L7M) (formula: C₄H₇NO₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	O	0	0
			6	4	1	1		
3	D	1	Total	C	N	O	0	0
			6	4	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	190	Total	O	0	0
			190	190		
4	B	209	Total	O	0	0
			209	209		
4	C	1	Total	O	0	0
			1	1		
4	D	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3C-like proteinase nsp45




- Molecule 1: 3C-like proteinase nsp45



- Molecule 2: (S)-3-((S)-3-cyclobutyl-2-((R)-2-cyclopentyl-N-methyl-2-pivalamidoacetamido)prop anamido)-N-cyclopropyl-2-oxo-4-((S)-2-oxopiperidin-3-yl)butanamide



- Molecule 2: (S)-3-((S)-3-cyclobutyl-2-((R)-2-cyclopentyl-N-methyl-2-pivalamidoacetamido)prop anamido)-N-cyclopropyl-2-oxo-4-((S)-2-oxopiperidin-3-yl)butanamide



X1	A1E8K2
	A1L7L3
	A1L7K4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.97Å 99.70Å 59.83Å 90.00° 108.76° 90.00°	Depositor
Resolution (Å)	46.96 – 1.37 46.96 – 1.37	Depositor EDS
% Data completeness (in resolution range)	98.5 (46.96-1.37) 98.5 (46.96-1.37)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	23.57 (at 1.37Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.215 , 0.248 0.216 , 0.248	Depositor DCC
R_{free} test set	1999 reflections (1.55%)	wwPDB-VP
Wilson B-factor (Å ²)	15.0	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5094	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PIV, A1EAK, A1L7M, A1L7K, A1L7L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/2359	0.72	2/3207 (0.1%)
1	B	0.50	0/2350	0.67	0/3198
All	All	0.50	0/4709	0.70	2/6405 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	222	ARG	CB-CG-CD	-8.14	92.58	111.30
1	A	222	ARG	CD-NE-CZ	-6.78	114.90	124.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	222	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2308	0	2257	16	0
1	B	2298	0	2229	18	0
2	C	37	0	9	1	0
2	D	37	0	9	0	0
3	C	6	0	0	0	0
3	D	6	0	0	0	0
4	A	190	0	0	1	0
4	B	209	0	0	0	0
4	C	1	0	0	0	0
4	D	2	0	0	0	0
All	All	5094	0	4504	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:MET:HG2	2:C:3:A1L7L:CD2	2.14	0.77
1:A:110:GLN:HG3	4:A:513:HOH:O	2.01	0.61
1:A:4:ARG:HH12	1:B:5:LYS:NZ	2.00	0.60
1:A:106:ILE:HG12	1:A:110:GLN:HB2	1.85	0.58
1:B:188:ARG:CZ	1:B:190:THR:HG21	2.34	0.57
1:A:4:ARG:HH12	1:B:5:LYS:HZ2	1.55	0.55
1:B:126:TYR:HE2	1:B:128:CYS:SG	2.33	0.52
1:B:288:GLU:HG2	1:B:291:PHE:HD2	1.74	0.51
1:B:56:ASP:HB3	1:B:60:ARG:NH2	2.26	0.50
1:B:288:GLU:HG2	1:B:291:PHE:CD2	2.47	0.50
1:A:6:MET:HE1	1:B:139:SER:HB3	1.95	0.48
1:A:126:TYR:HD1	1:B:6:MET:HG3	1.78	0.48
1:A:131:ARG:HG2	1:A:135:THR:O	2.15	0.47
1:A:6:MET:HE1	1:B:139:SER:CB	2.44	0.47
1:A:76:ARG:HA	1:A:76:ARG:HD3	1.53	0.46
1:A:31:TRP:CD2	1:A:95:ASN:HB2	2.51	0.46
1:A:69:GLN:NE2	1:A:74:GLN:OE1	2.49	0.46
1:B:271:LEU:HD22	1:B:276:MET:HG2	1.97	0.45
1:B:86:VAL:HG23	1:B:179:GLY:HA2	2.00	0.44
1:A:52:PRO:HD3	1:A:188:ARG:HG3	2.00	0.43
1:A:252:PRO:O	1:A:256:GLN:HG3	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:LEU:HD23	1:B:189:GLN:O	2.19	0.43
1:B:126:TYR:CE2	1:B:128:CYS:SG	3.10	0.43
1:B:167:LEU:HA	1:B:167:LEU:HD23	1.82	0.42
1:B:56:ASP:HB3	1:B:60:ARG:HH22	1.85	0.42
1:A:54:TYR:O	1:A:55:GLU:C	2.64	0.41
1:B:165:MET:HG2	1:B:173:ALA:HB3	2.02	0.41
1:A:233:VAL:HG11	1:A:269:LYS:HG3	2.02	0.40
1:B:31:TRP:CE2	1:B:95:ASN:HB2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/311 (95%)	287 (97%)	8 (3%)	1 (0%)	37	16
1	B	297/311 (96%)	291 (98%)	6 (2%)	0	100	100
All	All	593/622 (95%)	578 (98%)	14 (2%)	1 (0%)	44	20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	275	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/266 (97%)	257 (100%)	0	100	100
1	B	254/266 (96%)	254 (100%)	0	100	100
All	All	511/532 (96%)	511 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	ASN
1	A	69	GLN
1	A	119	ASN
1	B	19	GLN
1	B	214	ASN
1	B	273	GLN
1	B	274	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1L7K	C	4	2,3,1	10,12,13	4.25	6 (60%)	9,15,17	1.18	0
2	PIV	D	1	2	5,5,6	1.00	0	6,7,9	0.66	0
2	A1L7L	D	3	2	9,10,11	0.81	0	7,12,14	1.56	1 (14%)
2	PIV	C	1	2	5,5,6	1.05	0	6,7,9	0.71	0
2	A1EAK	C	2	2	8,9,10	2.51	6 (75%)	6,11,13	0.40	0
2	A1L7K	D	4	2,3,1	10,12,13	4.56	6 (60%)	9,15,17	1.60	1 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1EAK	D	2	2	8,9,10	2.42	5 (62%)	6,11,13	0.61	0
2	A1L7L	C	3	2	9,10,11	0.95	1 (11%)	7,12,14	2.06	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1L7K	C	4	2,3,1	-	1/5/17/19	0/1/1/1
2	PIV	D	1	2	-	0/3/3/6	-
2	A1L7L	D	3	2	-	1/5/14/16	0/1/1/1
2	PIV	C	1	2	-	0/3/3/6	-
2	A1EAK	C	2	2	-	2/3/13/15	0/1/1/1
2	A1L7K	D	4	2,3,1	-	1/5/17/19	0/1/1/1
2	A1EAK	D	2	2	-	1/3/13/15	0/1/1/1
2	A1L7L	C	3	2	-	1/5/14/16	0/1/1/1

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	4	A1L7K	CD2-NE2	11.00	1.52	1.33
2	C	4	A1L7K	CD2-NE2	9.90	1.50	1.33
2	D	4	A1L7K	O-C	5.96	1.43	1.19
2	C	4	A1L7K	O-C	5.54	1.42	1.19
2	C	4	A1L7K	CG-CD2	4.86	1.57	1.50
2	D	4	A1L7K	CG-CD2	4.49	1.57	1.50
2	D	4	A1L7K	CF-NE2	3.78	1.54	1.46
2	C	2	A1EAK	C14-C15	-3.38	1.37	1.51
2	D	2	A1EAK	C13-C12	-3.37	1.37	1.51
2	D	2	A1EAK	C14-C15	-3.24	1.38	1.51
2	C	2	A1EAK	C13-C12	-3.15	1.38	1.51
2	C	4	A1L7K	CF-NE2	3.05	1.53	1.46
2	C	2	A1EAK	CB-CA	2.87	1.58	1.53
2	D	2	A1EAK	C12-CB	2.78	1.61	1.53
2	C	2	A1EAK	C15-CB	2.76	1.61	1.53
2	D	2	A1EAK	C14-C13	2.73	1.67	1.48
2	C	2	A1EAK	C14-C13	2.68	1.67	1.48
2	D	4	A1L7K	CD1-CG	-2.58	1.46	1.53
2	C	4	A1L7K	OD2-CD2	-2.50	1.18	1.23
2	D	2	A1EAK	C15-CB	2.47	1.60	1.53
2	C	2	A1EAK	C12-CB	2.37	1.59	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	4	A1L7K	CE1-CD1	-2.26	1.47	1.53
2	C	4	A1L7K	CE1-CD1	-2.26	1.47	1.53
2	C	3	A1L7L	CB-CG	2.25	1.56	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	A1L7L	CG-CB-CA	-4.93	107.89	114.52
2	D	3	A1L7L	C01-N-CA	3.29	123.88	113.64
2	D	4	A1L7K	CF-NE2-CD2	-2.80	117.66	125.79
2	C	3	A1L7L	C01-N-CA	2.00	119.88	113.64

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	A1EAK	C-CA-CB-C12
2	C	2	A1EAK	C-CA-CB-C15
2	C	3	A1L7L	O-C-CA-CB
2	D	3	A1L7L	CA-CB-CG-CD2
2	C	4	A1L7K	O-C-CA-CB
2	D	4	A1L7K	O-C-CA-CB
2	D	2	A1EAK	C-CA-CB-C12

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	A1L7L	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A1L7M	D	101	2	6,6,7	2.31	3 (50%)	6,7,9	2.28	2 (33%)
3	A1L7M	C	101	2	6,6,7	2.30	3 (50%)	6,7,9	1.52	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1L7M	D	101	2	-	0/1/5/6	0/1/1/1
3	A1L7M	C	101	2	-	0/1/5/6	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	101	A1L7M	C5-N	4.18	1.47	1.33
3	D	101	A1L7M	C5-N	4.03	1.46	1.33
3	C	101	A1L7M	C1-C2	2.43	1.54	1.48
3	D	101	A1L7M	C-C2	2.35	1.53	1.48
3	D	101	A1L7M	C1-C2	2.35	1.53	1.48
3	C	101	A1L7M	C-C2	2.27	1.53	1.48

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	101	A1L7M	C-C2-N	-4.66	112.01	118.63
3	D	101	A1L7M	O-C5-N	-2.92	117.58	125.27
3	C	101	A1L7M	C1-C2-N	-2.48	115.11	118.63
3	C	101	A1L7M	C-C2-N	-2.27	115.41	118.63


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/311 (95%)	1.49	87 (29%)  	13, 25, 47, 80	0
1	B	299/311 (96%)	1.15	69 (23%)  	12, 21, 46, 78	0
2	C	0/4	-	-	-	-
2	D	0/4	-	-	-	-
All	All	597/630 (94%)	1.32	156 (26%)  	12, 23, 47, 80	0

All (156) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	223	PHE	8.5
1	B	3	PHE	7.4
1	A	275	GLY	6.5
1	B	300	CYS	6.2
1	B	223	PHE	6.1
1	B	275	GLY	6.1
1	A	154	TYR	5.8
1	B	194	ALA	5.8
1	B	294	PHE	5.6
1	B	193	ALA	5.3
1	A	222	ARG	5.3
1	B	277	ASN	5.1
1	B	154	TYR	4.9
1	A	255	ALA	4.7
1	A	274	ASN	4.6
1	B	274	ASN	4.6
1	A	64	HIS	4.6
1	A	224	THR	4.6
1	B	190	THR	4.4
1	A	59	ILE	4.4
1	B	297	VAL	4.2
1	B	280	THR	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	301	SER	4.1
1	A	212	VAL	4.1
1	A	257	THR	4.1
1	B	50	LEU	4.1
1	A	278	GLY	4.0
1	A	50	LEU	3.8
1	B	276	MET	3.8
1	A	276	MET	3.7
1	B	191	ALA	3.7
1	A	215	GLY	3.6
1	B	222	ARG	3.6
1	A	249	ILE	3.6
1	B	278	GLY	3.6
1	B	24	THR	3.5
1	B	301	SER	3.5
1	A	218	TRP	3.5
1	B	291	PHE	3.5
1	A	78	ILE	3.5
1	A	213	ILE	3.5
1	A	216	ASP	3.5
1	A	300	CYS	3.5
1	A	72	ASN	3.4
1	A	24	THR	3.4
1	B	285	ALA	3.4
1	A	23	GLY	3.4
1	A	259	ILE	3.4
1	A	56	ASP	3.4
1	A	55	GLU	3.4
1	B	283	GLY	3.4
1	B	155	ASP	3.4
1	B	189	GLN	3.3
1	A	220	LEU	3.3
1	A	258	GLY	3.3
1	B	279	ARG	3.3
1	B	59	ILE	3.3
1	A	277	ASN	3.2
1	B	46	SER	3.2
1	A	219	PHE	3.2
1	A	297	VAL	3.2
1	B	226	THR	3.2
1	B	22	CYS	3.2
1	A	280	THR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	282	LEU	3.2
1	A	73	VAL	3.1
1	B	286	LEU	3.1
1	A	298	ARG	3.1
1	B	60	ARG	3.0
1	A	49	MET	3.0
1	B	51	ASN	3.0
1	B	284	SER	3.0
1	B	207	TRP	3.0
1	A	54	TYR	2.9
1	B	299	GLN	2.9
1	A	45	THR	2.9
1	A	260	ALA	2.9
1	A	94	ALA	2.8
1	B	49	MET	2.8
1	B	195	GLY	2.8
1	A	294	PHE	2.8
1	A	279	ARG	2.8
1	A	209	TYR	2.8
1	A	66	PHE	2.8
1	A	43	ILE	2.8
1	A	210	ALA	2.7
1	A	285	ALA	2.7
1	B	196	THR	2.7
1	A	67	LEU	2.7
1	B	58	LEU	2.7
1	A	71	GLY	2.7
1	A	46	SER	2.6
1	A	238	ASN	2.6
1	B	298	ARG	2.6
1	A	139	SER	2.6
1	A	51	ASN	2.6
1	A	57	LEU	2.6
1	A	286	LEU	2.6
1	B	169	THR	2.6
1	B	168	PRO	2.6
1	B	56	ASP	2.5
1	B	221	ASN	2.5
1	A	196	THR	2.5
1	A	251	GLY	2.5
1	B	138	GLY	2.5
1	B	57	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	214	ASN	2.5
1	B	137	LYS	2.5
1	B	4	ARG	2.5
1	A	74	GLN	2.5
1	A	256	GLN	2.5
1	A	153	ASP	2.5
1	B	48	ASP	2.5
1	B	171	VAL	2.5
1	B	53	ASN	2.4
1	A	106	ILE	2.4
1	A	80	HIS	2.4
1	B	282	LEU	2.4
1	A	287	LEU	2.4
1	A	155	ASP	2.4
1	B	140	PHE	2.4
1	B	218	TRP	2.4
1	A	190	THR	2.3
1	A	211	ALA	2.3
1	B	5	LYS	2.3
1	B	188	ARG	2.3
1	A	281	ILE	2.3
1	A	234	ALA	2.3
1	A	77	VAL	2.3
1	A	188	ARG	2.2
1	A	93	THR	2.2
1	A	42	VAL	2.2
1	A	4	ARG	2.2
1	A	75	LEU	2.2
1	A	271	LEU	2.2
1	B	281	ILE	2.2
1	B	224	THR	2.2
1	B	167	LEU	2.2
1	B	287	LEU	2.2
1	A	70	ALA	2.2
1	A	48	ASP	2.1
1	B	153	ASP	2.1
1	B	186	VAL	2.1
1	A	87	LEU	2.1
1	A	253	LEU	2.1
1	A	221	ASN	2.1
1	A	291	PHE	2.1
1	B	184	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	76	ARG	2.0
1	B	134	PHE	2.0
1	A	152	ILE	2.0
1	B	170	GLY	2.0
1	B	296	VAL	2.0
1	A	191	ALA	2.0
1	B	45	THR	2.0
1	B	272	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PIV	D	1	6/7	0.85	0.14	29,30,33,35	0
2	A1EAK	C	2	9/10	0.89	0.16	20,25,39,39	0
2	A1L7L	D	3	10/11	0.91	0.12	20,25,36,38	0
2	A1EAK	D	2	9/10	0.92	0.11	19,26,34,40	0
2	A1L7K	D	4	12/13	0.92	0.10	20,23,26,26	0
2	PIV	C	1	6/7	0.93	0.10	23,26,30,31	0
2	A1L7K	C	4	12/13	0.94	0.08	16,20,22,23	0
2	A1L7L	C	3	10/11	0.94	0.11	20,23,34,36	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	A1L7M	C	101	6/7	0.93	0.10	20,27,29,31	0
3	A1L7M	D	101	6/7	0.97	0.07	18,20,21,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1L7M C 101:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1L7M D 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.