



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 19, 2026 – 02:07 PM EDT

PDB ID : 9NKH / pdb_00009nkh
Title : Conformational flexibility in HLA-B8: peptide tuning structural and dynamical changes
Authors : Li, L.; Bouvier, M.
Deposited on : 2025-02-28
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1306 (1.94-1.94)
Clashscore	180529	1400 (1.94-1.94)
Ramachandran outliers	177936	1387 (1.94-1.94)
Sidechain outliers	177891	1387 (1.94-1.94)
RSRZ outliers	164620	1306 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
1	D	275	
2	B	100	
2	E	100	
3	C	20	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	20	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '15%', a green segment in the middle labeled '45%', and a grey segment on the right labeled '55%'.</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5665 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	Total	C	N	O	S	0	0	0
			1906	1183	350	365	8			
1	D	228	Total	C	N	O	S	0	0	0
			1872	1162	341	361	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	CYS	GLU	engineered mutation	UNP A0A3G6II09
D	76	CYS	GLU	engineered mutation	UNP A0A3G6II09

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	0	0
			836	533	141	158	4			
2	E	100	Total	C	N	O	S	0	0	0
			836	533	141	158	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-PHE-LEU-LYS-LYS-LYS-TYR-CYS-LEU.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	9	Total	C	N	O	S	0	0	0
			73	51	11	10	1			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	F	9	73	51	11	10	1	0	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total	O	0	0
			32	32		
4	B	23	Total	O	0	0
			23	23		
4	D	8	Total	O	0	0
			8	8		
4	E	6	Total	O	0	0
			6	6		



- Molecule 3: ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-PHE-LEU-LYS-LYS-LYS-TYR-CYS-LEU



- Molecule 3: ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-ARG-ALA-PHE-LEU-LYS-LYS-LYS-TYR-CYS-LEU



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.68Å 86.01Å 70.43Å 90.00° 99.59° 90.00°	Depositor
Resolution (Å)	27.02 – 1.94 27.02 – 1.94	Depositor EDS
% Data completeness (in resolution range)	93.8 (27.02-1.94) 93.8 (27.02-1.94)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.93Å)	Xtrriage
Refinement program	PHENIX (1.21.1_5286)	Depositor
R, R_{free}	0.238 , 0.259 0.238 , 0.261	Depositor DCC
R_{free} test set	2760 reflections (4.58%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtrriage
Anisotropy	0.350	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 35.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.024 for l,-k,h	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5665	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/1952	0.47	0/2645
1	D	0.27	0/1919	0.45	0/2603
2	B	0.26	0/859	0.49	0/1162
2	E	0.21	0/859	0.46	0/1162
3	C	0.35	0/74	0.41	0/97
3	F	0.21	0/74	0.38	0/97
All	All	0.26	0/5737	0.46	0/7766

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1906	0	1778	15	0
1	D	1872	0	1730	19	0
2	B	836	0	803	5	0
2	E	836	0	803	13	0
3	C	73	0	79	0	0
3	F	73	0	79	0	0
4	A	32	0	0	0	0
4	B	23	0	0	0	0
4	D	8	0	0	0	0
4	E	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5665	0	5272	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:17:ASN:HD21	2:E:97:ARG:HH12	1.38	0.69
2:E:73:THR:O	2:E:75:LYS:N	2.25	0.69
1:D:82:ARG:NH2	1:D:89:GLU:OE2	2.26	0.68
1:D:126:LEU:HD11	1:D:130:LEU:HA	1.78	0.66
1:A:119:ASP:HB3	2:B:0:MET:HB2	1.83	0.59
1:D:119:ASP:HB3	2:E:0:MET:HG3	1.86	0.58
1:D:5:MET:HB2	1:D:168:LEU:HD13	1.87	0.57
2:E:73:THR:C	2:E:75:LYS:H	2.13	0.56
2:E:17:ASN:ND2	2:E:97:ARG:HH22	2.04	0.55
1:A:33:PHE:CD2	1:A:34:VAL:HG23	2.42	0.54
1:D:239:ARG:N	1:D:239:ARG:HE	2.06	0.54
1:A:5:MET:HB2	1:A:168:LEU:HD13	1.89	0.54
1:D:214:THR:HG23	1:D:262:GLN:HB3	1.91	0.53
2:B:4:THR:HG22	2:B:86:THR:HB	1.90	0.53
1:D:260:HIS:HA	1:D:270:LEU:O	2.09	0.52
2:E:4:THR:HG22	2:E:86:THR:HB	1.92	0.51
2:E:58:LYS:HD3	2:E:58:LYS:H	1.74	0.51
1:A:111:ARG:HG3	1:A:128:GLU:HG2	1.94	0.49
1:D:71:THR:O	1:D:75:ARG:HG3	2.12	0.49
1:A:28:VAL:HG23	1:A:33:PHE:CD1	2.48	0.48
1:D:67:PHE:O	1:D:71:THR:HG23	2.13	0.48
1:D:215:LEU:HD22	1:D:261:VAL:HG22	1.96	0.47
1:A:5:MET:HE3	1:A:33:PHE:CZ	2.50	0.47
1:A:67:PHE:O	1:A:71:THR:HG23	2.15	0.47
1:A:86:ASN:N	1:A:86:ASN:HD22	2.12	0.47
2:E:37:VAL:HG12	2:E:82:VAL:HG13	1.96	0.47
1:D:126:LEU:CD1	1:D:130:LEU:HA	2.44	0.45
1:D:35:ARG:CZ	2:E:53:ASP:HB3	2.46	0.45
2:B:17:ASN:HD21	2:B:74:GLU:HB2	1.82	0.44
1:D:60:TRP:CG	1:D:61:ASP:H	2.36	0.44
2:E:51:HIS:HB3	2:E:66:TYR:CD2	2.54	0.43
1:A:21:ARG:CZ	1:A:23:ILE:HD11	2.48	0.43
1:A:81:LEU:HD13	1:A:118:TYR:CD1	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ILE:HD11	1:A:200:THR:HG22	2.00	0.43
2:B:22:PHE:CE1	2:B:69:GLU:HG3	2.54	0.43
1:D:9:ASP:O	1:D:96:GLN:HA	2.18	0.43
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.54	0.42
1:A:274:TRP:CE2	1:A:275:GLU:HG2	2.55	0.42
1:D:200:THR:HG22	1:D:248:VAL:HG22	2.02	0.42
1:D:219:ARG:HB2	1:D:257:TYR:CE2	2.55	0.41
1:D:220:ASP:OD1	1:D:256:ARG:HD2	2.21	0.41
1:A:230:LEU:HD12	1:D:138:THR:HG21	2.02	0.41
2:E:50:GLU:HB2	2:E:67:TYR:CE1	2.56	0.41
2:E:58:LYS:HB2	2:E:58:LYS:HE2	1.90	0.41
2:E:97:ARG:H	2:E:97:ARG:HG2	1.39	0.41
1:A:183:ASP:N	1:A:183:ASP:OD1	2.54	0.40
1:D:266:LEU:CD2	1:D:270:LEU:HD23	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/275 (81%)	215 (97%)	7 (3%)	0	100	100
1	D	218/275 (79%)	211 (97%)	7 (3%)	0	100	100
2	B	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
2	E	98/100 (98%)	95 (97%)	2 (2%)	1 (1%)	13	5
3	C	7/20 (35%)	7 (100%)	0	0	100	100
3	F	7/20 (35%)	7 (100%)	0	0	100	100
All	All	650/790 (82%)	630 (97%)	19 (3%)	1 (0%)	44	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	74	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/234 (86%)	199 (99%)	2 (1%)	73	68
1	D	197/234 (84%)	194 (98%)	3 (2%)	60	52
2	B	95/95 (100%)	95 (100%)	0	100	100
2	E	95/95 (100%)	95 (100%)	0	100	100
3	C	7/14 (50%)	7 (100%)	0	100	100
3	F	7/14 (50%)	7 (100%)	0	100	100
All	All	602/686 (88%)	597 (99%)	5 (1%)	79	76

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	THR
1	A	200	THR
1	D	88	SER
1	D	197	HIS
1	D	223	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
2	B	2	GLN
2	B	13	HIS
2	B	17	ASN
2	E	13	HIS
2	E	17	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	232/275 (84%)	0.51	11 (4%) 37 41	33, 52, 87, 109	0
1	D	228/275 (82%)	0.79	15 (6%) 26 28	42, 65, 98, 123	0
2	B	100/100 (100%)	0.33	2 (2%) 64 70	31, 51, 82, 121	0
2	E	100/100 (100%)	0.68	6 (6%) 29 31	40, 67, 108, 118	0
3	C	9/20 (45%)	0.84	0 100 100	40, 46, 71, 73	0
3	F	9/20 (45%)	1.69	3 (33%) 1 1	53, 64, 73, 82	0
All	All	678/790 (85%)	0.62	37 (5%) 32 35	31, 59, 96, 123	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	0	ALA	5.1
1	D	60	TRP	4.9
1	A	167	TRP	4.7
1	A	168	LEU	4.6
1	D	36	PHE	4.4
2	E	40	LEU	3.8
1	A	171	TYR	3.5
1	A	109	LEU	2.9
1	A	110	LEU	2.9
3	F	8	LEU	2.7
1	A	225	THR	2.7
1	A	160	LEU	2.7
1	D	168	LEU	2.6
1	D	69	THR	2.6
1	D	165	VAL	2.5
1	D	247	VAL	2.5
2	E	22	PHE	2.5
1	D	182	ALA	2.5
2	E	70	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	167	TRP	2.4
1	A	33	PHE	2.4
1	D	197	HIS	2.3
1	D	20	PRO	2.3
2	E	37	VAL	2.3
1	A	20	PRO	2.3
1	D	228	THR	2.3
2	E	66	TYR	2.3
1	D	142	ILE	2.2
2	B	0	MET	2.1
1	D	201	LEU	2.1
1	A	36	PHE	2.1
1	D	91	GLY	2.1
2	B	47	GLU	2.1
2	E	78	TYR	2.1
3	F	6	TYR	2.0
1	D	274	TRP	2.0
1	A	182	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.