



Full wwPDB X-ray Structure Validation Report i

Dec 8, 2025 – 02:38 pm GMT

PDB ID : 9T5S / pdb_00009t5s
Title : Crystal structure of human PPARalpha in complex with co-activator PGC-1 alpha peptide and bempedoic acid
Authors : Useini, A.; Strater, N.; Kuenze, G.; Heiker, J.T.; Sheikh, B.N.
Deposited on : 2025-11-05
Resolution : 1.74 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1043 (1.74-1.74)
Clashscore	180529	1119 (1.74-1.74)
Ramachandran outliers	177936	1112 (1.74-1.74)
Sidechain outliers	177891	1112 (1.74-1.74)
RSRZ outliers	164620	1043 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 9718 atoms, of which 4758 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	271	Total	C	H	N	O	S	0	10	0
			4459	1413	2250	373	403	20			
1	C	271	Total	C	H	N	O	S	0	8	0
			4421	1402	2233	369	396	21			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	179	MET	-	initiating methionine	UNP Q07869
A	180	GLY	-	expression tag	UNP Q07869
A	181	SER	-	expression tag	UNP Q07869
A	182	SER	-	expression tag	UNP Q07869
A	183	HIS	-	expression tag	UNP Q07869
A	184	HIS	-	expression tag	UNP Q07869
A	185	HIS	-	expression tag	UNP Q07869
A	186	HIS	-	expression tag	UNP Q07869
A	187	HIS	-	expression tag	UNP Q07869
A	188	HIS	-	expression tag	UNP Q07869
A	189	SER	-	expression tag	UNP Q07869
A	190	SER	-	expression tag	UNP Q07869
A	191	GLY	-	expression tag	UNP Q07869
A	192	LEU	-	expression tag	UNP Q07869
A	193	VAL	-	expression tag	UNP Q07869
A	194	PRO	-	expression tag	UNP Q07869
A	195	ARG	-	expression tag	UNP Q07869
A	196	GLY	-	expression tag	UNP Q07869
A	197	SER	-	expression tag	UNP Q07869
A	198	HIS	-	expression tag	UNP Q07869
A	199	MET	-	expression tag	UNP Q07869
C	179	MET	-	initiating methionine	UNP Q07869
C	180	GLY	-	expression tag	UNP Q07869
C	181	SER	-	expression tag	UNP Q07869
C	182	SER	-	expression tag	UNP Q07869

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	183	HIS	-	expression tag	UNP Q07869
C	184	HIS	-	expression tag	UNP Q07869
C	185	HIS	-	expression tag	UNP Q07869
C	186	HIS	-	expression tag	UNP Q07869
C	187	HIS	-	expression tag	UNP Q07869
C	188	HIS	-	expression tag	UNP Q07869
C	189	SER	-	expression tag	UNP Q07869
C	190	SER	-	expression tag	UNP Q07869
C	191	GLY	-	expression tag	UNP Q07869
C	192	LEU	-	expression tag	UNP Q07869
C	193	VAL	-	expression tag	UNP Q07869
C	194	PRO	-	expression tag	UNP Q07869
C	195	ARG	-	expression tag	UNP Q07869
C	196	GLY	-	expression tag	UNP Q07869
C	197	SER	-	expression tag	UNP Q07869
C	198	HIS	-	expression tag	UNP Q07869
C	199	MET	-	expression tag	UNP Q07869

- Molecule 2 is a protein called Peroxisome proliferator-activated receptor gamma coactivator 1-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	12	Total	C	H	N	O	0	0	0
			198	60	109	15	14			
2	D	10	Total	C	H	N	O	0	0	0
			174	53	98	12	11			

- Molecule 3 is Bempedoic acid (CCD ID: A1JTY) (formula: C₁₉H₃₆O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			58	19	34	5		

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	H	O	0	0
			58	19	34	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	192	Total	O			0	10
			197	197				
4	B	6	Total	O			0	1
			7	7				
4	C	141	Total	O			0	4
			145	145				
4	D	1	Total	O			0	0
			1	1				


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peroxisome proliferator-activated receptor alpha




- Molecule 1: Peroxisome proliferator-activated receptor alpha



- Molecule 2: Peroxisome proliferator-activated receptor gamma coactivator 1-alpha



- Molecule 2: Peroxisome proliferator-activated receptor gamma coactivator 1-alpha



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.25 \AA 99.75 \AA 61.61 \AA 90.00° 99.80° 90.00°	Depositor
Resolution (\AA)	38.54 – 1.74 38.54 – 1.74	Depositor EDS
% Data completeness (in resolution range)	79.1 (38.54-1.74) 79.1 (38.54-1.74)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.02 (at 1.74 \AA)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.178, 0.211 0.178, 0.209	Depositor DCC
R_{free} test set	2988 reflections (4.06%)	wwPDB-VP
Wilson B-factor (\AA^2)	26.3	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/ \AA^3), B_{sol} (\AA^2)	0.41, 35.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.020 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9718	wwPDB-VP
Average B, all atoms (\AA^2)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: A1JTY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2273	0.48	0/3062
1	C	0.24	0/2233	0.49	0/3009
2	B	0.20	0/89	0.37	0/119
2	D	0.20	0/76	0.41	0/101
All	All	0.24	0/4671	0.48	0/6291

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2209	2250	2214	13	0
1	C	2188	2233	2216	10	0
2	B	89	109	108	0	0
2	D	76	98	97	1	0
3	A	24	34	0	0	0
3	C	24	34	0	1	0
4	A	197	0	0	0	0
4	B	7	0	0	0	0
4	C	145	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	1	0	0	0	0
All	All	4960	4758	4635	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263[A]:ILE:HD13	1:A:274[A]:HIS:CG	2.22	0.73
1:A:321:LEU:HD21	1:A:355:MET:HE1	1.75	0.68
1:A:321:LEU:HD11	1:A:330:MET:HE1	1.76	0.67
1:C:206:LEU:HD23	1:C:407:VAL:HG21	1.77	0.66
1:A:258:LEU:HD23	1:A:263[B]:ILE:HD11	1.85	0.58
1:A:300:LEU:HD21	1:A:400:MET:HE1	1.87	0.56
1:C:339:ILE:HD12	3:C:501:A1JTY:C20	2.36	0.55
1:A:299:ASN:ND2	1:C:446:ILE:HD13	2.21	0.54
1:A:263[A]:ILE:HD13	1:A:274[A]:HIS:ND1	2.27	0.49
1:A:263[A]:ILE:HG22	1:A:263[A]:ILE:O	2.12	0.49
1:A:258:LEU:HD23	1:A:263[B]:ILE:CD1	2.41	0.49
1:C:203:LEU:HD22	1:C:407:VAL:HG22	1.97	0.47
1:C:220[A]:MET:HE1	1:C:335:GLY:HA3	1.96	0.47
1:A:423:PHE:HB3	1:A:424:PRO:HD3	1.96	0.46
1:A:384:CYS:O	1:A:401:GLN:HB2	2.14	0.46
1:A:263[A]:ILE:O	1:A:263[A]:ILE:CG2	2.63	0.46
1:C:423:PHE:HB3	1:C:424:PRO:HD3	1.98	0.45
1:C:403:GLY:O	1:C:407:VAL:HG23	2.17	0.45
1:A:330:MET:SD	1:A:355:MET:HE2	2.56	0.45
1:C:370:LEU:HD11	1:C:426:LEU:HD21	2.00	0.44
1:C:302:LEU:HD22	2:D:695:LEU:HD21	2.01	0.43
1:C:456:LEU:HB2	1:C:461:GLN:HG2	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	279/290 (96%)	274 (98%)	5 (2%)	0	100 100
1	C	275/290 (95%)	272 (99%)	3 (1%)	0	100 100
2	B	10/19 (53%)	10 (100%)	0	0	100 100
2	D	8/19 (42%)	8 (100%)	0	0	100 100
All	All	572/618 (93%)	564 (99%)	8 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	247/253 (98%)	247 (100%)	0	100 100
1	C	242/253 (96%)	242 (100%)	0	100 100
2	B	10/16 (62%)	10 (100%)	0	100 100
2	D	9/16 (56%)	9 (100%)	0	100 100
All	All	508/538 (94%)	508 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	219	ASN
1	A	299	ASN
1	A	393	ASN
2	B	699	ASN
1	C	219	ASN
1	C	261	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A1JTY	C	501	-	23,23,23	0.63	0	31,31,31	2.19	5 (16%)
3	A1JTY	A	501	-	23,23,23	0.86	1 (4%)	31,31,31	1.38	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1JTY	C	501	-	-	11/30/30/30	-
3	A1JTY	A	501	-	-	18/30/30/30	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	A1JTY	C3-C2	3.06	1.55	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	501	A1JTY	O22-C21-C18	-8.62	110.59	122.87
3	C	501	A1JTY	O24-C2-C3	-6.52	113.58	122.87
3	A	501	A1JTY	O22-C21-C18	-4.92	115.86	122.87
3	A	501	A1JTY	O24-C2-C3	-4.53	116.41	122.87
3	C	501	A1JTY	O23-C21-C18	2.99	120.54	114.30
3	C	501	A1JTY	C6-C3-C2	2.44	113.23	108.15
3	C	501	A1JTY	O1-C2-C3	2.43	119.38	114.30
3	A	501	A1JTY	C6-C3-C2	2.17	112.66	108.15

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	A1JTY	C5-C3-C6-C7
3	A	501	A1JTY	C2-C3-C6-C7
3	A	501	A1JTY	C17-C18-C21-O23
3	C	501	A1JTY	O24-C2-C3-C6
3	A	501	A1JTY	C9-C10-C11-O12
3	A	501	A1JTY	C9-C10-C11-C13
3	A	501	A1JTY	C3-C6-C7-C8
3	C	501	A1JTY	C4-C3-C6-C7
3	C	501	A1JTY	C5-C3-C6-C7
3	A	501	A1JTY	C6-C7-C8-C9
3	C	501	A1JTY	C6-C7-C8-C9
3	A	501	A1JTY	C20-C18-C21-O22
3	A	501	A1JTY	C20-C18-C21-O23
3	A	501	A1JTY	C19-C18-C21-O23
3	A	501	A1JTY	O1-C2-C3-C6
3	A	501	A1JTY	O24-C2-C3-C6
3	C	501	A1JTY	O1-C2-C3-C4
3	C	501	A1JTY	O1-C2-C3-C5
3	C	501	A1JTY	O1-C2-C3-C6
3	C	501	A1JTY	C7-C8-C9-C10
3	A	501	A1JTY	C4-C3-C6-C7
3	C	501	A1JTY	C2-C3-C6-C7
3	C	501	A1JTY	C11-C13-C14-C15
3	A	501	A1JTY	O1-C2-C3-C5
3	C	501	A1JTY	O24-C2-C3-C4
3	A	501	A1JTY	C11-C10-C9-C8
3	A	501	A1JTY	C17-C18-C21-O22
3	A	501	A1JTY	C19-C18-C21-O22

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	501	A1JTY	O24-C2-C3-C5


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	501	A1JTY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	271/290 (93%)	-0.25	7 (2%) 57 64	14, 28, 49, 80	7 (2%)
1	C	271/290 (93%)	-0.02	7 (2%) 57 64	16, 31, 55, 84	5 (1%)
2	B	12/19 (63%)	0.10	1 (8%) 19 24	29, 38, 50, 72	0
2	D	10/19 (52%)	1.59	3 (30%) 1 1	41, 55, 73, 102	0
All	All	564/618 (91%)	-0.10	18 (3%) 50 58	14, 30, 55, 102	12 (2%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	697	PRO	7.2
1	C	468	TYR	5.0
1	A	263[A]	ILE	3.2
2	B	699	ASN	3.1
1	A	233	ALA	2.9
1	C	274[A]	HIS	2.8
1	C	396	HIS	2.8
2	D	696	ALA	2.7
1	A	198	HIS	2.4
1	A	231	GLY	2.4
1	A	274[A]	HIS	2.3
1	A	226[A]	ARG	2.3
1	C	198	HIS	2.2
1	C	467	MET	2.2
1	A	219	ASN	2.2
1	C	466	ASP	2.1
1	C	387	ASP	2.1
2	D	695	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no oligosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A1JTY	A	501	24/24	0.88	0.15	27,50,71,80	0
3	A1JTY	C	501	24/24	0.91	0.12	25,46,60,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.