



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 02:32 AM EDT

PDB ID : 1B49  
Title : DCMP HYDROXYMETHYLASE FROM T4 (PHOSPHATE-BOUND)  
Authors : Song, H.K.; Sohn, S.H.; Suh, S.W.  
Deposited on : 1999-01-06  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	 % 81% 15% . .
1	C	246	 % 80% 15% . .

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (DEOXYCYTIDYLATE HYDROXYMETHYLASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	Total 1972	1261	329	372	10	0	0	0
1	C	241	Total 1972	1261	329	372	10	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			O	P		
2	A	1	Total 5	4 1	0	0
2	C	1	Total 5	4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	68	Total 68	O 68	0	0
3	C	70	Total 70	O 70	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (DEOXYCYTIDYLATE HYDROXYMETHYLASE)



- Molecule 1: PROTEIN (DEOXYCYTIDYLATE HYDROXYMETHYLASE)



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	176.34Å 53.27Å 75.20Å 90.00° 115.21° 90.00°	Depositor
Resolution (Å)	8.00 – 2.30 15.86 – 2.30	Depositor EDS
% Data completeness (in resolution range)	91.2 (8.00-2.30) 92.5 (15.86-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.32 (at 2.29Å)	Xtrriage
Refinement program	X-PLOR 3.843	Depositor
R, $R_{free}$	0.159 , 0.202 0.157 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.9	Xtrriage
Anisotropy	0.317	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 25.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.477 for -h-2*1,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4092	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/2021	0.49	0/2733
1	C	0.37	0/2021	0.49	0/2733
All	All	0.37	0/4042	0.49	0/5466

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	56	TYR	Sidechain
1	C	56	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1972	0	1908	26	0
1	C	1972	0	1908	25	0
2	A	5	0	0	1	0
2	C	5	0	0	1	0
3	A	68	0	0	3	0
3	C	70	0	0	4	0
All	All	4092	0	3816	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:ASN:ND2	1:C:166:ASN:ND2	2.33	0.77
1:A:153:GLN:HE22	1:C:213:GLY:HA3	1.59	0.68
1:C:124:ARG:HD2	3:C:367:HOH:O	1.98	0.64
1:A:150:ASN:HB2	1:A:166:ASN:ND2	2.12	0.64
1:A:161:ILE:HD12	1:A:191:VAL:HG22	1.82	0.60
1:A:227:HIS:CG	1:A:236:ILE:HG12	2.38	0.59
1:A:153:GLN:NE2	1:C:213:GLY:HA3	2.19	0.57
1:A:213:GLY:HA3	1:C:153:GLN:NE2	2.20	0.57
1:C:150:ASN:HB2	1:C:166:ASN:ND2	2.20	0.57
1:A:213:GLY:HA3	1:C:153:GLN:HE22	1.70	0.56
1:C:168:ARG:NH2	2:C:301:PO4:O2	2.37	0.56
1:A:85:VAL:HG22	1:A:144:SER:O	2.05	0.56
1:A:124:ARG:HD2	3:A:366:HOH:O	2.06	0.55
1:C:85:VAL:HG22	1:C:144:SER:O	2.07	0.55
1:C:227:HIS:CG	1:C:236:ILE:HG12	2.42	0.54
1:A:168:ARG:NH2	2:A:300:PO4:O2	2.37	0.53
1:A:155:LEU:HD13	1:C:37:ILE:HG23	1.89	0.53
1:C:139:ASN:HA	1:C:143:MET:O	2.08	0.52
1:C:61:LEU:HB2	3:C:317:HOH:O	2.09	0.52
1:A:64:TYR:O	1:A:97:GLY:HA3	2.11	0.51
1:A:37:ILE:HG23	1:C:155:LEU:HD13	1.91	0.51
1:C:53:ASN:C	1:C:53:ASN:HD22	2.15	0.50
1:C:81:ILE:HG13	1:C:82:TRP:N	2.26	0.49
1:C:118:GLN:NE2	3:C:370:HOH:O	2.46	0.49
1:C:64:TYR:O	1:C:97:GLY:HA3	2.13	0.49
1:A:5:SER:HB2	1:A:42:VAL:HG23	1.95	0.48
1:A:53:ASN:HD22	1:A:53:ASN:C	2.18	0.47
1:A:118:GLN:NE2	3:A:368:HOH:O	2.48	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:ILE:HG12	1:A:161:ILE:HG12	1.96	0.47
1:A:139:ASN:HA	1:A:143:MET:O	2.15	0.46
1:A:81:ILE:HG13	1:A:82:TRP:N	2.31	0.46
1:A:2:ILE:HD12	1:A:2:ILE:N	2.31	0.46
1:C:182:TRP:O	1:C:186:VAL:HG23	2.15	0.46
1:A:8:VAL:HG13	1:A:48:ILE:HG12	1.98	0.45
1:C:100:ILE:HA	1:C:108:GLN:HB2	1.98	0.45
1:A:100:ILE:HA	1:A:108:GLN:HB2	1.98	0.44
1:C:17:LEU:HD13	1:C:229:TRP:CH2	2.52	0.43
1:A:237:SER:HB3	1:A:239:LYS:HE2	1.99	0.43
1:A:227:HIS:ND1	1:A:236:ILE:HG12	2.34	0.43
1:A:221:HIS:HD2	3:A:311:HOH:O	2.02	0.42
1:C:5:SER:HB2	1:C:42:VAL:HG23	2.03	0.41
1:C:156:ILE:HG12	1:C:161:ILE:HG12	2.02	0.40
1:C:221:HIS:HD2	3:C:313:HOH:O	2.03	0.40
1:C:223:TYR:HB2	1:C:241:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/246 (97%)	231 (97%)	8 (3%)	0	100	100
1	C	239/246 (97%)	232 (97%)	7 (3%)	0	100	100
All	All	478/492 (97%)	463 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/217 (99%)	201 (94%)	13 (6%)	18	25
1	C	214/217 (99%)	201 (94%)	13 (6%)	18	25
All	All	428/434 (99%)	402 (94%)	26 (6%)	18	25

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ASP
1	A	17	LEU
1	A	22	LYS
1	A	29	THR
1	A	53	ASN
1	A	85	VAL
1	A	103	GLU
1	A	124	ARG
1	A	147	MET
1	A	171	ASP
1	A	236	ILE
1	A	237	SER
1	A	240	ASP
1	C	4	ASP
1	C	17	LEU
1	C	22	LYS
1	C	29	THR
1	C	53	ASN
1	C	85	VAL
1	C	103	GLU
1	C	124	ARG
1	C	147	MET
1	C	171	ASP
1	C	236	ILE
1	C	237	SER
1	C	240	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	A	58	GLN
1	A	83	GLN
1	A	84	GLN
1	A	118	GLN
1	A	119	ASN
1	A	153	GLN
1	A	166	ASN
1	A	183	GLN
1	A	221	HIS
1	C	53	ASN
1	C	58	GLN
1	C	83	GLN
1	C	84	GLN
1	C	118	GLN
1	C	119	ASN
1	C	153	GLN
1	C	166	ASN
1	C	183	GLN
1	C	221	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	C	301	-	4,4,4	1.51	1 (25%)	6,6,6	0.78	0
2	PO4	A	300	-	4,4,4	1.16	1 (25%)	6,6,6	0.73	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	PO4	P-O1	2.99	1.57	1.50
2	A	300	PO4	P-O1	2.29	1.55	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	301	PO4	1	0
2	A	300	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	241/246 (97%)	-0.66	3 (1%) 79 83	6, 17, 55, 86	0
1	C	241/246 (97%)	-0.68	2 (0%) 86 89	5, 17, 52, 85	0
All	All	482/492 (97%)	-0.67	5 (1%) 82 86	5, 17, 55, 86	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	28	LYS	4.9
1	C	28	LYS	4.6
1	A	29	THR	3.9
1	C	29	THR	2.6
1	A	31	VAL	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	A	300	5/5	0.95	0.12	47,50,52,53	0
2	PO4	C	301	5/5	0.95	0.12	46,46,53,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.