



Full wwPDB NMR Structure Validation Report ⓘ

Mar 8, 2026 – 02:32 PM UTC

PDB ID : 1C7W / pdb_00001c7w
Title : NMR SOLUTION STRUCTURE OF THE CALCIUM-BOUND C-TERMINAL DOMAIN (W81-S161) OF CALCIUM VECTOR PROTEIN FROM AMPHIOXUS
Authors : Theret, I.; Baladi, S.; Cox, J.A.; Sakamoto, H.; Craescu, C.T.
Deposited on : 2000-03-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	81	

2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1049 atoms, of which 510 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CALCIUM VECTOR PROTEIN.


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	68	1049	341	510	80	115	3	0

4 Residue-property plots [i](#)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: CALCIUM VECTOR PROTEIN

Chain A: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *Distance Geometry (DGII) and Simulated Annealing (Discover)*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Felix	structure solution	97.0
Felix	refinement	97.0

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.55	0/546 (0.0%)	2.84	45/731 (6.2%)
All	All	0.55	0/546 (0.0%)	2.84	45/731 (6.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0	1
All	All	0	1

There are no bond-length outliers.

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	GLU	O-C-N	-33.65	80.88	122.27
1	A	122	LEU	N-CA-C	11.37	127.82	109.85
1	A	110	PHE	CA-CB-CG	10.80	124.60	113.80
1	A	116	LYS	N-CA-CB	-10.38	93.97	111.31
1	A	116	LYS	N-CA-C	9.54	122.31	108.86
1	A	143	ASP	CA-CB-CG	9.35	121.95	112.60
1	A	121	PRO	N-CA-CB	-8.99	93.81	103.25
1	A	144	ILE	N-CA-CB	8.71	115.98	110.50
1	A	102	ASP	CA-CB-CG	8.43	121.03	112.60
1	A	122	LEU	N-CA-CB	-8.34	97.68	111.20
1	A	127	VAL	CA-CB-CG1	8.29	124.50	110.40
1	A	124	ASP	CA-CB-CG	7.89	120.49	112.60
1	A	139	ASN	CA-CB-CG	7.59	120.19	112.60
1	A	115	GLN	CA-CB-CG	7.46	129.03	114.10
1	A	88	GLU	CA-C-N	7.17	132.50	120.88
1	A	88	GLU	C-N-CA	7.17	132.50	120.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	144	ILE	CA-C-O	-7.07	112.84	118.85
1	A	139	ASN	OD1-CG-ND2	-6.86	115.74	122.60
1	A	137	ASP	CA-CB-CG	6.68	119.28	112.60
1	A	121	PRO	N-CD-CG	-6.50	93.45	103.20
1	A	148	MET	CA-CB-CG	6.31	126.73	114.10
1	A	106	ASP	CA-CB-CG	6.06	118.66	112.60
1	A	152	LYS	N-CA-C	5.92	120.37	113.20
1	A	147	PHE	CA-CB-CG	5.90	119.70	113.80
1	A	144	ILE	CB-CA-C	-5.88	108.11	114.35
1	A	142	ILE	CA-CB-CG1	5.86	120.36	110.40
1	A	109	GLU	CB-CG-CD	5.83	122.50	112.60
1	A	135	ASP	CA-CB-CG	5.62	118.22	112.60
1	A	96	VAL	CA-CB-CG1	5.56	119.85	110.40
1	A	152	LYS	CA-CB-CG	5.51	125.12	114.10
1	A	134	ALA	N-CA-C	5.46	119.66	112.89
1	A	88	GLU	CA-C-O	-5.45	113.70	119.97
1	A	98	ASP	CA-CB-CG	5.45	118.05	112.60
1	A	104	VAL	CA-CB-CG1	5.44	119.65	110.40
1	A	97	PHE	N-CA-C	5.40	117.92	111.71
1	A	127	VAL	CB-CA-C	5.39	119.66	112.22
1	A	99	ALA	N-CA-C	5.34	116.96	111.03
1	A	90	ILE	CA-C-O	-5.25	115.22	121.05
1	A	114	MET	N-CA-C	5.22	121.92	110.80
1	A	93	ALA	N-CA-C	5.18	116.78	111.03
1	A	93	ALA	CA-C-O	-5.16	115.58	120.90
1	A	115	GLN	OE1-CD-NE2	-5.10	117.50	122.60
1	A	100	ASN	CB-CG-ND2	5.05	123.98	116.40
1	A	150	LEU	CA-CB-CG	5.04	133.95	116.30
1	A	115	GLN	CB-CA-C	-5.00	100.46	110.42

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	88	GLU	Mainchain

6.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes

averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	539	510	509	4
All	All	539	510	509	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:114:MET:HG3	1:A:115:GLN:HG2	0.48	1.84
1:A:90:ILE:HG23	1:A:147:PHE:CE1	0.45	2.45
1:A:90:ILE:HG23	1:A:147:PHE:HE1	0.45	1.72
1:A:99:ALA:CB	1:A:109:GLU:HG3	0.40	2.47

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	66/81 (81%)	59 (89%)	6 (9%)	1 (2%)	11	57
All	All	66/81 (81%)	59 (89%)	6 (9%)	1 (2%)	11	57

All 1 Ramachandran outliers are listed below.

Mol	Chain	Res	Type
1	A	121	PRO

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	58/70 (83%)	46 (79%)	12 (21%)	3	32
All	All	58/70 (83%)	46 (79%)	12 (21%)	3	32

All 12 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	91	LEU
1	A	96	VAL
1	A	104	VAL
1	A	116	LYS
1	A	122	LEU
1	A	127	VAL
1	A	131	MET
1	A	141	VAL
1	A	142	ILE
1	A	150	LEU
1	A	152	LYS
1	A	154	SER

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided