



# Full wwPDB NMR Structure Validation Report ⓘ

Mar 5, 2026 – 10:01 PM UTC

PDB ID : 1GJH / pdb\_00001gjh  
Title : HUMAN BCL-2, ISOFORM 2  
Authors : Petros, A.M.; Medek, A.; Nettlesheim, D.G.; Kim, D.H.; Yoon, H.S.; Swift, K.;  
Matayoshi, E.D.; Oltersdorf, T.; Fesik, S.W.  
Deposited on : 2001-05-31

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	166	

## 2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2609 atoms, of which 1259 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called PROTEIN (APOPTOSIS REGULATOR BCL-2 WITH PUTATIVE FLEXIBLE LOOP REPLACED WITH A PORTION OF APOPTOSIS REGULATOR BCL-X PROTEIN).

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	164	2609	847	1259	242	254	7	0

## 4 Residue-property plots [i](#)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PROTEIN (APOPTOSIS REGULATOR BCL-2 WITH PUTATIVE FLEXIBLE LOOP REPLACED WITH A PORTION OF APOPTOSIS REGULATOR BCL-X PROTEIN)



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNX	refinement	2000

No chemical shift data was provided.

## 6 Model quality

### 6.1 Standard geometry

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1350	1259	1258	44
All	All	1350	1259	1258	44

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:119:LEU:HD21	1:A:157:MET:HE3	0.85	1.45
1:A:181:LEU:HG	1:A:185:LEU:HD12	0.70	1.64
1:A:162:VAL:HG22	1:A:167:SER:HA	0.68	1.65
1:A:167:SER:O	1:A:170:VAL:HG13	0.64	1.93
1:A:30:TRP:CE2	1:A:32:ALA:HB2	0.62	2.29
1:A:7:THR:HG22	1:A:48:GLU:HG2	0.59	1.75
1:A:15:VAL:HG12	1:A:19:ILE:CD1	0.57	2.30
1:A:95:LEU:HD23	1:A:95:LEU:N	0.54	2.18
1:A:169:LEU:CD2	1:A:173:ILE:HD11	0.52	2.35
1:A:112:PHE:CZ	1:A:149:ALA:HB1	0.51	2.41
1:A:19:ILE:HD12	1:A:174:ALA:HB2	0.51	1.82
1:A:104:PHE:CZ	1:A:108:TYR:CE1	0.50	3.00
1:A:147:ILE:HD13	1:A:189:ILE:HD11	0.50	1.84
1:A:129:ARG:O	1:A:133:VAL:HG23	0.49	2.08
1:A:36:VAL:O	1:A:36:VAL:HG13	0.49	2.08

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:133:VAL:HG11	1:A:153:PHE:CE1	0.48	2.44
1:A:48:GLU:O	1:A:92:VAL:HG12	0.48	2.07
1:A:19:ILE:CD1	1:A:174:ALA:HB2	0.48	2.39
1:A:30:TRP:CD2	1:A:32:ALA:HB2	0.47	2.44
1:A:11:ASN:O	1:A:15:VAL:HG23	0.47	2.10
1:A:26:ARG:NH2	1:A:159:VAL:HG21	0.47	2.25
1:A:173:ILE:HG22	1:A:177:MET:SD	0.47	2.50
1:A:162:VAL:HG13	1:A:167:SER:OG	0.46	2.09
1:A:14:ILE:CD1	1:A:195:TRP:CZ2	0.45	3.00
1:A:17:LYS:CE	1:A:94:HIS:CG	0.45	3.00
1:A:96:THR:HG22	1:A:198:PHE:CE2	0.45	2.46
1:A:36:VAL:O	1:A:36:VAL:HG22	0.45	2.12
1:A:130:PHE:CD1	1:A:131:ALA:N	0.45	2.84
1:A:151:PHE:CZ	1:A:181:LEU:CD1	0.45	3.00
1:A:142:VAL:HG12	1:A:143:ASN:N	0.44	2.28
1:A:14:ILE:CD1	1:A:195:TRP:CE2	0.44	3.01
1:A:112:PHE:HZ	1:A:149:ALA:HB1	0.44	1.72
1:A:169:LEU:HD23	1:A:173:ILE:HG12	0.43	1.91
1:A:189:ILE:HG22	1:A:190:GLN:N	0.43	2.27
1:A:7:THR:HG22	1:A:48:GLU:CG	0.43	2.42
1:A:42:GLU:O	1:A:43:ALA:HB3	0.42	2.15
1:A:181:LEU:CG	1:A:185:LEU:HD12	0.41	2.43
1:A:21:TYR:CD1	1:A:21:TYR:C	0.41	2.97
1:A:180:TYR:CD1	1:A:180:TYR:C	0.41	2.98
1:A:173:ILE:CD1	1:A:176:TRP:CZ3	0.41	3.03
1:A:186:HIS:CD2	1:A:186:HIS:C	0.41	2.99
1:A:49:SER:O	1:A:94:HIS:CD2	0.40	2.74
1:A:150:PHE:HE1	1:A:181:LEU:HD12	0.40	1.77
1:A:96:THR:CG2	1:A:198:PHE:CE2	0.40	3.04

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	162/166 (98%)	131 (81%)	23 (14%)	8 (5%)	<b>3</b> <b>24</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	162/166 (98%)	131 (81%)	23 (14%)	8 (5%)	3 24

All 8 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	4	ALA
1	A	38	GLU
1	A	41	THR
1	A	48	GLU
1	A	111	ASP
1	A	165	GLU
1	A	204	PRO
1	A	205	SER

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	140/141 (99%)	96 (69%)	44 (31%)	1 14
All	All	140/141 (99%)	96 (69%)	44 (31%)	1 14

All 44 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	3	HIS
1	A	6	ARG
1	A	14	ILE
1	A	17	LYS
1	A	21	TYR
1	A	23	LEU
1	A	24	SER
1	A	25	GLN
1	A	34	ASP
1	A	40	ARG
1	A	41	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	42	GLU
1	A	48	GLU
1	A	95	LEU
1	A	97	LEU
1	A	98	ARG
1	A	105	SER
1	A	106	ARG
1	A	109	ARG
1	A	110	ARG
1	A	111	ASP
1	A	118	GLN
1	A	119	LEU
1	A	120	HIS
1	A	121	LEU
1	A	122	THR
1	A	130	PHE
1	A	132	THR
1	A	135	GLU
1	A	137	LEU
1	A	139	ARG
1	A	148	VAL
1	A	150	PHE
1	A	151	PHE
1	A	158	CYS
1	A	159	VAL
1	A	161	SER
1	A	164	ARG
1	A	167	SER
1	A	178	THR
1	A	187	THR
1	A	189	ILE
1	A	199	VAL
1	A	200	GLU

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided