



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:10 PM UTC

PDB ID : 1H6N / pdb_00001h6n
Title : Formation of a tyrosyl radical intermediate in *Proteus mirabilis* catalase by directed mutagenesis and consequences for nucleotide reactivity
Authors : Andreoletti, P.; Sainz, G.; Jaquinod, M.; Gagnon, J.; Jouve, H.M.
Deposited on : 2001-06-20
Resolution : 2.11 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	OMT	A	53	-	-	X	-



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	4	2	2	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	325	Total 325	O 325	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CATALASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	109.90Å 109.90Å 249.78Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.67 – 2.11 29.67 – 2.11	Depositor EDS
% Data completeness (in resolution range)	97.9 (29.67-2.11) 98.0 (29.67-2.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.12Å)	Xtrriage
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.220 , 0.236 0.211 , 0.228	Depositor DCC
R_{free} test set	2554 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtrriage
Anisotropy	0.685	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4272	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: OMT, HEM, SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	1/3995 (0.0%)	0.97	19/5404 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	359	CYS	C-N	7.41	1.42	1.33

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	336	SER	N-CA-C	8.27	121.04	111.11
1	A	419	PHE	N-CA-C	7.04	122.54	113.88
1	A	95	VAL	N-CA-C	6.58	116.82	111.62
1	A	226	ASP	N-CA-C	6.57	118.99	111.11
1	A	195	GLY	N-CA-C	-6.49	106.61	114.66
1	A	149	ARG	N-CA-C	6.31	120.56	109.96
1	A	47	VAL	N-CA-C	6.09	118.12	108.87
1	A	414	GLU	N-CA-C	-6.08	105.13	112.54
1	A	163	ASP	N-CA-C	-5.65	105.02	111.07
1	A	134	LEU	N-CA-C	5.60	118.82	110.14
1	A	97	GLY	N-CA-C	5.36	118.09	111.93
1	A	80	SER	N-CA-C	5.24	118.46	111.75
1	A	184	LEU	CA-C-N	5.12	125.40	119.92
1	A	184	LEU	C-N-CA	5.12	125.40	119.92
1	A	463	VAL	N-CA-C	-5.09	105.88	110.82
1	A	94	THR	N-CA-C	-5.09	101.95	109.94
1	A	96	ALA	N-CA-C	5.08	117.94	111.69
1	A	376	SER	N-CA-C	5.08	119.46	113.16
1	A	266	LYS	N-CA-C	5.03	117.50	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3895	0	3705	49	0
2	A	43	0	30	1	0
3	A	4	0	3	0	0
4	A	5	0	0	0	0
5	A	325	0	0	9	0
All	All	4272	0	3738	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:LYS:HE3	1:A:266:LYS:HA	1.20	1.16
1:A:359:CYS:HB2	1:A:360:PRO:HD2	1.42	1.00
1:A:266:LYS:HA	1:A:266:LYS:CE	2.01	0.89
1:A:17:ASN:HD22	1:A:35:TRP:HE1	1.22	0.86
1:A:266:LYS:HE3	1:A:266:LYS:CA	2.05	0.81
1:A:464:HIS:HD2	1:A:466:GLU:H	1.38	0.72
1:A:53:OMT:HE1	1:A:144:ASN:HD22	1.59	0.67
1:A:429:LEU:HD23	1:A:437:MET:HE1	1.79	0.65
1:A:139:LYS:HE2	1:A:139:LYS:HA	1.80	0.63
1:A:359:CYS:HB2	1:A:360:PRO:CD	2.25	0.61
1:A:423:ARG:HD3	1:A:463:VAL:O	2.03	0.59
1:A:124:MET:HE2	1:A:215:PHE:CE1	2.38	0.58
1:A:433:GLU:HG2	1:A:437:MET:CE	2.34	0.57
1:A:254:PRO:HG2	1:A:297:LEU:HB2	1.88	0.56
1:A:278:LEU:HD12	1:A:329:MET:HG3	1.85	0.56
1:A:132:PHE:HB3	1:A:278:LEU:HD13	1.89	0.55
1:A:349:HIS:HD2	5:A:2241:HOH:O	1.89	0.54
1:A:53:OMT:CE	1:A:144:ASN:HD22	2.19	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:OMT:HE1	1:A:144:ASN:ND2	2.21	0.53
1:A:433:GLU:HG2	1:A:437:MET:HE3	1.91	0.53
1:A:17:ASN:ND2	1:A:35:TRP:HE1	1.97	0.52
1:A:310:GLN:HB3	1:A:352:ILE:HD13	1.90	0.52
1:A:53:OMT:HE2	1:A:140:PHE:HE2	1.75	0.52
1:A:135:ARG:HG2	5:A:2188:HOH:O	2.10	0.52
1:A:299:ARG:NH2	5:A:2201:HOH:O	2.42	0.51
1:A:266:LYS:HE2	5:A:2183:HOH:O	2.09	0.51
1:A:17:ASN:HD21	1:A:39:LYS:NZ	2.09	0.51
1:A:53:OMT:HE2	1:A:140:PHE:CE2	2.46	0.50
1:A:53:OMT:HG3	5:A:2043:HOH:O	2.13	0.48
1:A:23:ALA:O	1:A:27:GLY:HA3	2.14	0.48
1:A:464:HIS:CD2	1:A:466:GLU:HB2	2.49	0.48
1:A:208:ARG:NH2	1:A:400:PRO:O	2.47	0.47
1:A:148:LYS:O	1:A:156:ARG:HG3	2.15	0.47
1:A:180:SER:HA	1:A:437:MET:HE2	1.97	0.47
1:A:179:MET:HG3	1:A:441:ILE:HG12	1.97	0.46
1:A:53:OMT:HE3	1:A:53:OMT:HB3	1.75	0.45
1:A:91:ARG:HD3	2:A:600:HEM:O1D	2.16	0.45
1:A:450:LYS:HD3	1:A:450:LYS:HA	1.77	0.44
1:A:192:HIS:CD2	1:A:221:ILE:HD11	2.54	0.43
1:A:146:ILE:HB	1:A:161:LYS:HA	2.00	0.43
1:A:189:ARG:NH1	5:A:2143:HOH:O	2.26	0.43
1:A:140:PHE:HZ	5:A:2043:HOH:O	2.01	0.42
1:A:54:HIS:CE1	1:A:95:VAL:HG22	2.54	0.42
1:A:189:ARG:HH11	1:A:189:ARG:HD3	1.67	0.42
1:A:349:HIS:CD2	5:A:2241:HOH:O	2.69	0.42
1:A:380:ILE:HG23	1:A:382:TYR:CE2	2.55	0.42
1:A:393:GLN:HA	1:A:394:PRO:HD2	1.93	0.41
1:A:212:LYS:HB2	1:A:261:GLN:HB2	2.03	0.41
1:A:396:PHE:HB3	5:A:2284:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	476/484 (98%)	455 (96%)	20 (4%)	1 (0%)	43 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	359	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	410/414 (99%)	390 (95%)	20 (5%)	22 21

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	VAL
1	A	72	LYS
1	A	117	THR
1	A	135	ARG
1	A	147	VAL
1	A	156	ARG
1	A	172	LEU
1	A	201	PHE
1	A	244	LEU
1	A	255	ARG
1	A	266	LYS
1	A	271	VAL
1	A	278	LEU
1	A	286	ASP
1	A	297	LEU
1	A	330	LEU
1	A	389	VAL
1	A	429	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	450	LYS
1	A	466	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	121	ASN
1	A	144	ASN
1	A	145	HIS
1	A	174	GLN
1	A	300	ASN
1	A	310	GLN
1	A	317	ASN
1	A	341	HIS
1	A	454	GLN
1	A	464	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OMT	A	53	1	8,9,10	2.51	2 (25%)	6,12,14	1.76	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OMT	A	53	1	-	6/7/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	53	OMT	CG-SD	-5.11	1.71	1.78
1	A	53	OMT	CE-SD	-4.44	1.61	1.75

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	OMT	OD2-SD-OD1	-2.89	111.79	117.22

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	53	OMT	C-CA-CB-CG
1	A	53	OMT	O-C-CA-CB
1	A	53	OMT	CB-CG-SD-OD1
1	A	53	OMT	CB-CG-SD-OD2
1	A	53	OMT	CB-CG-SD-CE
1	A	53	OMT	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	53	OMT	7	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	600	5,1	50,50,50	1.56	10 (20%)	67,82,82	0.83	3 (4%)
4	SO4	A	800	-	4,4,4	0.85	0	6,6,6	0.41	0
3	ACT	A	700	-	3,3,3	1.29	0	3,3,3	1.64	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	600	5,1	-	5/14/54/54	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	HEM	CBC-CAC	4.94	1.54	1.30
2	A	600	HEM	CBB-CAB	4.75	1.53	1.30
2	A	600	HEM	FE-NA	2.68	2.04	1.95
2	A	600	HEM	CAC-C3C	2.51	1.54	1.47
2	A	600	HEM	O2A-CGA	-2.34	1.23	1.30
2	A	600	HEM	FE-NC	2.17	2.02	1.95
2	A	600	HEM	CAB-C3B	2.10	1.53	1.47
2	A	600	HEM	FE-NB	2.09	2.01	1.94
2	A	600	HEM	CMC-C2C	2.06	1.55	1.50
2	A	600	HEM	O1A-CGA	2.05	1.28	1.22

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	HEM	CBB-CAB-C3B	-2.72	113.91	127.53
2	A	600	HEM	CBC-CAC-C3C	-2.55	114.80	127.53
2	A	600	HEM	O1A-CGA-CBA	-2.46	115.30	123.09
3	A	700	ACT	O-C-CH3	-2.24	113.33	122.53

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	600	HEM	CAA-CBA-CGA-O2A
2	A	600	HEM	CAD-CBD-CGD-O2D
2	A	600	HEM	CAD-CBD-CGD-O1D
2	A	600	HEM	CAA-CBA-CGA-O1A
2	A	600	HEM	C2B-C3B-CAB-CBB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	475/484 (98%)	0.45	25 (5%) 32 35	24, 49, 65, 83	3 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	362	HIS	6.3
1	A	361	PHE	5.9
1	A	360	PRO	4.5
1	A	479	GLU	4.3
1	A	359	CYS	4.1
1	A	475	ILE	3.6
1	A	227	ASP	3.1
1	A	466	GLU	3.1
1	A	431	ASP	2.9
1	A	81	GLU	2.8
1	A	404	GLU	2.7
1	A	266	LYS	2.7
1	A	477	VAL	2.7
1	A	250	ARG	2.6
1	A	395	ASP	2.6
1	A	478	LEU	2.4
1	A	397	LYS	2.4
1	A	99[A]	ARG	2.3
1	A	228	GLU	2.2
1	A	284	HIS	2.2
1	A	474	ALA	2.1
1	A	363	ASN	2.1
1	A	205	ASP	2.1
1	A	286	ASP	2.1
1	A	271	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	OMT	A	53	10/11	0.93	0.14	43,45,51,51	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	A	700	4/4	0.77	0.27	68,69,69,69	0
4	SO4	A	800	5/5	0.89	0.12	88,88,88,88	0
2	HEM	A	600	43/43	0.94	0.09	40,42,43,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.