



Full wwPDB NMR Structure Validation Report ⓘ

Feb 14, 2022 – 12:05 AM EST

PDB ID : 1I9K
Title : THE RNA I-MOTIF
Authors : Snoussi, K.; Nonin-Lecomte, S.; Lerou, J.L.
Deposited on : 2001-03-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26

2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 736 atoms, of which 268 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called 5'-R(*UP*CP*CP*CP*CP*C)-3'.


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	6	184	54	67	17	41	5	0
1	B	6	184	54	67	17	41	5	0
1	C	6	184	54	67	17	41	5	0
1	D	6	184	54	67	17	41	5	0

4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: 5'-R(*UP*CP*CP*CP*CP*C)-3'

Chain A:  100%




- Molecule 1: 5'-R(*UP*CP*CP*CP*CP*C)-3'

Chain B:  100%




- Molecule 1: 5'-R(*UP*CP*CP*CP*CP*C)-3'

Chain C:  100%



- Molecule 1: 5'-R(*UP*CP*CP*CP*CP*C)-3'

Chain D:  100%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *NMR-based simulated annealing*.

Of the 50 calculated structures, 1 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.851

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	0	0	0	0
1	B	0	0	0	0
1	C	0	0	0	0
1	D	0	0	0	0
All	All	0	0	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	6/6 (100%)	5 (83%)	3 (50%)	0.00
1	B	6/6 (100%)	5 (83%)	3 (50%)	0.00
1	C	6/6 (100%)	4 (67%)	2 (33%)	0.00
1	D	6/6 (100%)	4 (67%)	3 (50%)	0.00
All	All	24/24 (100%)	18 (75%)	11 (46%)	0.00

The overall RNA backbone suiteness is 0.00.

All RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	2	C
1	A	3	C
1	A	4	C
1	A	5	C
1	A	6	C
1	B	2	C
1	B	3	C
1	B	4	C
1	B	5	C
1	B	6	C
1	C	2	C
1	C	3	C
1	C	4	C
1	C	5	C
1	D	2	C
1	D	3	C
1	D	4	C
1	D	5	C

All RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	1	U
1	A	2	C
1	A	4	C
1	B	1	U
1	B	2	C
1	B	4	C
1	C	1	U
1	C	2	C
1	D	1	U
1	D	2	C
1	D	4	C

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided