



# Full wwPDB NMR Structure Validation Report ⓘ

Mar 7, 2026 – 04:14 AM UTC

PDB ID : 1IDG / pdb\_00001idg  
Title : THE NMR SOLUTION STRUCTURE OF THE COMPLEX FORMED BETWEEN ALPHA-BUNGAROTOXIN AND AN 18MER COGNATE PEPTIDE  
Authors : Zeng, H.; Moise, L.; Grant, M.A.; Hawrot, E.  
Deposited on : 2001-04-04

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	74	
2	B	19	

## 2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1394 atoms, of which 678 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called ALPHA-BUNGAROTOXIN.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	74	1085	338	534	97	105	11	0

- Molecule 2 is a protein called ACETYLCHOLINE RECEPTOR PROTEIN, ALPHA CHAIN.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
2	B	18	309	111	144	26	26	2	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	199	HSL	-	cloning artifact	UNP P02710


## 4 Residue-property plots [i](#)

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ALPHA-BUNGAROTOXIN



- Molecule 2: ACETYLCHOLINE RECEPTOR PROTEIN, ALPHA CHAIN



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry and simulated annealing*.

Of the 1 calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNSsolve	refinement	1.0

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	2.01	12/565 ( 2.1%)	3.14	32/767 ( 4.2%)
2	B	1.61	0/175 ( 0.0%)	2.62	18/241 ( 7.5%)
All	All	1.93	12/740 ( 1.6%)	3.02	50/1008 ( 5.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	2	25
2	B	1	10
All	All	3	35

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	72	ARG	CA-C	14.00	1.67	1.52
1	A	31	ALA	CA-C	7.51	1.62	1.52
1	A	20	GLU	CA-C	7.14	1.68	1.52
1	A	73	PRO	CA-C	6.54	1.61	1.52
1	A	72	ARG	C-N	6.45	1.49	1.33
1	A	32	PHE	CA-C	6.21	1.61	1.52
1	A	17	PRO	CA-C	6.15	1.57	1.52
1	A	52	LYS	CA-C	5.31	1.59	1.52
1	A	68	HIS	CA-C	5.29	1.59	1.52
1	A	4	HIS	CG-CD2	5.21	1.41	1.35
1	A	68	HIS	ND1-CE1	5.13	1.37	1.32
1	A	68	HIS	CG-CD2	5.08	1.41	1.35

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	72	ARG	N-CA-C	-31.34	49.92	108.85
1	A	73	PRO	N-CA-C	-29.60	51.50	112.47
1	A	32	PHE	N-CA-C	-26.98	53.34	110.80
1	A	20	GLU	N-CA-C	-22.55	47.85	111.00
1	A	73	PRO	N-CA-CB	17.16	121.27	103.25
1	A	73	PRO	CA-N-CD	-16.73	88.58	112.00
1	A	32	PHE	CA-C-O	-11.31	104.33	120.51
2	B	186	HIS	ND1-CE1-NE2	11.18	119.58	108.40
1	A	68	HIS	ND1-CE1-NE2	11.10	119.50	108.40
1	A	4	HIS	ND1-CE1-NE2	10.94	119.34	108.40
1	A	63	ASP	CA-CB-CG	10.74	123.34	112.60
1	A	17	PRO	N-CA-C	10.10	123.02	110.70
1	A	73	PRO	N-CD-CG	9.55	117.53	103.20
1	A	4	HIS	CA-CB-CG	9.32	123.12	113.80
1	A	68	HIS	CE1-NE2-CD2	-8.99	100.01	109.00
1	A	72	ARG	N-CA-CB	8.97	127.14	110.27
2	B	186	HIS	CE1-NE2-CD2	-8.95	100.05	109.00
1	A	4	HIS	CE1-NE2-CD2	-8.82	100.18	109.00
1	A	31	ALA	CA-C-O	-8.75	108.00	120.51
1	A	73	PRO	O-C-N	-8.46	111.22	122.64
2	B	184	TRP	N-CA-CB	8.38	124.74	110.50
1	A	32	PHE	CB-CA-C	8.29	126.91	110.42
1	A	72	ARG	CB-CA-C	8.18	121.09	110.13
2	B	187	TRP	O-C-N	-7.65	112.42	122.59
2	B	196	THR	CA-CB-CG2	7.37	123.03	110.50
1	A	32	PHE	O-C-N	-7.28	112.91	122.59
2	B	188	VAL	O-C-N	-7.13	111.59	123.00
1	A	71	GLN	N-CA-C	6.84	125.38	110.80
2	B	188	VAL	CA-CB-CG2	6.83	122.00	110.40
2	B	188	VAL	CB-CA-C	6.78	124.28	111.40
2	B	187	TRP	CB-CA-C	6.67	123.69	110.42
2	B	186	HIS	CB-CA-C	6.13	121.75	110.10
2	B	189	TYR	N-CA-CB	6.12	120.90	110.50
2	B	190	TYR	CA-CB-CG	6.03	124.75	113.90
1	A	31	ALA	CA-C-N	5.97	132.94	121.54
1	A	31	ALA	C-N-CA	5.97	132.94	121.54
1	A	31	ALA	O-C-N	-5.72	114.99	122.59
2	B	195	ASP	O-C-N	-5.58	114.43	122.36
2	B	186	HIS	CG-ND1-CE1	-5.57	99.83	109.30
1	A	24	TYR	N-CA-CB	-5.55	101.11	110.49
1	A	68	HIS	CG-ND1-CE1	-5.48	99.98	109.30
1	A	73	PRO	CA-C-O	-5.38	110.82	120.60
2	B	186	HIS	CB-CG-CD2	-5.35	124.25	131.20

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	GLY	O-C-N	-5.34	117.80	122.88
2	B	183	GLY	O-C-N	-5.28	115.83	122.70
2	B	185	LYS	O-C-N	-5.25	115.60	122.59
1	A	4	HIS	CG-ND1-CE1	-5.15	100.55	109.30
2	B	190	TYR	N-CA-CB	5.12	119.12	110.77
1	A	68	HIS	CA-CB-CG	5.08	118.88	113.80
1	A	68	HIS	O-C-N	-5.00	115.57	121.32

All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	6	THR	CB
1	A	11	ILE	CB
2	B	196	THR	CB

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	5	THR	Peptide
1	A	6	THR	Peptide
1	A	8	THR	Peptide
1	A	13	ALA	Peptide
1	A	16	CYS	Peptide
1	A	18	PRO	Peptide
1	A	20	GLU	Mainchain,Peptide
1	A	21	ASN	Peptide
1	A	24	TYR	Sidechain
1	A	31	ALA	Mainchain,Peptide
1	A	32	PHE	Mainchain,Peptide
1	A	36	ARG	Sidechain
1	A	54	TYR	Sidechain
1	A	55	GLU	Peptide
1	A	63	ASP	Peptide
1	A	68	HIS	Peptide
1	A	70	LYS	Peptide
1	A	72	ARG	Mainchain,Peptide,Sidechain
1	A	73	PRO	Mainchain,Peptide
2	B	181	TYR	Peptide,Sidechain
2	B	184	TRP	Peptide
2	B	186	HIS	Peptide
2	B	189	TYR	Sidechain
2	B	190	TYR	Peptide,Sidechain

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Group
2	B	194	PRO	Peptide
2	B	197	PRO	Peptide
2	B	198	TYR	Sidechain

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	551	534	532	1
2	B	165	144	140	2
All	All	716	678	672	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
2:B:183:GLY:H	2:B:184:TRP:HB3	0.46	1.70
2:B:192:CYS:C	2:B:193:CYS:SG	0.45	3.00
1:A:23:CYS:SG	1:A:42:LEU:HD13	0.41	2.56

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	72/74 (97%)	30 (42%)	16 (22%)	26 (36%)	0	0
2	B	16/19 (84%)	3 (19%)	4 (25%)	9 (56%)	0	0
All	All	88/93 (95%)	33 (38%)	20 (23%)	35 (40%)	0	0

All 35 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in

the ensemble.

Mol	Chain	Res	Type
1	A	6	THR
1	A	12	SER
1	A	14	VAL
1	A	15	THR
1	A	17	PRO
1	A	18	PRO
1	A	20	GLU
1	A	21	ASN
1	A	24	TYR
1	A	31	ALA
1	A	32	PHE
1	A	33	CYS
1	A	41	GLU
1	A	42	LEU
1	A	44	CYS
1	A	48	CYS
1	A	51	LYS
1	A	53	PRO
1	A	56	GLU
1	A	59	CYS
1	A	62	THR
1	A	64	LYS
1	A	67	PRO
1	A	68	HIS
1	A	71	GLN
1	A	73	PRO
2	B	182	ARG
2	B	183	GLY
2	B	184	TRP
2	B	185	LYS
2	B	188	VAL
2	B	189	TYR
2	B	191	THR
2	B	194	PRO
2	B	197	PRO

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/65 (100%)	50 (77%)	15 (23%)	2	28
2	B	17/17 (100%)	9 (53%)	8 (47%)	0	2
All	All	82/82 (100%)	59 (72%)	23 (28%)	1	20

All 23 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	3	CYS
1	A	12	SER
1	A	29	CYS
1	A	36	ARG
1	A	38	LYS
1	A	41	GLU
1	A	42	LEU
1	A	44	CYS
1	A	54	TYR
1	A	56	GLU
1	A	57	VAL
1	A	59	CYS
1	A	60	CYS
1	A	64	LYS
1	A	70	LYS
2	B	182	ARG
2	B	184	TRP
2	B	185	LYS
2	B	189	TYR
2	B	190	TYR
2	B	191	THR
2	B	193	CYS
2	B	196	THR

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided