



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 12:56 PM EDT

PDB ID : 1JDC
Title : MUTANT (E219Q) MALTOTETRAOSE-FORMING EXO-AMYLASE
COCRYSTALLIZED WITH MALTOTETRAOSE (CRYSTAL TYPE 1)
Authors : Yoshioka, Y.; Hasegawa, K.; Matsuura, Y.; Katsube, Y.; Kubota, M.
Deposited on : 1997-06-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	
2	B	4	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3556 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1,4-ALPHA MALTOTETRAHYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	418	3297	2070	597	620	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	219	GLN	GLU	engineered mutation	UNP P13507
A	334	ASP	SER	conflict	UNP P13507

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	4	45	24	21	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	2	2	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	212	212	212	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1,4-ALPHA MALTOTETRAHYDROLASE



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.10Å 138.40Å 50.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.90 26.17 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.90) 75.1 (26.17-1.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.89Å)	Xtrriage
Refinement program	PROFFT, X-PLOR	Depositor
R, R_{free}	0.174 , (Not available) 0.165 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtrriage
Anisotropy	0.318	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 58.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3556	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	1/3402 (0.0%)	1.80	51/4631 (1.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	ARG	NE-CZ	-5.81	1.25	1.33

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	202	ARG	CD-NE-CZ	40.25	179.94	123.60
1	A	346	ARG	CD-NE-CZ	26.81	161.13	123.60
1	A	61	ARG	CD-NE-CZ	22.52	155.13	123.60
1	A	61	ARG	NE-CZ-NH2	21.60	131.10	120.30
1	A	175	ARG	NE-CZ-NH1	19.77	130.18	120.30
1	A	175	ARG	CD-NE-CZ	16.24	146.34	123.60
1	A	202	ARG	NE-CZ-NH2	16.14	128.37	120.30
1	A	417	ARG	NE-CZ-NH1	15.93	128.27	120.30
1	A	248	ARG	NE-CZ-NH1	15.44	128.02	120.30
1	A	346	ARG	NE-CZ-NH1	14.97	127.78	120.30
1	A	61	ARG	NE-CZ-NH1	-11.51	114.55	120.30
1	A	358	ARG	NE-CZ-NH1	10.83	125.71	120.30
1	A	417	ARG	CD-NE-CZ	10.14	137.79	123.60
1	A	137	ARG	NE-CZ-NH1	10.13	125.36	120.30
1	A	155	ARG	NE-CZ-NH2	-9.70	115.45	120.30
1	A	11	ARG	NE-CZ-NH1	9.38	124.99	120.30
1	A	196	ARG	NE-CZ-NH1	9.24	124.92	120.30
1	A	316	ARG	NE-CZ-NH1	-8.63	115.98	120.30
1	A	390	ASP	CB-CG-OD1	8.56	126.00	118.30
1	A	196	ARG	NE-CZ-NH2	-8.49	116.06	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	417	ARG	NE-CZ-NH2	-8.35	116.12	120.30
1	A	248	ARG	CD-NE-CZ	8.07	134.90	123.60
1	A	346	ARG	NE-CZ-NH2	-8.06	116.27	120.30
1	A	358	ARG	CD-NE-CZ	7.57	134.19	123.60
1	A	358	ARG	NE-CZ-NH2	-7.25	116.67	120.30
1	A	316	ARG	CD-NE-CZ	-7.18	113.55	123.60
1	A	212	ASP	CB-CG-OD2	-6.75	112.23	118.30
1	A	261	ARG	NE-CZ-NH1	6.62	123.61	120.30
1	A	233	ARG	NE-CZ-NH1	6.62	123.61	120.30
1	A	152	ASP	CB-CG-OD1	6.57	124.22	118.30
1	A	61	ARG	CA-CB-CG	6.30	127.25	113.40
1	A	175	ARG	NH1-CZ-NH2	-6.19	112.59	119.40
1	A	175	ARG	NE-CZ-NH2	-6.17	117.21	120.30
1	A	283	ARG	CB-CA-C	-5.95	98.50	110.40
1	A	281	ARG	NE-CZ-NH2	5.86	123.23	120.30
1	A	191	ARG	NE-CZ-NH1	5.83	123.21	120.30
1	A	202	ARG	NH1-CZ-NH2	-5.74	113.09	119.40
1	A	137	ARG	NE-CZ-NH2	-5.69	117.46	120.30
1	A	255	ASP	CB-CG-OD2	5.67	123.41	118.30
1	A	171	TYR	CB-CG-CD2	-5.59	117.64	121.00
1	A	332	TYR	CB-CG-CD1	-5.53	117.68	121.00
1	A	124	ASP	CB-CG-OD2	5.44	123.20	118.30
1	A	373	VAL	CB-CA-C	5.34	121.56	111.40
1	A	195	VAL	CB-CA-C	5.30	121.47	111.40
1	A	152	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	A	244	ASP	CB-CG-OD1	-5.27	113.56	118.30
1	A	248	ARG	NH1-CZ-NH2	-5.16	113.73	119.40
1	A	244	ASP	CB-CG-OD2	5.14	122.93	118.30
1	A	40	ARG	CD-NE-CZ	-5.12	116.43	123.60
1	A	352	ARG	NE-CZ-NH1	5.07	122.83	120.30
1	A	279	ASP	CB-CG-OD1	5.03	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3297	0	3015	80	0
2	B	45	0	39	4	0
3	A	2	0	0	0	0
4	A	212	0	0	11	0
All	All	3556	0	3054	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:PRO:HD3	1:A:334:ASP:OD2	1.47	1.14
1:A:317:GLN:H	1:A:317:GLN:HE21	1.01	0.94
1:A:61:ARG:HG3	1:A:84:ASN:O	1.67	0.94
1:A:167:HIS:HD2	1:A:169:GLN:H	1.19	0.86
1:A:409:SER:HB3	1:A:412:GLN:HB2	1.61	0.82
1:A:277:ASN:HD22	1:A:279:ASP:H	1.28	0.80
1:A:145:ASN:ND2	1:A:155:ARG:H	1.81	0.79
1:A:316:ARG:NH1	1:A:316:ARG:HG2	1.98	0.77
1:A:167:HIS:CD2	1:A:169:GLN:H	2.04	0.75
1:A:317:GLN:H	1:A:317:GLN:NE2	1.83	0.74
1:A:193:ASP:OD1	2:B:1:GLC:H1	1.88	0.74
1:A:317:GLN:HE21	1:A:317:GLN:N	1.81	0.72
1:A:358:ARG:HD3	4:A:607:HOH:O	1.89	0.71
1:A:409:SER:O	1:A:410:ASN:HB2	1.90	0.71
1:A:145:ASN:ND2	1:A:155:ARG:HG2	2.07	0.69
1:A:145:ASN:HD21	1:A:155:ARG:H	1.40	0.69
1:A:316:ARG:CG	1:A:316:ARG:HH11	2.07	0.68
1:A:301:GLN:NE2	4:A:703:HOH:O	2.22	0.67
1:A:316:ARG:NH1	1:A:316:ARG:CG	2.56	0.67
1:A:325:SER:HB2	1:A:326:PRO:CD	2.25	0.65
1:A:143:PRO:HD2	4:A:771:HOH:O	1.96	0.64
1:A:334:ASP:OD1	1:A:338:ASP:OD2	2.16	0.63
1:A:216:CYS:O	1:A:252:PRO:HD2	1.98	0.63
1:A:347:GLN:OE1	1:A:406:VAL:HG22	1.98	0.63
1:A:145:ASN:HD22	1:A:155:ARG:HG2	1.64	0.62
1:A:167:HIS:HD2	1:A:169:GLN:N	1.94	0.62
1:A:407:ASN:C	1:A:407:ASN:HD22	2.05	0.61
1:A:24:HIS:HD2	1:A:26:ASN:H	1.47	0.60
1:A:325:SER:HB2	1:A:326:PRO:HD2	1.82	0.60
1:A:122:TYR:O	1:A:125:LYS:NZ	2.35	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:ARG:HH12	1:A:204:ASN:HD22	1.52	0.58
1:A:366:HIS:HE1	1:A:375:THR:OG1	1.88	0.56
1:A:417:ARG:HH11	1:A:417:ARG:HG3	1.70	0.56
1:A:145:ASN:HD21	1:A:154:ASP:HB3	1.73	0.54
1:A:299:PRO:CD	1:A:334:ASP:OD2	2.38	0.54
1:A:292:ASN:ND2	1:A:295:THR:H	2.06	0.53
1:A:237:SER:H	1:A:240:GLN:HE21	1.56	0.53
1:A:157:ILE:HG21	2:B:4:GLC:H3	1.90	0.53
1:A:24:HIS:HE1	4:A:658:HOH:O	1.91	0.53
1:A:404:GLU:HG2	4:A:634:HOH:O	2.10	0.52
1:A:61:ARG:HD2	1:A:82:ASP:OD1	2.09	0.52
1:A:130:PRO:HG2	1:A:133:GLN:NE2	2.25	0.51
1:A:1:ASP:HA	1:A:16:ASP:OD2	2.10	0.51
1:A:42:GLN:O	1:A:45:THR:HG22	2.11	0.51
1:A:170:VAL:HA	1:A:173:MET:HE3	1.93	0.49
1:A:404:GLU:CG	4:A:634:HOH:O	2.60	0.49
1:A:2:GLN:H	1:A:2:GLN:HE21	1.60	0.49
1:A:358:ARG:HE	1:A:358:ARG:HB3	1.51	0.48
1:A:366:HIS:HB2	1:A:373:VAL:HG22	1.95	0.48
1:A:26:ASN:HD22	1:A:29:ARG:HE	1.62	0.47
1:A:237:SER:H	1:A:240:GLN:NE2	2.13	0.47
1:A:319:TYR:OH	1:A:335:HIS:HD2	1.97	0.47
1:A:366:HIS:CE1	1:A:375:THR:OG1	2.66	0.47
1:A:62:ASP:O	1:A:75:GLY:HA2	2.14	0.46
1:A:17:GLU:OE2	1:A:108:LYS:HE3	2.15	0.46
1:A:116:ASN:OD1	1:A:197:GLY:HA3	2.17	0.45
1:A:292:ASN:ND2	1:A:294:ASP:H	2.13	0.45
1:A:40:ARG:HD2	1:A:97:GLN:HB3	1.98	0.45
1:A:340:GLY:HA2	4:A:556:HOH:O	2.16	0.45
1:A:199:ALA:HA	1:A:200:PRO:HD3	1.83	0.45
1:A:155:ARG:HA	1:A:162:ASP:OD2	2.16	0.45
1:A:353:ARG:NH2	4:A:554:HOH:O	2.49	0.45
1:A:319:TYR:O	1:A:323:LEU:HB2	2.16	0.45
1:A:334:ASP:HB3	1:A:339:TRP:CE3	2.52	0.44
1:A:40:ARG:NH2	4:A:732:HOH:O	2.51	0.44
1:A:394:PRO:HD3	1:A:416:TRP:CE2	2.53	0.44
1:A:71:LYS:HE2	4:A:680:HOH:O	2.18	0.43
1:A:319:TYR:OH	1:A:335:HIS:CD2	2.71	0.43
1:A:31:ALA:N	1:A:32:PRO:CD	2.82	0.43
1:A:61:ARG:HD3	1:A:81:HIS:CE1	2.54	0.42
1:A:207:MET:HE3	1:A:211:ALA:O	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:TYR:HA	1:A:123:PRO:HD3	1.86	0.42
1:A:157:ILE:CD1	2:B:3:GLC:H3	2.49	0.42
1:A:142:ASP:HA	1:A:143:PRO:HD2	1.94	0.42
1:A:277:ASN:O	1:A:283:ARG:CG	2.68	0.42
1:A:218:GLY:HA3	1:A:245:TRP:CH2	2.55	0.41
1:A:409:SER:O	1:A:410:ASN:CB	2.64	0.41
1:A:182:ARG:HA	1:A:187:ALA:HB3	2.02	0.41
1:A:193:ASP:OD1	2:B:1:GLC:C1	2.62	0.41
1:A:196:ARG:HG2	4:A:509:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	416/429 (97%)	401 (96%)	14 (3%)	1 (0%)	47 38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	157	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	333/337 (99%)	307 (92%)	26 (8%)	12 5

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	12	TYR
1	A	61	ARG
1	A	79	PHE
1	A	85	LYS
1	A	95	LEU
1	A	102	LEU
1	A	119	ASN
1	A	162	ASP
1	A	181	LEU
1	A	195	VAL
1	A	196	ARG
1	A	202	ARG
1	A	229	ASN
1	A	277	ASN
1	A	292	ASN
1	A	316	ARG
1	A	317	GLN
1	A	323	LEU
1	A	353	ARG
1	A	364	SER
1	A	373	VAL
1	A	397	VAL
1	A	407	ASN
1	A	409	SER
1	A	417	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	8	ASN
1	A	24	HIS
1	A	26	ASN
1	A	81	HIS
1	A	119	ASN
1	A	128	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	133	GLN
1	A	145	ASN
1	A	167	HIS
1	A	180	ASN
1	A	204	ASN
1	A	240	GLN
1	A	264	ASN
1	A	277	ASN
1	A	292	ASN
1	A	317	GLN
1	A	335	HIS
1	A	366	HIS
1	A	388	ASN
1	A	393	ASN
1	A	407	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	12,12,12	1.04	1 (8%)	17,17,17	2.43	4 (23%)
2	GLC	B	2	2	11,11,12	1.02	2 (18%)	15,15,17	1.94	4 (26%)
2	GLC	B	3	2	11,11,12	0.56	0	15,15,17	1.28	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	4	2	11,11,12	0.59	0	15,15,17	1.60	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	2/2/19/22	0/1/1/1
2	GLC	B	4	2	-	2/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	GLC	O4-C4	-2.40	1.37	1.43
2	B	2	GLC	O4-C4	-2.22	1.37	1.43
2	B	2	GLC	C1-C2	-2.02	1.47	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C1-O5-C5	6.71	126.33	113.66
2	B	1	GLC	O5-C1-C2	4.83	118.91	110.28
2	B	2	GLC	C1-O5-C5	4.49	118.28	112.19
2	B	4	GLC	C1-C2-C3	3.88	114.44	109.67
2	B	4	GLC	O4-C4-C3	3.79	119.11	110.35
2	B	3	GLC	C1-O5-C5	3.77	117.29	112.19
2	B	1	GLC	O2-C2-C1	3.40	117.06	109.16
2	B	2	GLC	O5-C5-C6	2.89	111.74	107.20
2	B	2	GLC	O3-C3-C2	-2.29	105.62	109.99
2	B	1	GLC	O1-C1-C2	2.06	114.84	109.03
2	B	2	GLC	C1-C2-C3	2.04	112.17	109.67

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	3	GLC	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	3	GLC	C4-C5-C6-O6
2	B	4	GLC	C4-C5-C6-O6
2	B	4	GLC	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	GLC	1	0
2	B	4	GLC	1	0
2	B	1	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	418/429 (97%)	-0.39	4 (0%) 82 84	9, 16, 31, 52	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ASP	6.2
1	A	2	GLN	4.8
1	A	146	TYR	4.8
1	A	3	ALA	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	4	11/12	0.96	0.08	16,17,18,19	0
2	GLC	B	3	11/12	0.97	0.07	14,15,17,17	0
2	GLC	B	1	12/12	0.98	0.09	10,12,12,13	0
2	GLC	B	2	11/12	0.98	0.05	12,12,13,14	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	A	452	1/1	0.96	0.19	59,59,59,59	0
3	CA	A	451	1/1	1.00	0.07	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.