



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 03:11 AM EDT

PDB ID : 1JY1
Title : CRYSTAL STRUCTURE OF HUMAN TYROSYL-DNA PHOSPHODI-
ESTERASE (TDP1)
Authors : Davies, D.R.; Interthal, H.; Champoux, J.J.; Hol, W.G.J.
Deposited on : 2001-09-10
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TYROSYL-DNA PHOSPHODIESTERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	439	3505	2273	590	631	3	8	0	0	0

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	145	LEU	-	expression tag	UNP Q9NUW8
A	146	GLU	-	expression tag	UNP Q9NUW8
A	147	ASP	-	expression tag	UNP Q9NUW8
A	148	PRO	-	expression tag	UNP Q9NUW8
A	157	MSE	MET	modified residue	UNP Q9NUW8
A	266	MSE	MET	modified residue	UNP Q9NUW8
A	267	MSE	MET	modified residue	UNP Q9NUW8
A	322	ASN	ASP	engineered mutation	UNP Q9NUW8
A	328	THR	MET	engineered mutation	UNP Q9NUW8
A	386	MSE	MET	modified residue	UNP Q9NUW8
A	420	MSE	MET	modified residue	UNP Q9NUW8
A	491	MSE	MET	modified residue	UNP Q9NUW8
A	498	MSE	MET	modified residue	UNP Q9NUW8
A	533	MSE	MET	modified residue	UNP Q9NUW8
A	548	LEU	PHE	engineered mutation	UNP Q9NUW8
A	567	MSE	MET	modified residue	UNP Q9NUW8
A	604	MSE	MET	modified residue	UNP Q9NUW8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	345	Total	O	0	0
			345	345		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TYROSYL-DNA PHOSPHODIESTERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	186.44Å 52.03Å 56.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.69 19.95 – 1.65	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.69) 92.8 (19.95-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.29 (at 1.65Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.198 , 0.229 0.197 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.234	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3850	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	1/3610 (0.0%)	0.95	7/4896 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	498	MSE	SE-CE	-5.48	1.63	1.95

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	ASP	CB-CG-OD2	8.28	125.75	118.30
1	A	407	ASP	CB-CG-OD2	7.91	125.42	118.30
1	A	159	ASP	CB-CG-OD2	7.64	125.17	118.30
1	A	551	ASP	CB-CG-OD2	6.51	124.16	118.30
1	A	147	ASP	CB-CG-OD2	5.73	123.46	118.30
1	A	479	LYS	CD-CE-NZ	-5.09	99.99	111.70
1	A	599	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3505	0	3424	48	1
2	A	345	0	0	2	1
All	All	3850	0	3424	48	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:VAL:HG11	1:A:420:MSE:CE	1.72	1.18
1:A:394:VAL:HG11	1:A:420:MSE:HE3	1.11	1.10
1:A:607:PRO:O	1:A:608:SER:HB2	1.52	1.10
1:A:394:VAL:CG1	1:A:420:MSE:HE3	1.92	0.97
1:A:394:VAL:CG1	1:A:420:MSE:CE	2.44	0.96
1:A:389:ALA:CB	1:A:423:LEU:CD1	2.53	0.86
1:A:291:GLN:HB2	1:A:452:GLU:OE2	1.77	0.85
1:A:228:HIS:HD2	1:A:230:ASP:H	1.30	0.79
1:A:498:MSE:HE1	1:A:507:ILE:HG21	1.68	0.76
1:A:389:ALA:CB	1:A:423:LEU:HD12	2.16	0.75
1:A:394:VAL:CG1	1:A:420:MSE:HE1	2.17	0.74
1:A:389:ALA:HB3	1:A:423:LEU:CD1	2.23	0.68
1:A:607:PRO:O	1:A:608:SER:CB	2.39	0.66
1:A:421:LEU:HD12	1:A:421:LEU:C	2.16	0.66
1:A:389:ALA:HB1	1:A:423:LEU:HD12	1.80	0.64
1:A:498:MSE:CE	1:A:507:ILE:HG21	2.28	0.63
1:A:389:ALA:HB1	1:A:423:LEU:CD1	2.29	0.62
1:A:389:ALA:HB3	1:A:423:LEU:HD12	1.82	0.61
1:A:389:ALA:CB	1:A:423:LEU:HD13	2.33	0.56
1:A:218:GLU:OE1	1:A:218:GLU:N	2.29	0.55
1:A:228:HIS:HD2	1:A:230:ASP:N	2.01	0.54
1:A:389:ALA:HB3	1:A:423:LEU:HD13	1.92	0.52
1:A:498:MSE:HE1	1:A:507:ILE:CG2	2.39	0.51
1:A:310:HIS:O	1:A:310:HIS:ND1	2.37	0.50
1:A:555:VAL:O	1:A:556:LYS:CB	2.59	0.50
1:A:319:PHE:CG	1:A:350:THR:HG21	2.48	0.48
1:A:457:GLY:HA3	1:A:588:TRP:CZ2	2.49	0.48
1:A:344:LYS:HD3	1:A:344:LYS:HA	1.61	0.48
1:A:304:ARG:NH2	2:A:1047:HOH:O	2.48	0.47
1:A:183:LEU:CD1	1:A:298:LEU:HD11	2.45	0.46
1:A:383:ALA:HB1	1:A:423:LEU:HD22	1.98	0.46
1:A:395:VAL:HB	1:A:498:MSE:HG3	1.99	0.45
1:A:452:GLU:HB2	1:A:456:ALA:HB2	1.97	0.45
1:A:498:MSE:CE	1:A:507:ILE:CG2	2.95	0.45
1:A:176:PRO:HG2	2:A:994:HOH:O	2.17	0.45
1:A:421:LEU:C	1:A:421:LEU:CD1	2.85	0.44
1:A:302:TYR:CD1	1:A:347:LEU:HA	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:228:HIS:CD2	1:A:230:ASP:H	2.20	0.44
1:A:554:LYS:HB2	1:A:569:THR:HG22	2.01	0.43
1:A:454:TYR:CD2	1:A:599:ASP:HB3	2.54	0.42
1:A:392:TRP:CD1	1:A:502:PRO:HD3	2.55	0.42
1:A:398:PHE:CE2	1:A:401:VAL:HG22	2.54	0.41
1:A:203:ASN:OD1	1:A:283:ASN:HA	2.20	0.41
1:A:504:PHE:CE1	1:A:577:PRO:HB3	2.56	0.41
1:A:158:LEU:HD21	1:A:298:LEU:CD1	2.51	0.41
1:A:394:VAL:HG22	1:A:499:ARG:HD2	2.01	0.41
1:A:527:LYS:O	1:A:530:THR:HG23	2.21	0.40
1:A:216:PRO:O	1:A:217:PRO:C	2.59	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:845:HOH:O	2:A:845:HOH:O[2_655]	1.85	0.35
1:A:365:SER:OG	1:A:380:LYS:NZ[2_655]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	431/464 (93%)	422 (98%)	8 (2%)	1 (0%)	47 30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	371	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	378/393 (96%)	368 (97%)	10 (3%)	46 28

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	GLU
1	A	310	HIS
1	A	331	ASN
1	A	337	GLU
1	A	344	LYS
1	A	452	GLU
1	A	584	LYS
1	A	590	TRP
1	A	596	LYS
1	A	608	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	GLN
1	A	228	HIS
1	A	331	ASN
1	A	363	GLN
1	A	397	GLN
1	A	531	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	431/464 (92%)	0.14	23 (5%) 26 29	10, 20, 36, 47	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	389	ALA	7.4
1	A	608	SER	5.9
1	A	308	GLY	4.4
1	A	590	TRP	4.2
1	A	307	ASP	4.0
1	A	310	HIS	3.4
1	A	423	LEU	3.3
1	A	530	THR	3.2
1	A	528	ASN	3.2
1	A	180	SER	3.1
1	A	391	SER	3.0
1	A	148	PRO	3.0
1	A	390	GLU	2.9
1	A	147	ASP	2.6
1	A	145	LEU	2.5
1	A	596	LYS	2.5
1	A	246	GLU	2.4
1	A	309	THR	2.3
1	A	152	GLN	2.3
1	A	584	LYS	2.2
1	A	434	SER	2.1
1	A	556	LYS	2.0
1	A	366	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.