



Full wwPDB NMR Structure Validation Report ⓘ

Mar 5, 2026 – 09:00 PM UTC

PDB ID : 1KA6 / pdb_00001ka6
Title : SAP/SH2D1A bound to peptide n-pY
Authors : Hwang, P.M.; Li, C.; Morra, M.; Lillywhite, J.; Gertler, F.; Terhorst, C.; Kay, L.E.; Pawson, T.; Forman-Kay, J.; Li, S.-C.
Deposited on : 2001-10-31

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	128	 55% 27% 16%
2	B	9	 44% 33% 22%

2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1847 atoms, of which 931 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called SH2 DOMAIN PROTEIN 1A.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	107	1699	543	854	142	157	3	0

- Molecule 2 is a protein called peptide n-pY.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	B	9	148	43	77	13	14	1	1

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	275	ARG	LYS	engineered mutation	UNP Q13291
B	281	PTR	TYR	modified residue	UNP Q13291

5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	structure solution	1.0
ARIA	refinement	1.0

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, NH2

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	845	854	851	14
2	B	71	77	71	1
All	All	916	931	922	14

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:20:LEU:HB3	1:A:46:LEU:HD13	0.68	1.66
1:A:28:SER:HA	1:A:100:TYR:O	0.65	1.91
1:A:42:CYS:HB3	1:A:53:THR:HG23	0.60	1.73
1:A:4:VAL:HG11	1:A:31:LEU:HD22	0.56	1.78
1:A:17:GLU:HA	1:A:51:ILE:HG12	0.52	1.80
1:A:8:HIS:HA	1:A:104:LYS:HG3	0.49	1.84
1:A:21:LEU:HD11	2:B:278:LEU:HD22	0.47	1.85
1:A:29:TYR:CD2	1:A:98:LEU:HB3	0.47	2.45
1:A:72:VAL:HG11	1:A:93:GLY:HA3	0.46	1.86
1:A:29:TYR:CE2	1:A:98:LEU:HB3	0.45	2.47
1:A:25:LEU:O	1:A:46:LEU:HD23	0.41	2.14
1:A:44:CYS:HB3	1:A:51:ILE:HG23	0.41	1.92

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:6:VAL:HG12	1:A:102:VAL:C	0.40	2.42
1:A:29:TYR:HA	1:A:44:CYS:O	0.40	2.16

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	105/128 (82%)	95 (90%)	9 (9%)	1 (1%)	15 65
2	B	6/9 (67%)	2 (33%)	3 (50%)	1 (17%)	0 3
All	All	111/137 (81%)	97 (87%)	12 (11%)	2 (2%)	9 52

All 2 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	6	VAL
2	B	276	LYS

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	92/109 (84%)	74 (80%)	18 (20%)	3 33
2	B	6/6 (100%)	2 (33%)	4 (67%)	0 0
All	All	98/115 (85%)	76 (78%)	22 (22%)	2 29

All 22 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	12	SER
1	A	14	GLU
1	A	18	LYS
1	A	23	THR
1	A	30	LEU
1	A	32	ARG
1	A	36	SER
1	A	53	THR
1	A	57	SER
1	A	63	SER
1	A	65	SER
1	A	74	LYS
1	A	78	ARG
1	A	81	LYS
1	A	89	LYS
1	A	104	LYS
1	A	105	LYS
1	A	106	SER
2	B	275	ARG
2	B	276	LYS
2	B	277	SER
2	B	278	LEU

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	PTR	B	281	2	15,16,17	0.80	1 (6%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	PTR	B	281	2	17,22,24	0.53	0 (0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	B	281	2	-	0,10,11,13	0,1,1,1

All bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	281	PTR	P-OH	2.02	1.63	1.59

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided