



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 10:36 AM EDT

PDB ID : 1LZ1
Title : REFINEMENT OF HUMAN LYSOZYME AT 1.5 ANGSTROMS RESOLUTION. ANALYSIS OF NON-BONDED AND HYDROGEN-BOND INTERACTIONS
Authors : Artymiuk, P.J.; Blake, C.C.F.
Deposited on : 1984-10-12
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	130	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1029 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN LYSOZYME.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	130	1029	633	200	186	10	0	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HUMAN LYSOZYME



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.13Å 60.99Å 32.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.50	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	0.177 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1029	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.32	10/1049 (1.0%)	2.53	58/1416 (4.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	112	TRP	NE1-CE2	-7.21	1.28	1.37
1	A	109	TRP	NE1-CE2	-7.04	1.28	1.37
1	A	64	TRP	NE1-CE2	-6.98	1.28	1.37
1	A	34	TRP	NE1-CE2	-5.60	1.30	1.37
1	A	35	GLU	CD-OE2	-5.51	1.19	1.25
1	A	45	TYR	CD1-CE1	5.29	1.47	1.39
1	A	28	TRP	NE1-CE2	-5.20	1.30	1.37
1	A	7	GLU	CD-OE2	-5.19	1.20	1.25
1	A	4	GLU	CD-OE2	-5.03	1.20	1.25
1	A	20	TYR	CD1-CE1	5.02	1.46	1.39

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	113	ARG	NE-CZ-NH2	-24.63	107.99	120.30
1	A	107	ARG	NE-CZ-NH2	19.56	130.08	120.30
1	A	14	ARG	NE-CZ-NH1	18.69	129.64	120.30
1	A	98	ARG	NE-CZ-NH1	-17.84	111.38	120.30
1	A	115	ARG	NE-CZ-NH2	16.34	128.47	120.30
1	A	45	TYR	CB-CG-CD1	-13.02	113.19	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	50	ARG	NE-CZ-NH1	12.24	126.42	120.30
1	A	62	ARG	NE-CZ-NH2	12.19	126.40	120.30
1	A	119	ARG	NE-CZ-NH2	12.14	126.37	120.30
1	A	4	GLU	OE1-CD-OE2	12.02	137.73	123.30
1	A	18	ASP	CB-CG-OD1	10.71	127.94	118.30
1	A	98	ARG	NE-CZ-NH2	10.41	125.50	120.30
1	A	107	ARG	NH1-CZ-NH2	-10.32	108.05	119.40
1	A	18	ASP	CB-CG-OD2	-10.25	109.08	118.30
1	A	20	TYR	CB-CG-CD1	-9.79	115.13	121.00
1	A	45	TYR	CG-CD1-CE1	-9.62	113.60	121.30
1	A	4	GLU	CG-CD-OE1	-8.96	100.38	118.30
1	A	5	ARG	NE-CZ-NH2	-8.84	115.88	120.30
1	A	107	ARG	CD-NE-CZ	8.04	134.86	123.60
1	A	113	ARG	NH1-CZ-NH2	7.91	128.10	119.40
1	A	69	LYS	CA-CB-CG	7.81	130.58	113.40
1	A	124	TYR	CB-CG-CD1	-7.69	116.39	121.00
1	A	45	TYR	CD1-CG-CD2	7.55	126.20	117.90
1	A	124	TYR	CB-CG-CD2	7.49	125.49	121.00
1	A	119	ARG	NE-CZ-NH1	-7.28	116.66	120.30
1	A	113	ARG	NE-CZ-NH1	7.17	123.88	120.30
1	A	34	TRP	CA-CB-CG	-6.86	100.66	113.70
1	A	87	ASP	CB-CG-OD2	-6.70	112.27	118.30
1	A	82	SER	N-CA-CB	-6.69	100.46	110.50
1	A	62	ARG	NH1-CZ-NH2	-6.68	112.05	119.40
1	A	14	ARG	NE-CZ-NH2	-6.66	116.97	120.30
1	A	53	ASP	CB-CG-OD2	-6.51	112.44	118.30
1	A	69	LYS	CB-CG-CD	-6.21	95.46	111.60
1	A	38	TYR	CB-CG-CD1	-6.15	117.31	121.00
1	A	20	TYR	CG-CD1-CE1	-6.06	116.45	121.30
1	A	110	VAL	CG1-CB-CG2	6.01	120.53	110.90
1	A	88	ASN	OD1-CG-ND2	-5.98	108.14	121.90
1	A	7	GLU	CG-CD-OE2	-5.90	106.50	118.30
1	A	33	LYS	CB-CG-CD	5.86	126.85	111.60
1	A	38	TYR	CG-CD1-CE1	-5.71	116.73	121.30
1	A	41	ARG	NE-CZ-NH1	-5.69	117.46	120.30
1	A	88	ASN	CB-CG-OD1	5.66	132.92	121.60
1	A	14	ARG	NH1-CZ-NH2	-5.63	113.21	119.40
1	A	21	ARG	CD-NE-CZ	5.60	131.45	123.60
1	A	115	ARG	NH1-CZ-NH2	-5.60	113.25	119.40
1	A	21	ARG	NE-CZ-NH1	-5.51	117.55	120.30
1	A	72	GLY	CA-C-O	-5.51	110.69	120.60
1	A	54	TYR	CB-CG-CD1	-5.48	117.71	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	7	GLU	OE1-CD-OE2	5.42	129.80	123.30
1	A	5	ARG	NE-CZ-NH1	5.37	122.99	120.30
1	A	114	ASN	CA-CB-CG	-5.36	101.60	113.40
1	A	67	ASP	CB-CG-OD1	-5.27	113.56	118.30
1	A	73	ALA	CA-C-O	5.21	131.05	120.10
1	A	123	GLN	N-CA-CB	-5.20	101.25	110.60
1	A	74	VAL	CB-CA-C	-5.14	101.64	111.40
1	A	101	ARG	NE-CZ-NH1	5.13	122.87	120.30
1	A	104	GLN	CA-CB-CG	5.09	124.59	113.40
1	A	39	ASN	CB-CG-OD1	5.00	131.61	121.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	107	ARG	Sidechain
1	A	113	ARG	Sidechain
1	A	115	ARG	Sidechain
1	A	122	ARG	Sidechain
1	A	14	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1029	0	992	20	1
All	All	1029	0	992	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:ASP:OD1	1:A:122:ARG:HD2	1.47	1.12

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:ARG:NH1	1:A:112:TRP:CH2	2.42	0.87
1:A:107:ARG:HH12	1:A:112:TRP:HH2	1.33	0.76
1:A:120:ASP:OD1	1:A:122:ARG:CD	2.32	0.76
1:A:107:ARG:NH1	1:A:112:TRP:CZ3	2.56	0.74
1:A:107:ARG:NH1	1:A:112:TRP:HH2	1.89	0.70
1:A:111:ALA:O	1:A:115:ARG:HG2	1.90	0.69
1:A:74:VAL:O	1:A:74:VAL:HG12	2.03	0.59
1:A:62:ARG:O	1:A:73:ALA:HA	2.02	0.58
1:A:14:ARG:CZ	1:A:14:ARG:HB3	2.34	0.58
1:A:14:ARG:CZ	1:A:14:ARG:CB	2.82	0.52
1:A:110:VAL:HG12	1:A:114:ASN:ND2	2.26	0.50
1:A:122:ARG:HB3	1:A:126:GLN:HE21	1.77	0.49
1:A:110:VAL:HG12	1:A:114:ASN:HD21	1.76	0.49
1:A:49:ASP:O	1:A:62:ARG:NH2	2.34	0.48
1:A:74:VAL:O	1:A:74:VAL:CG1	2.62	0.46
1:A:107:ARG:HH11	1:A:112:TRP:HZ3	1.65	0.45
1:A:49:ASP:CG	1:A:62:ARG:HH12	2.20	0.44
1:A:25:LEU:HD23	1:A:130:VAL:HG21	2.02	0.42
1:A:84:LEU:HD22	1:A:92:ALA:HA	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ARG:CD	1:A:127:GLY:O[2_565]	2.11	0.09

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	128/130 (98%)	126 (98%)	2 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	105/105 (100%)	99 (94%)	6 (6%)	20 2

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	14	ARG
1	A	69	LYS
1	A	107	ARG
1	A	115	ARG
1	A	122	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	114	ASN
1	A	117	GLN
1	A	126	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.