



Full wwPDB NMR Structure Validation Report ⓘ

Oct 17, 2021 – 12:02 AM EDT

PDB ID : 1RGR
Title : Cyclic Peptides Targeting PDZ Domains of PSD-95: Structural Basis for Enhanced Affinity and Enzymatic Stability
Authors : Piserchio, A.; Salinas, G.D.; Li, T.; Marshall, J.; Spaller, M.R.; Mierke, D.F.
Deposited on : 2003-11-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.23.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	99	58% (green), 12% (yellow), 21% (cyan), 6% (grey)
2	B	6	50% (green), 50% (yellow)

2 Ensemble composition and analysis

This entry contains 22 models. Model 7 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:63-A:69, A:75-A:87, A:92-A:108, A:115-A:149, B:1-B:6 (78)	0.36	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	3, 4, 9, 10, 13, 20, 22
2	5, 11, 15, 16, 21
3	2, 8, 14, 18, 19
4	1, 7
Single-model clusters	6; 12; 17

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1531 atoms, of which 768 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Presynaptic density protein 95.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	93	1411	440	706	126	138	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	155	HIS	-	expression tag	UNP P31016
A	156	HIS	-	expression tag	UNP P31016
A	157	HIS	-	expression tag	UNP P31016
A	158	HIS	-	expression tag	UNP P31016
A	159	HIS	-	expression tag	UNP P31016
A	160	HIS	-	expression tag	UNP P31016


- Molecule 2 is a protein called postsynaptic protein CRIPT peptide.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
2	B	6	110	35	57	8	10	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	LYS	GLN	engineered mutation	GB 3098551
B	5	GLU	SER	engineered mutation	GB 3098551

- Molecule 3 is BETA-ALANINE (three-letter code: BAL) (formula: C₃H₇NO₂).




Mol	Chain	Residues	Atoms				
			Total	C	H	N	O
3	B	1	10	3	5	1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Presynaptic density protein 95



- Molecule 2: postsynaptic protein CRIPT peptide




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Presynaptic density protein 95



- Molecule 2: postsynaptic protein CRIPT peptide







4.2.2 Score per residue for model 2

- Molecule 1: Presynaptic density protein 95

Chain A: 51% 19% 21% 6%



- Molecule 2: postsynaptic protein CRIPT peptide


Chain B: 83% 17%



4.2.3 Score per residue for model 3

- Molecule 1: Presynaptic density protein 95

Chain A: 53% 18% 21% 6%



- Molecule 2: postsynaptic protein CRIPT peptide


Chain B: 50% 33% 17%




4.2.4 Score per residue for model 4

- Molecule 1: Presynaptic density protein 95

Chain A: 54% 14% 21% 6%




- Molecule 2: postsynaptic protein CRIPT peptide




4.2.5 Score per residue for model 5

- Molecule 1: Presynaptic density protein 95




- Molecule 2: postsynaptic protein CRIPT peptide



4.2.6 Score per residue for model 6

- Molecule 1: Presynaptic density protein 95



- Molecule 2: postsynaptic protein CRIPT peptide






4.2.7 Score per residue for model 7 (medoid)

- Molecule 1: Presynaptic density protein 95

Chain A: 56% 16% 21% 6%



- Molecule 2: postsynaptic protein CRIPT peptide


Chain B: 67% 33%



4.2.8 Score per residue for model 8

- Molecule 1: Presynaptic density protein 95

Chain A: 51% 18% 21% 6%



- Molecule 2: postsynaptic protein CRIPT peptide


Chain B: 67% 33%



4.2.9 Score per residue for model 9

- Molecule 1: Presynaptic density protein 95

Chain A: 51% 17% 5% 21% 6%



HIS
HIS
HIS
HIS
HIS
HIS

- Molecule 2: postsynaptic protein CRIPT peptide

Chain B:  33% 67%

Y1
K2
K3
V6

4.2.10 Score per residue for model 10

- Molecule 1: Presynaptic density protein 95

Chain A:  56% 11% 5% 21% 6%

E62 E65 G70 G71 N72 S73 G74 L75 A80 N85 I88 G89 D90 D91 P92 K98 I99 Q107 E108 G109 R110 L111 R112 V113 M114 L118 M121 V125 R126 A130 I143 V144 R145 M149 R150 R151 K152 P153 P154 HIS HIS HIS HIS HIS HIS

- Molecule 2: postsynaptic protein CRIPT peptide

Chain B:  67% 33%

Y1
V6

4.2.11 Score per residue for model 11

- Molecule 1: Presynaptic density protein 95

Chain A:  57% 11% 5% 21% 6%

E62 E69 G71 N72 S73 G74 L75 F77 A80 N85 I88 G89 D90 P91 P92 I99 Q107 E108 G109 R110 L111 R112 V113 M114 M121 V125 R126 H130 A140 I143 V144 R145 V148 M149 R150 R151 K152 P153 P154 HIS HIS HIS HIS HIS HIS

- Molecule 2: postsynaptic protein CRIPT peptide

Chain B:  50% 50%

Y1
K2
K3
T4
E5
V6

4.2.12 Score per residue for model 12

- Molecule 1: Presynaptic density protein 95

Chain A:  51% 17% 21% 6%




- Molecule 2: postsynaptic protein CRIPT peptide



4.2.13 Score per residue for model 13

- Molecule 1: Presynaptic density protein 95




- Molecule 2: postsynaptic protein CRIPT peptide




4.2.14 Score per residue for model 14

- Molecule 1: Presynaptic density protein 95



- Molecule 2: postsynaptic protein CRIPT peptide






4.2.15 Score per residue for model 15

- Molecule 1: Presynaptic density protein 95

Chain A: 47% 21% • 21% 6%



- Molecule 2: postsynaptic protein CRIPT peptide

Chain B: 50% 33% 17%



4.2.16 Score per residue for model 16

- Molecule 1: Presynaptic density protein 95

Chain A: 53% 16% •• 21% 6%



- Molecule 2: postsynaptic protein CRIPT peptide


Chain B: 83% 17%



4.2.17 Score per residue for model 17

- Molecule 1: Presynaptic density protein 95

Chain A: 55% 15% • 21% 6%




- Molecule 2: postsynaptic protein CRIPT peptide




4.2.18 Score per residue for model 18

- Molecule 1: Presynaptic density protein 95





- Molecule 2: postsynaptic protein CRIPT peptide




4.2.19 Score per residue for model 19

- Molecule 1: Presynaptic density protein 95




- Molecule 2: postsynaptic protein CRIPT peptide




4.2.20 Score per residue for model 20

- Molecule 1: Presynaptic density protein 95




- Molecule 2: postsynaptic protein CRIPT peptide



4.2.21 Score per residue for model 21

- Molecule 1: Presynaptic density protein 95




- Molecule 2: postsynaptic protein CRIPT peptide



4.2.22 Score per residue for model 22

- Molecule 1: Presynaptic density protein 95



- Molecule 2: postsynaptic protein CRIPT peptide



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics, cartesian MD in water*.

Of the 100 calculated structures, 22 were deposited, based on the following criterion: *structures with acceptable covalent geometry, structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
CNS	refinement	1.1

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: BAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.83±0.03	0±0/551 (0.0± 0.0%)	1.38±0.03	4±1/750 (0.5± 0.1%)
2	B	0.87±0.04	0±0/53 (0.0± 0.0%)	1.35±0.11	0±0/67 (0.5± 0.7%)
All	All	0.83	0/13288 (0.0%)	1.38	94/17974 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.7±0.6
2	B	0.0±0.0	0.1±0.3
All	All	0	40

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	126	ARG	NE-CZ-NH1	10.40	125.50	120.30	6	22
1	A	145	ARG	NE-CZ-NH1	7.82	124.21	120.30	1	22
2	B	6	VAL	CA-CB-CG1	7.69	122.44	110.90	3	8
1	A	126	ARG	NE-CZ-NH2	-7.32	116.64	120.30	6	13
1	A	77	PHE	CB-CG-CD1	-6.28	116.40	120.80	11	2
1	A	92	PRO	N-CA-CB	-6.24	95.74	102.60	13	6
1	A	105	ALA	N-CA-CB	-5.75	102.05	110.10	15	1
1	A	118	LEU	CB-CA-C	5.63	120.90	110.20	21	2
1	A	85	ASN	N-CA-CB	-5.39	100.89	110.60	16	2
1	A	95	PHE	CB-CG-CD2	-5.36	117.05	120.80	2	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	77	PHE	CB-CG-CD2	-5.21	117.16	120.80	19	1
1	A	96	ILE	CB-CA-C	5.20	121.99	111.60	22	2
1	A	145	ARG	NE-CZ-NH2	-5.16	117.72	120.30	12	5
1	A	75	LEU	CB-CG-CD2	5.11	119.69	111.00	4	1
1	A	140	ALA	N-CA-CB	-5.08	102.99	110.10	13	2
1	A	147	TYR	CB-CG-CD2	-5.06	117.96	121.00	22	1
1	A	134	VAL	CA-CB-CG1	5.06	118.48	110.90	2	1
1	A	118	LEU	CB-CG-CD2	5.04	119.57	111.00	12	1
1	A	118	LEU	CA-CB-CG	5.04	126.88	115.30	1	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	80	ALA	Peptide	22
1	A	130	HIS	Sidechain	9
1	A	87	HIS	Peptide	3
2	B	1	TYR	Sidechain	2
1	A	95	PHE	Peptide	2
1	A	63	TYR	Sidechain	1
1	A	128	VAL	Peptide	1

6.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	541	538	536	4±2
2	B	53	57	57	0±1
All	All	13178	13200	13139	91

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:118:LEU:HD11	1:A:149:MET:H	0.67	1.49	12	3
1:A:121:ASN:HD21	1:A:140:ALA:HB2	0.66	1.49	5	17
1:A:134:VAL:HG22	2:B:6:VAL:HG21	0.64	1.69	12	1
1:A:75:LEU:H	1:A:75:LEU:HD23	0.62	1.53	5	1
1:A:79:ILE:HG22	1:A:96:ILE:HG12	0.59	1.75	19	7
1:A:118:LEU:HD21	1:A:149:MET:HB3	0.57	1.77	13	4
1:A:126:ARG:HD2	1:A:127:GLU:H	0.55	1.62	14	2
1:A:79:ILE:HG22	1:A:96:ILE:HD13	0.54	1.79	9	1
1:A:75:LEU:H	1:A:75:LEU:HD22	0.53	1.62	13	1
1:A:75:LEU:HD21	2:B:6:VAL:HG22	0.51	1.83	14	2
1:A:75:LEU:HD11	1:A:144:VAL:HG11	0.51	1.81	10	2
1:A:123:VAL:HG12	1:A:125:VAL:HG13	0.50	1.82	2	2
1:A:76:GLY:HA3	1:A:103:GLY:HA3	0.47	1.86	8	4
1:A:118:LEU:HA	1:A:126:ARG:HB3	0.46	1.85	21	4
1:A:134:VAL:HG23	2:B:6:VAL:HB	0.45	1.89	3	1
1:A:137:LEU:HD11	1:A:146:LEU:HD21	0.45	1.88	4	1
1:A:75:LEU:HD13	2:B:6:VAL:HG22	0.45	1.88	20	2
1:A:75:LEU:HD23	2:B:6:VAL:HG12	0.45	1.89	11	1
1:A:83:THR:HG22	1:A:92:PRO:HB2	0.45	1.88	5	3
1:A:134:VAL:HG23	2:B:6:VAL:HG22	0.45	1.88	21	1
1:A:125:VAL:HG11	1:A:133:ALA:HA	0.45	1.87	2	4
1:A:87:HIS:HB2	2:B:1:TYR:CG	0.44	2.48	13	1
1:A:118:LEU:HD23	1:A:149:MET:HB2	0.44	1.88	15	2
1:A:94:ILE:HG13	1:A:133:ALA:CB	0.44	2.43	6	1
1:A:78:SER:O	1:A:96:ILE:HG23	0.43	2.13	20	1
1:A:131:SER:HA	1:A:134:VAL:HG12	0.43	1.89	5	4
1:A:126:ARG:H	1:A:128:VAL:HG22	0.43	1.73	6	1
1:A:117:ILE:HD12	1:A:137:LEU:HD21	0.43	1.90	4	1
1:A:75:LEU:HD23	2:B:6:VAL:HG22	0.43	1.89	15	1
1:A:121:ASN:HD21	1:A:140:ALA:CB	0.43	2.26	10	1
1:A:75:LEU:HB2	2:B:6:VAL:HG13	0.43	1.90	15	1
1:A:75:LEU:HD23	1:A:75:LEU:N	0.43	2.29	4	3
1:A:75:LEU:HD12	1:A:144:VAL:HG21	0.42	1.92	13	1
1:A:120:VAL:HG21	1:A:137:LEU:HD12	0.42	1.91	5	1
1:A:79:ILE:CG2	1:A:96:ILE:HD13	0.41	2.45	9	1
1:A:115:ASP:HB3	1:A:148:VAL:HG11	0.41	1.91	9	1
1:A:117:ILE:HD12	1:A:137:LEU:HD11	0.41	1.93	5	1
1:A:140:ALA:HB1	1:A:144:VAL:HA	0.41	1.93	15	1
1:A:117:ILE:CD1	1:A:137:LEU:HD21	0.40	2.46	4	1
1:A:137:LEU:O	1:A:140:ALA:HB3	0.40	2.15	12	1
1:A:95:PHE:HA	1:A:115:ASP:O	0.40	2.17	8	1
1:A:120:VAL:HG21	1:A:137:LEU:HD22	0.40	1.94	8	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	72/99 (73%)	61±2 (84±2%)	7±2 (10±3%)	4±1 (6±1%)	3	21
2	B	4/6 (67%)	4±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
All	All	1672/2310 (72%)	1424 (85%)	155 (9%)	93 (6%)	3	22

All 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	125	VAL	21
1	A	126	ARG	20
1	A	121	ASN	17
1	A	92	PRO	11
1	A	122	GLU	9
1	A	86	PRO	7
1	A	115	ASP	4
1	A	75	LEU	1
1	A	141	GLY	1
1	A	140	ALA	1
1	A	76	GLY	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	58/81 (72%)	50±2 (86±3%)	8±2 (14±3%)	7	47
2	B	6/6 (100%)	4±1 (71±11%)	2±1 (29±11%)	2	18
All	All	1408/1914 (74%)	1196 (85%)	212 (15%)	6	44

All 33 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	143	ILE	22
1	A	99	ILE	20
1	A	107	GLN	18
1	A	85	ASN	16
2	B	1	TYR	14
1	A	125	VAL	12
1	A	118	LEU	11
2	B	5	GLU	11
1	A	77	PHE	10
1	A	65	GLU	10
1	A	75	LEU	10
1	A	148	VAL	8
2	B	3	LYS	7
1	A	126	ARG	6
1	A	98	LYS	5
1	A	68	LEU	4
1	A	135	GLU	4
1	A	134	VAL	3
2	B	4	THR	2
1	A	117	ILE	2
2	B	2	LYS	2
1	A	139	GLU	2
2	B	6	VAL	2
1	A	122	GLU	2
1	A	83	THR	1
1	A	127	GLU	1
1	A	129	THR	1
1	A	92	PRO	1
1	A	97	THR	1
1	A	64	GLU	1
1	A	69	GLU	1
1	A	66	ILE	1
1	A	78	SER	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
3	BAL	B	7	2	4,4,5	0.70±0.06	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
3	BAL	B	7	2	3,3,5	2.62±1.02	1±0 (25±13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BAL	B	7	2	-	0±0,1,2,3	-

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)	Models	
								Worst	Total
3	B	7	BAL	CB-CA-C	6.81	121.54	111.42	14	17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided