



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 15, 2023 – 04:56 AM EDT

PDB ID : 1TP5  
Title : Crystal structure of PDZ3 domain of PSD-95 protein complexed with a peptide ligand KKETWV  
Authors : Saro, D.; Wawrzak, Z.; Martin, P.; Vickrey, J.; Paredes, A.; Kovari, L.; Spaller, M.  
Deposited on : 2004-06-15  
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	119	<p>2% (Poor fit), 85% (0 outliers), 9% (1 outlier), 4% (2 outliers), 2% (3+ outliers), 0% (Not modelled)</p>
2	B	6	<p>67% (0 outliers), 17% (1 outlier), 17% (2 outliers)</p>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Presynaptic density protein 95.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	115	863	533	158	172	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	297	GLY	-	cloning artifact	UNP P31016
A	298	SER	-	cloning artifact	UNP P31016
A	299	PRO	-	cloning artifact	UNP P31016
A	300	GLU	-	cloning artifact	UNP P31016
A	301	PHE	-	cloning artifact	UNP P31016
A	403	ASN	-	cloning artifact	UNP P31016
A	404	SER	-	cloning artifact	UNP P31016
A	405	ARG	-	cloning artifact	UNP P31016
A	406	VAL	-	cloning artifact	UNP P31016
A	407	ASP	-	cloning artifact	UNP P31016
A	408	SER	-	cloning artifact	UNP P31016
A	409	SER	-	cloning artifact	UNP P31016
A	410	GLY	-	cloning artifact	UNP P31016
A	411	ARG	-	cloning artifact	UNP P31016
A	412	ILE	-	cloning artifact	UNP P31016
A	413	VAL	-	cloning artifact	UNP P31016
A	414	THR	-	cloning artifact	UNP P31016
A	415	ASP	-	cloning artifact	UNP P31016

- Molecule 2 is a protein called LYS-LYS-GLU-THR-TRP-VAL peptide ligand.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	56	37	9	10	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	176	Total 176	O 176	0	0
3	B	11	Total 11	O 11	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Presynaptic density protein 95



- Molecule 2: LYS-LYS-GLU-THR-TRP-VAL peptide ligand



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.98Å 89.98Å 89.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.65 – 1.54 27.13 – 1.54	Depositor EDS
% Data completeness (in resolution range)	96.4 (19.65-1.54) 96.4 (27.13-1.54)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.61 (at 1.54Å)	Xtrriage
Refinement program	REFMAC 5.1.24, XTALVIEW	Depositor
R, $R_{free}$	0.191 , 0.229 0.206 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.6	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 45.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1106	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/874	0.87	0/1177
2	B	1.31	1/57 (1.8%)	0.85	0/74
All	All	0.82	1/931 (0.1%)	0.87	0/1251

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	424	TRP	CB-CG	-7.42	1.36	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	863	0	847	12	0
2	B	56	0	57	4	0
3	A	176	0	0	5	1
3	B	11	0	0	0	1
All	All	1106	0	904	12	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:GLU:O	1:A:334:GLU:HG2	1.67	0.94
1:A:396:GLU:OE1	3:A:168:HOH:O	1.93	0.85
1:A:342:LEU:HB2	2:B:424:TRP:HH2	1.49	0.77
1:A:301:PHE:N	3:A:170:HOH:O	2.28	0.65
1:A:326:ASN:OD1	2:B:424:TRP:CD2	2.53	0.62
1:A:331:GLU:O	1:A:334:GLU:CG	2.47	0.56
1:A:352:GLU:CD	3:A:137:HOH:O	2.50	0.50
1:A:326:ASN:OD1	2:B:424:TRP:CE2	2.67	0.47
1:A:399:ARG:NE	3:A:61:HOH:O	2.47	0.47
1:A:310:GLU:HB3	3:A:31:HOH:O	2.14	0.47
1:A:307:ILE:HD11	1:A:411:ARG:HG2	1.96	0.46
1:A:342:LEU:HB2	2:B:424:TRP:CH2	2.40	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:183:HOH:O	3:A:183:HOH:O[14_555]	2.12	0.08
3:B:182:HOH:O	3:B:182:HOH:O[14_555]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	113/119 (95%)	112 (99%)	1 (1%)	0	100	100
2	B	4/6 (67%)	4 (100%)	0	0	100	100
All	All	117/125 (94%)	116 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	89/94 (95%)	83 (93%)	6 (7%)	16 1
2	B	6/6 (100%)	5 (83%)	1 (17%)	2 0
All	All	95/100 (95%)	88 (93%)	7 (7%)	13 1

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	304	GLU
1	A	310	GLU
1	A	312	ARG
1	A	318	ARG
1	A	334	GLU
1	A	399	ARG
2	B	420	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	381	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	115/119 (96%)	-0.01	2 (1%) 70 74	10, 16, 29, 40	0
2	B	6/6 (100%)	2.80	4 (66%) 0 0	21, 36, 40, 42	0
All	All	121/125 (96%)	0.13	6 (4%) 28 32	10, 17, 36, 42	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	424	TRP	5.5
1	A	301	PHE	4.8
2	B	422	GLU	4.5
1	A	304	GLU	4.0
2	B	420	LYS	2.6
2	B	421	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.