



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 01:21 AM UTC

PDB ID : 1U6N / pdb_00001u6n
Title : Solution Structure of an Oligodeoxynucleotide Containing a Butadiene Derived N1 b-Hydroxyalkyl Adduct on Deoxyinosine in the Human N-ras Codon 61 Sequence
Authors : Scholdberg, T.A.; Merritt, W.K.; Dean, S.M.; Kowalczyk, A.; Harris, T.M.; Harris, C.M.; Lloyd, R.S.; Stone, M.P.
Deposited on : 2004-07-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR



The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	11	
2	B	11	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
1	A	2BD	6	1	-

2 Ensemble composition and analysis

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 706 atoms, of which 256 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*GP*AP*CP*(2BD)P*AP*GP*AP*AP*G)-3'.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	11	361	112	128	50	61	10	0

- Molecule 2 is a DNA chain called 5'-D(*CP*TP*TP*CP*TP*TP*GP*TP*CP*CP*G)-3'.


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	B	11	345	106	128	32	69	10	0

4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-D(*CP*GP*GP*AP*CP*(2BD)P*AP*GP*AP*AP*G)-3'

Chain A: 



- Molecule 2: 5'-D(*CP*TP*TP*CP*TP*TP*GP*TP*CP*CP*G)-3'

Chain B: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry, simulated annealing, molecular dynamics, matrix relaxation, torsion angle dynamics*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: ?.

The authors did not provide any information on software used for structure solution, optimization or refinement.

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2BD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.73	0/233 (0.0%)	1.36	2/356 (0.6%)
2	B	0.76	0/240 (0.0%)	1.29	1/368 (0.3%)
All	All	0.75	0/473 (0.0%)	1.33	3/724 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	1	0
All	All	1	0

There are no bond-length outliers.

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	DC	O4'-C1'-N1	12.41	127.01	108.40
1	A	1	DC	N1-C1'-C2'	-6.08	104.39	113.50
2	B	17	DT	N1-C1'-C2'	5.66	121.99	113.50

All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	6	2BD	C3'

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	233	128	127	30
2	B	217	128	128	9
All	All	450	256	255	37

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 52.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:6:2BD:CAN	1:A:6:2BD:N1	1.54	1.70
1:A:5:DC:C2'	1:A:6:2BD:C5'	0.95	2.43
1:A:5:DC:H2''	1:A:6:2BD:C5'	0.95	1.90
1:A:6:2BD:H1'	1:A:7:DA:C8	0.91	1.99
1:A:5:DC:H2''	1:A:6:2BD:O5'	0.86	1.71
1:A:6:2BD:H2'	1:A:6:2BD:N3	0.85	1.86
1:A:5:DC:C1'	1:A:6:2BD:H5'	0.84	2.02
1:A:6:2BD:N1	1:A:6:2BD:CAO	0.83	2.40
1:A:5:DC:C2'	1:A:6:2BD:H5'	0.82	2.04
1:A:5:DC:H1'	1:A:6:2BD:C5'	0.81	2.05
1:A:5:DC:C1'	1:A:6:2BD:C5'	0.78	2.62
1:A:5:DC:H2''	1:A:6:2BD:H5'	0.78	1.56
1:A:5:DC:H1'	1:A:6:2BD:H5'	0.77	1.55
1:A:5:DC:C2'	1:A:6:2BD:O5'	0.73	2.34
1:A:6:2BD:CAN	1:A:6:2BD:C2	0.71	2.68
1:A:6:2BD:C1'	1:A:7:DA:C8	0.70	2.75
1:A:6:2BD:C8	1:A:7:DA:C5	0.69	2.74
2:B:16:DT:H2'	2:B:17:DT:C5	0.67	2.25
1:A:6:2BD:N1	1:A:6:2BD:HAO1	0.65	2.05
1:A:6:2BD:C2	1:A:6:2BD:HAO1	0.65	2.22
2:B:16:DT:H2'	2:B:17:DT:C6	0.64	2.28
1:A:6:2BD:CAO	1:A:6:2BD:C2	0.60	2.79
1:A:4:DA:H2''	1:A:5:DC:C6	0.58	2.34
2:B:16:DT:C2'	2:B:17:DT:C6	0.57	2.87
1:A:6:2BD:N3	1:A:6:2BD:C2'	0.57	2.67
1:A:6:2BD:H1'	1:A:7:DA:N9	0.56	2.15
2:B:19:DT:H2''	2:B:20:DC:OP2	0.53	2.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:4:DA:H2''	1:A:5:DC:C5	0.53	2.39
2:B:14:DT:H2''	2:B:15:DC:O5'	0.53	2.03
1:A:6:2BD:C8	1:A:7:DA:C6	0.52	2.93
1:A:5:DC:H1'	1:A:6:2BD:H5''	0.48	1.81
1:A:5:DC:C3'	1:A:6:2BD:C5'	0.47	2.88
2:B:15:DC:H2''	2:B:16:DT:OP2	0.45	2.11
1:A:3:DG:N2	2:B:21:DC:C2	0.44	2.86
2:B:20:DC:C2	2:B:21:DC:C5	0.44	3.06
1:A:4:DA:N6	2:B:18:DG:O6	0.43	2.52
1:A:3:DG:C6	1:A:4:DA:C6	0.42	3.07

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	2BD	A	6	1	24,28,29	2.57	9 (37%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	2BD	A	6	1	29,40,43	2.76	13 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2BD	A	6	1	1,1,6,6	0,14,29,30	0,3,3,3

All bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	2BD	CAN-N1	6.11	1.70	1.48
1	A	6	2BD	C6-N1	5.09	1.49	1.38
1	A	6	2BD	O4'-C1'	4.64	1.52	1.42
1	A	6	2BD	C2'-C3'	3.90	1.43	1.52
1	A	6	2BD	C2-N1	3.44	1.44	1.36
1	A	6	2BD	C8-N9	3.38	1.29	1.37
1	A	6	2BD	C1'-N9	3.16	1.56	1.48
1	A	6	2BD	C5-N7	2.92	1.33	1.39
1	A	6	2BD	C5-C6	2.31	1.39	1.45

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	2BD	O3'-C3'-C2'	5.66	130.56	110.88
1	A	6	2BD	C2-N3-C4	5.65	123.61	112.53
1	A	6	2BD	C2'-C3'-C4'	5.39	113.73	102.80
1	A	6	2BD	C5-C4-N3	5.25	119.55	127.27
1	A	6	2BD	C2'-C1'-N9	3.92	104.02	113.81
1	A	6	2BD	O6-C6-C5	3.57	115.40	124.59
1	A	6	2BD	CAN-CAL-CAM	3.56	108.97	125.49
1	A	6	2BD	O4'-C4'-C5'	2.92	118.68	109.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	2BD	N9-C4-N3	2.84	133.38	126.90
1	A	6	2BD	C5-C6-N1	2.82	119.81	112.54
1	A	6	2BD	N1-C2-N3	2.61	122.64	125.84
1	A	6	2BD	C4-C5-N7	2.21	107.17	110.67
1	A	6	2BD	O4'-C4'-C3'	2.18	100.68	105.65

All chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms
1	A	6	2BD	C3'

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided