



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 27, 2024 – 05:48 pm BST

PDB ID : 1UO1
Title : Structure Based Engineering of Internal Molecular Surfaces Of Four Helix Bundles
Authors : Yadav, M.K.; Redman, J.E.; Alvarez-Gutierrez, J.M.; Zhang, Y.; Stout, C.D.; Ghadiri, M.R.
Deposited on : 2003-09-15
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	34	
1	B	34	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 523 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	31	261	166	46	48	1	0	0	0
1	B	32	262	166	47	48	1	0	0	1

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	ILE	LEU	conflict	UNP P03069
A	9	LEU	VAL	conflict	UNP P03069
A	12	ILE	LEU	conflict	UNP P03069
A	16	LEU	ASN	conflict	UNP P03069
A	19	ILE	LEU	conflict	UNP P03069
A	23	LEU	VAL	conflict	UNP P03069
A	26	THR	LEU	engineered mutation	UNP P03069
A	30	LEU	VAL	conflict	UNP P03069
B	5	ILE	LEU	conflict	UNP P03069
B	9	LEU	VAL	conflict	UNP P03069
B	12	ILE	LEU	conflict	UNP P03069
B	16	LEU	ASN	conflict	UNP P03069
B	19	ILE	LEU	conflict	UNP P03069
B	23	LEU	VAL	conflict	UNP P03069
B	26	THR	LEU	engineered mutation	UNP P03069
B	30	LEU	VAL	conflict	UNP P03069

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A: 



- Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	79.39Å 79.39Å 79.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.90 – 2.50 28.07 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (55.90-2.50) 100.0 (28.07-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.36 (at 2.51Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.231 , 0.302 0.231 , 0.296	Depositor DCC
R_{free} test set	235 reflections (4.20%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 61.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	523	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.39	26/262 (9.9%)	2.30	12/346 (3.5%)
1	B	3.52	31/263 (11.8%)	2.70	24/348 (6.9%)
All	All	3.46	57/525 (10.9%)	2.51	36/694 (5.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (57) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	20	GLU	CD-OE2	19.35	1.47	1.25
1	B	11	GLU	CD-OE1	19.14	1.46	1.25
1	A	6	GLU	CD-OE2	14.91	1.42	1.25
1	B	22	GLU	CD-OE1	14.35	1.41	1.25
1	A	10	GLU	CD-OE2	11.96	1.38	1.25
1	B	6	GLU	CD-OE2	11.49	1.38	1.25
1	A	20	GLU	CD-OE1	11.45	1.38	1.25
1	B	17	TYR	CE1-CZ	-10.78	1.24	1.38
1	A	28	LYS	CD-CE	10.53	1.77	1.51
1	B	10	GLU	CD-OE1	9.52	1.36	1.25
1	A	8	LYS	CD-CE	9.37	1.74	1.51
1	B	25	ARG	NE-CZ	9.12	1.45	1.33
1	B	17	TYR	CG-CD2	-8.66	1.27	1.39
1	A	14	SER	CB-OG	-8.64	1.31	1.42
1	B	16	LEU	C-O	-8.60	1.07	1.23
1	B	20	GLU	CD-OE1	7.87	1.34	1.25
1	B	17	TYR	CB-CG	-7.84	1.39	1.51
1	B	22	GLU	CD-OE2	-7.81	1.17	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	8	LYS	CD-CE	7.62	1.70	1.51
1	B	26	THR	CA-CB	-7.36	1.34	1.53
1	B	25	ARG	CG-CD	7.23	1.70	1.51
1	A	22	GLU	CD-OE2	-7.22	1.17	1.25
1	A	17	TYR	CD2-CE2	7.18	1.50	1.39
1	B	20	GLU	CD-OE2	-7.16	1.17	1.25
1	B	26	THR	CA-C	-7.05	1.34	1.52
1	B	6	GLU	CG-CD	6.92	1.62	1.51
1	A	4	GLN	CD-OE1	6.87	1.39	1.24
1	B	23	LEU	CG-CD2	-6.84	1.26	1.51
1	A	28	LYS	CG-CD	6.80	1.75	1.52
1	B	28	LYS	CE-NZ	6.69	1.65	1.49
1	A	25	ARG	CB-CG	-6.66	1.34	1.52
1	B	17	TYR	CZ-OH	6.54	1.49	1.37
1	B	25	ARG	CD-NE	6.48	1.57	1.46
1	B	23	LEU	C-O	6.40	1.35	1.23
1	B	8	LYS	CG-CD	6.35	1.74	1.52
1	B	7	ASP	C-O	6.29	1.35	1.23
1	A	17	TYR	CG-CD2	-6.26	1.31	1.39
1	A	7	ASP	CB-CG	6.24	1.64	1.51
1	A	28	LYS	CE-NZ	6.19	1.64	1.49
1	B	28	LYS	CD-CE	6.08	1.66	1.51
1	A	24	ALA	CA-CB	6.07	1.65	1.52
1	A	8	LYS	C-O	-5.97	1.11	1.23
1	B	11	GLU	CG-CD	5.96	1.60	1.51
1	A	6	GLU	CD-OE1	5.93	1.32	1.25
1	B	8	LYS	CE-NZ	5.86	1.63	1.49
1	B	8	LYS	CB-CG	5.76	1.68	1.52
1	A	3	LYS	CA-C	-5.69	1.38	1.52
1	B	30	LEU	C-O	-5.65	1.12	1.23
1	A	11	GLU	C-O	-5.57	1.12	1.23
1	B	15	LYS	CE-NZ	5.44	1.62	1.49
1	A	21	ASN	C-O	-5.43	1.13	1.23
1	B	15	LYS	C-O	-5.38	1.13	1.23
1	A	18	HIS	CA-CB	-5.35	1.42	1.53
1	A	17	TYR	CZ-OH	5.10	1.46	1.37
1	A	9	LEU	CA-CB	5.07	1.65	1.53
1	A	6	GLU	CG-CD	5.06	1.59	1.51
1	A	26	THR	CB-CG2	5.02	1.69	1.52

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	31	GLY	O-C-N	-13.22	101.55	122.70
1	B	7	ASP	CB-CG-OD1	11.29	128.46	118.30
1	B	22	GLU	CG-CD-OE1	10.61	139.52	118.30
1	B	25	ARG	NE-CZ-NH2	9.70	125.15	120.30
1	B	13	LEU	CB-CG-CD2	-9.68	94.54	111.00
1	B	31	GLY	CA-C-N	8.99	136.98	117.20
1	A	16	LEU	CB-CG-CD2	-8.80	96.04	111.00
1	A	25	ARG	NE-CZ-NH1	-8.69	115.95	120.30
1	B	11	GLU	CG-CD-OE2	-8.55	101.20	118.30
1	B	16	LEU	CB-CG-CD2	-8.45	96.64	111.00
1	B	22	GLU	CG-CD-OE2	-8.30	101.71	118.30
1	A	22	GLU	OE1-CD-OE2	-8.17	113.50	123.30
1	B	10	GLU	OE1-CD-OE2	6.90	131.58	123.30
1	A	27	LYS	CD-CE-NZ	6.87	127.49	111.70
1	B	25	ARG	NH1-CZ-NH2	-6.44	112.31	119.40
1	A	17	TYR	CG-CD1-CE1	6.34	126.37	121.30
1	B	9	LEU	CB-CG-CD1	-6.33	100.23	111.00
1	A	10	GLU	CG-CD-OE1	-6.28	105.74	118.30
1	B	30	LEU	CB-CG-CD2	-6.18	100.50	111.00
1	B	28	LYS	CD-CE-NZ	5.93	125.33	111.70
1	A	14	SER	CB-CA-C	-5.91	98.87	110.10
1	B	11	GLU	CG-CD-OE1	5.82	129.95	118.30
1	B	2	MET	CG-SD-CE	5.74	109.39	100.20
1	B	22	GLU	OE1-CD-OE2	-5.65	116.52	123.30
1	A	22	GLU	CB-CA-C	-5.53	99.33	110.40
1	B	6	GLU	CG-CD-OE2	5.52	129.34	118.30
1	B	23	LEU	CA-CB-CG	5.46	127.86	115.30
1	B	10	GLU	CA-CB-CG	-5.43	101.46	113.40
1	A	21	ASN	CB-CA-C	-5.34	99.72	110.40
1	A	19	ILE	CA-CB-CG2	-5.34	100.22	110.90
1	A	4	GLN	CB-CG-CD	5.13	124.94	111.60
1	B	15	LYS	CB-CA-C	-5.11	100.19	110.40
1	B	17	TYR	CB-CG-CD1	-5.07	117.96	121.00
1	A	6	GLU	CB-CA-C	-5.05	100.30	110.40
1	B	17	TYR	CB-CG-CD2	5.02	124.01	121.00
1	B	20	GLU	N-CA-CB	-5.01	101.59	110.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	26	THR	Mainchain
1	B	4	GLN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	261	0	285	13	0
1	B	262	0	285	4	0
All	All	523	0	570	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:LYS:CD	1:A:8:LYS:CE	1.74	1.66
1:A:5:ILE:CD1	1:A:5:ILE:CG1	1.74	1.57
1:A:28:LYS:CD	1:A:28:LYS:CE	1.77	1.56
1:A:28:LYS:CD	1:A:28:LYS:CG	1.75	1.53
1:A:8:LYS:CD	1:A:8:LYS:NZ	2.55	0.70
1:B:1:ARG:HG2	1:B:1:ARG:HH11	1.55	0.70
1:A:3:LYS:HE2	1:A:7:ASP:OD1	1.92	0.69
1:A:8:LYS:CE	1:A:8:LYS:CG	2.67	0.69
1:A:5:ILE:CD1	1:A:5:ILE:CB	2.73	0.60
1:B:8:LYS:O	1:B:12:ILE:HG13	2.01	0.60
1:A:28:LYS:CD	1:A:28:LYS:CB	2.77	0.59
1:A:1:ARG:HG3	1:A:3:LYS:H	1.69	0.57
1:A:30:LEU:O	1:A:31:GLY:C	2.51	0.47
1:A:6:GLU:HG3	1:B:5:ILE:HG13	1.97	0.46
1:A:2:MET:HG3	1:B:5:ILE:CD1	2.47	0.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	29/34 (85%)	25 (86%)	4 (14%)	0	100	100
1	B	30/34 (88%)	24 (80%)	5 (17%)	1 (3%)	4	5
All	All	59/68 (87%)	49 (83%)	9 (15%)	1 (2%)	9	16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	6	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	29/31 (94%)	22 (76%)	7 (24%)	0	1
1	B	29/31 (94%)	21 (72%)	8 (28%)	0	0
All	All	58/62 (94%)	43 (74%)	15 (26%)	0	1

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	MET
1	A	4	GLN
1	A	13	LEU
1	A	14	SER
1	A	23	LEU
1	A	28	LYS
1	A	30	LEU
1	B	2	MET
1	B	3	LYS
1	B	8	LYS
1	B	15	LYS
1	B	23	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	27	LYS
1	B	29	LEU
1	B	30	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	21	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	31/34 (91%)	-0.40	0 100 100	17, 30, 57, 62	0
1	B	32/34 (94%)	-0.37	0 100 100	15, 30, 64, 66	0
All	All	63/68 (92%)	-0.38	0 100 100	15, 30, 62, 66	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.