



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 11, 2024 – 09:21 PM EST

PDB ID : 1X9Z  
Title : Crystal structure of the MutL C-terminal domain  
Authors : Guarne, A.; Ramon-Maiques, S.; Wolff, E.M.; Ghirlando, R.; Hu, X.; Miller, J.H.; Yang, W.  
Deposited on : 2004-08-24  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	188	 7% 74% 22% ..
1	B	188	 11% 66% 28% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NA	A	423	-	-	-	X
4	GOL	B	402	-	X	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3140 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA mismatch repair protein mutL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	182	1409	898	250	256	3	2	0	1	0
1	B	182	1418	904	251	256	4	3	0	4	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	428	GLY	-	cloning artifact	UNP P23367
A	429	SER	-	cloning artifact	UNP P23367
A	430	HIS	-	cloning artifact	UNP P23367
A	431	MET	-	cloning artifact	UNP P23367
A	566	MSE	MET	modified residue	UNP P23367
A	574	MSE	MET	modified residue	UNP P23367
B	428	GLY	-	cloning artifact	UNP P23367
B	429	SER	-	cloning artifact	UNP P23367
B	430	HIS	-	cloning artifact	UNP P23367
B	431	MET	-	cloning artifact	UNP P23367
B	566	MSE	MET	modified residue	UNP P23367
B	574	MSE	MET	modified residue	UNP P23367

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cl	0	0
			2	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula:  $C_3H_8O$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	3	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	3	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	154	Total	O	0	0
			154	154		
6	B	136	Total	O	0	0
			136	136		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.99Å 95.99Å 140.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.61 – 2.10 19.61 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.4 (19.61-2.10) 94.3 (19.61-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.55 (at 2.09Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.236 , 0.278 0.229 , 0.269	Depositor DCC
$R_{free}$ test set	3678 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.9	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3140	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.96 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9372e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA, CL, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/1437	0.66	0/1953
1	B	0.39	0/1458	0.63	0/1980
All	All	0.40	0/2895	0.64	0/3933

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1409	0	1465	52	0
1	B	1418	0	1476	57	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	7	6	0
5	A	4	0	8	2	0
5	B	4	0	8	0	0
6	A	154	0	0	8	0
6	B	136	0	0	15	0
All	All	3140	0	2972	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:464:GLU:HG2	5:A:411:IPA:H32	1.44	0.95
1:B:479:VAL:HG12	4:B:402:GOL:O1	1.71	0.89
1:A:562:ALA:HA	1:A:566:MSE:CE	2.03	0.89
1:B:456:SER:HB3	1:B:600:LEU:HD11	1.55	0.86
1:A:562:ALA:HA	1:A:566:MSE:HE3	1.60	0.81
1:A:568:GLU:HA	6:A:230:HOH:O	1.87	0.75
1:A:468:ARG:HB3	1:A:566:MSE:HE1	1.71	0.72
1:A:551:VAL:HA	6:A:323:HOH:O	1.89	0.71
1:A:485:LEU:HG	1:A:531:ARG:HH22	1.57	0.70
1:B:456:SER:CB	1:B:600:LEU:HD11	2.21	0.70
1:B:536:GLN:H	1:B:536:GLN:HE21	1.43	0.66
1:A:538:LEU:CD1	1:A:565:LEU:HB3	2.27	0.65
1:B:566:MSE:HE3	6:B:312:HOH:O	1.95	0.65
1:B:517:ASP:HB3	6:B:303:HOH:O	1.96	0.65
1:A:465[A]:ARG:HD2	1:A:563:ARG:HG2	1.80	0.64
1:A:560:TRP:HZ3	1:A:565:LEU:HD11	1.62	0.64
1:A:465[A]:ARG:HD3	1:A:563:ARG:HA	1.80	0.64
1:A:465[B]:ARG:HD2	1:A:563:ARG:HG2	1.80	0.64
1:B:503:ALA:HB2	1:B:552:PHE:CD2	2.33	0.64
1:A:486:ILE:O	1:A:486:ILE:HG13	1.99	0.63
1:B:479:VAL:HB	4:B:402:GOL:H12	1.79	0.63
1:B:569:HIS:HB3	6:B:319:HOH:O	1.97	0.63
1:B:479:VAL:HB	4:B:402:GOL:C1	2.27	0.63
1:A:562:ALA:HA	1:A:566:MSE:HE2	1.81	0.62
1:A:538:LEU:HD12	1:A:565:LEU:HB3	1.80	0.62
1:A:568:GLU:H	1:A:568:GLU:CD	2.02	0.62
1:B:444:SER:HA	1:B:571:GLN:HE22	1.67	0.59
1:B:486:ILE:HG23	1:B:486:ILE:O	2.03	0.59
1:B:536:GLN:H	1:B:536:GLN:NE2	2.01	0.58
1:B:539:ILE:HB	1:B:540:PRO:HD3	1.85	0.57
1:B:465[B]:ARG:HH21	1:B:469:GLN:NE2	2.03	0.56
1:B:465[B]:ARG:HH21	1:B:469:GLN:HE22	1.53	0.56
1:A:466:TRP:CG	1:A:591:LEU:HD11	2.40	0.56
1:A:495:GLU:HG2	6:A:181:HOH:O	2.07	0.55
1:B:586:ARG:HH11	1:B:587:LEU:HG	1.70	0.55
1:B:465[B]:ARG:NH2	1:B:469:GLN:HE22	2.05	0.55
1:A:536:GLN:NE2	1:B:491:LYS:H	2.03	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:456:SER:HB3	1:B:600:LEU:CD1	2.34	0.54
1:A:541:GLU:HB3	1:A:565:LEU:CD2	2.38	0.54
1:A:433:GLN:HB2	1:A:592:VAL:HG11	1.90	0.54
1:A:538:LEU:HD12	1:A:565:LEU:HD13	1.89	0.53
1:B:434:SER:HB2	1:B:585:GLU:OE2	2.08	0.53
1:B:493:SER:OG	1:B:496:GLU:HG3	2.08	0.53
1:B:465[B]:ARG:HD2	1:B:563:ARG:HG2	1.91	0.53
1:B:439:LEU:HD11	1:B:450:GLU:HB2	1.91	0.53
1:B:479:VAL:CG1	4:B:402:GOL:O1	2.52	0.53
1:B:586:ARG:NH1	1:B:587:LEU:HG	2.24	0.52
6:A:107:HOH:O	1:B:496:GLU:HG2	2.10	0.52
1:B:465[A]:ARG:NE	6:B:305:HOH:O	2.43	0.52
1:A:516:SER:HB2	1:A:520:HIS:O	2.10	0.51
1:B:464:GLU:OE1	1:B:568:GLU:HG2	2.09	0.51
1:B:469:GLN:HG2	6:B:157:HOH:O	2.10	0.51
1:A:444:SER:HA	1:A:571:GLN:OE1	2.11	0.51
1:B:466:TRP:CG	1:B:591:LEU:HD11	2.45	0.51
1:A:468:ARG:HB3	1:A:566:MSE:CE	2.41	0.50
1:A:560:TRP:CZ3	1:A:565:LEU:HG	2.47	0.50
1:B:444:SER:HA	1:B:571:GLN:NE2	2.26	0.50
1:B:567:SER:N	6:B:312:HOH:O	2.45	0.50
1:B:465[B]:ARG:HD2	6:B:308:HOH:O	2.12	0.49
1:B:465[B]:ARG:HD3	6:B:293:HOH:O	2.11	0.49
1:A:433:GLN:HB3	6:A:264:HOH:O	2.12	0.49
1:A:444:SER:HB3	1:A:571:GLN:HE22	1.77	0.49
1:B:595:PRO:HG3	6:B:318:HOH:O	2.13	0.49
1:A:592:VAL:O	1:A:595:PRO:HD3	2.12	0.49
1:B:468:ARG:NH1	6:B:306:HOH:O	2.40	0.49
1:B:586:ARG:HH11	1:B:586:ARG:HG3	1.78	0.48
1:A:466:TRP:CB	1:A:591:LEU:HD11	2.43	0.48
1:B:465[A]:ARG:NH1	6:B:292:HOH:O	2.47	0.48
1:B:534:ASN:HB3	1:B:536:GLN:HE22	1.79	0.47
1:B:537:ILE:O	1:B:541:GLU:HG3	2.15	0.47
1:B:443:HIS:O	1:B:445:ASP:N	2.47	0.47
1:B:518:ALA:N	6:B:303:HOH:O	2.47	0.47
1:B:534:ASN:CA	1:B:536:GLN:HE22	2.28	0.46
1:B:438:VAL:HA	1:B:449:LEU:HD23	1.96	0.46
1:B:479:VAL:CG1	4:B:402:GOL:C1	2.94	0.46
1:A:569:HIS:HB2	5:A:411:IPA:H31	1.96	0.45
1:A:493:SER:OG	1:A:496:GLU:HG3	2.17	0.45
1:A:485:LEU:CG	1:A:531:ARG:HH22	2.28	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:541:GLU:HB3	1:A:565:LEU:HD22	1.98	0.45
1:A:560:TRP:HZ3	1:A:565:LEU:CD1	2.29	0.45
1:B:519:GLN:N	6:B:303:HOH:O	2.50	0.44
1:A:439:LEU:O	1:A:574:MSE:HE1	2.17	0.44
1:A:538:LEU:CD1	1:A:565:LEU:CB	2.94	0.44
1:A:596:PRO:HB2	6:A:134:HOH:O	2.17	0.44
1:A:541:GLU:HB3	1:A:565:LEU:HD21	2.00	0.43
1:B:438:VAL:HG12	1:B:574[B]:MSE:SE	2.70	0.42
1:A:460:LEU:N	1:A:461:PRO:CD	2.82	0.42
1:A:497:LYS:HE3	1:A:516:SER:OG	2.19	0.42
1:A:537:ILE:O	1:A:541:GLU:HG3	2.20	0.42
1:A:560:TRP:CZ3	1:A:565:LEU:CG	3.03	0.42
1:B:473:THR:HA	1:B:474:PRO:HD3	1.92	0.42
1:B:497:LYS:HE3	1:B:516:SER:HB2	2.01	0.42
1:B:479:VAL:CB	4:B:402:GOL:C1	2.95	0.42
1:A:541:GLU:CB	1:A:565:LEU:HD22	2.50	0.41
1:A:606:HIS:HB2	1:A:607:PRO:HD3	2.03	0.41
1:B:566:MSE:HB3	6:B:312:HOH:O	2.20	0.41
1:B:482:GLN:HG3	1:B:531:ARG:NH2	2.36	0.41
1:A:486:ILE:O	1:A:486:ILE:CG1	2.67	0.41
1:A:505:SER:HB2	6:A:34:HOH:O	2.19	0.41
1:A:561:ILE:HG22	1:A:566:MSE:HE2	2.03	0.41
1:A:465[B]:ARG:HD3	6:A:268:HOH:O	2.21	0.41
1:B:530:LEU:CD2	1:B:565:LEU:HD21	2.51	0.41
1:B:533:GLN:HG2	6:B:296:HOH:O	2.20	0.40
1:A:595:PRO:HA	1:A:596:PRO:HD2	1.98	0.40
1:A:439:LEU:HD11	1:A:450:GLU:HB2	2.04	0.40
1:A:568:GLU:HB2	1:B:495:GLU:HG2	2.03	0.40
1:B:485:LEU:HA	1:B:485:LEU:HD23	1.60	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/188 (96%)	173 (96%)	6 (3%)	2 (1%)	12	8
1	B	184/188 (98%)	176 (96%)	4 (2%)	4 (2%)	5	2
All	All	365/376 (97%)	349 (96%)	10 (3%)	6 (2%)	7	4

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	444	SER
1	B	444	SER
1	B	518	ALA
1	B	570	ALA
1	B	568	GLU
1	A	570	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/156 (99%)	152 (99%)	2 (1%)	65	72
1	B	157/156 (101%)	153 (98%)	4 (2%)	42	47
All	All	311/312 (100%)	305 (98%)	6 (2%)	58	59

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	486	ILE
1	A	519	GLN
1	B	480[A]	CYS
1	B	480[B]	CYS
1	B	536	GLN
1	B	594	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	519	GLN
1	A	532	GLN
1	A	536	GLN
1	B	469	GLN
1	B	534	ASN
1	B	536	GLN
1	B	569	HIS
1	B	571	GLN
1	B	576	GLN
1	B	601	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	IPA	B	412	-	3,3,3	0.33	0	3,3,3	1.80	1 (33%)
4	GOL	A	401	-	5,5,5	0.25	0	5,5,5	0.63	0
4	GOL	B	402	-	5,5,5	3.69	4 (80%)	5,5,5	1.97	2 (40%)
5	IPA	A	411	-	3,3,3	0.24	0	3,3,3	1.69	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	401	-	-	4/4/4/4	-
4	GOL	B	402	-	-	3/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	402	GOL	C3-C2	4.98	1.70	1.51
4	B	402	GOL	O2-C2	4.16	1.55	1.43
4	B	402	GOL	O3-C3	3.87	1.58	1.42
4	B	402	GOL	O1-C1	-2.73	1.31	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	402	GOL	O2-C2-C3	3.31	122.88	109.18
5	B	412	IPA	C3-C2-C1	-2.94	91.37	113.38
5	A	411	IPA	C3-C2-C1	-2.80	92.45	113.38
4	B	402	GOL	O2-C2-C1	2.45	119.33	109.18

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	402	GOL	O1-C1-C2-C3
4	B	402	GOL	O2-C2-C3-O3
4	B	402	GOL	O1-C1-C2-O2
4	A	401	GOL	O1-C1-C2-C3
4	A	401	GOL	C1-C2-C3-O3
4	A	401	GOL	O1-C1-C2-O2
4	A	401	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	402	GOL	6	0
5	A	411	IPA	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	180/188 (95%)	0.30	13 (7%) 23 25	21, 37, 57, 72	1 (0%)
1	B	180/188 (95%)	0.43	20 (11%) 12 12	21, 39, 64, 74	3 (1%)
All	All	360/376 (95%)	0.37	33 (9%) 16 17	21, 38, 61, 74	4 (1%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	479	VAL	5.5
1	B	475	GLY	5.2
1	B	568	GLU	4.4
1	B	478	PRO	4.4
1	B	477	ALA	4.3
1	B	479	VAL	4.2
1	A	477	ALA	3.9
1	A	485	LEU	3.7
1	A	614	ASP	3.6
1	B	570	ALA	3.3
1	A	565	LEU	3.2
1	A	478	PRO	3.2
1	B	569	HIS	3.1
1	A	568	GLU	3.1
1	B	567	SER	2.9
1	B	613	LYS	2.9
1	A	567	SER	2.8
1	B	480[A]	CYS	2.8
1	B	485	LEU	2.8
1	A	480	CYS	2.6
1	B	482	GLN	2.6
1	A	475	GLY	2.4
1	A	570	ALA	2.4
1	B	433	GLN	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	517	ASP	2.3
1	B	443	HIS	2.3
1	A	433	GLN	2.2
1	A	564	ASN	2.2
1	B	495	GLU	2.1
1	B	483	PRO	2.1
1	B	534	ASN	2.1
1	B	591	LEU	2.1
1	B	486	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NA	A	423	1/1	0.58	0.46	105,105,105,105	0
4	GOL	A	401	6/6	0.73	0.17	66,68,70,76	0
5	IPA	A	411	4/4	0.78	0.18	55,56,58,60	0
5	IPA	B	412	4/4	0.78	0.25	59,61,62,62	0
4	GOL	B	402	6/6	0.82	0.27	37,40,42,43	6
2	CL	A	422	1/1	0.99	0.07	34,34,34,34	0
2	CL	A	421	1/1	0.99	0.04	27,27,27,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.