



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2023 – 07:25 PM EDT

PDB ID : 2B4J  
Title : Structural basis for the recognition between HIV-1 integrase and LEDGF/p75  
Authors : Cherepanov, P.; Ambrosio, A.L.; Rahman, S.; Ellenberger, T.; Engelman, A.  
Deposited on : 2005-09-24  
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	 5% 67% 14% 19%
1	B	166	 3% 70% 15% 13%
2	C	98	 3% 74% 7% 16%
2	D	98	 5% 72% 9% 16%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	403	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3757 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrase (IN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	135	1048	671	179	193	5	12	0	0
1	B	145	1127	720	195	208	4	10	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	GLY	-	cloning artifact	UNP P12497
A	48	SER	-	cloning artifact	UNP P12497
A	49	HIS	-	cloning artifact	UNP P12497
A	185	LYS	PHE	engineered mutation	UNP P12497
B	47	GLY	-	cloning artifact	UNP P12497
B	48	SER	-	cloning artifact	UNP P12497
B	49	HIS	-	cloning artifact	UNP P12497
B	185	LYS	PHE	engineered mutation	UNP P12497


- Molecule 2 is a protein called PC4 and SFRS1 interacting protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	82	664	412	121	124	7	4	0	0
2	D	82	664	412	121	124	7	18	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	345	GLY	-	cloning artifact	UNP O75475
C	346	SER	-	cloning artifact	UNP O75475
D	345	GLY	-	cloning artifact	UNP O75475
D	346	SER	-	cloning artifact	UNP O75475

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	55	Total	O	0	0
			55	55		
5	B	58	Total	O	0	0
			58	58		
5	C	58	Total	O	0	0
			58	58		
5	D	49	Total	O	0	0
			49	49		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Integrase (IN)




- Molecule 1: Integrase (IN)



- Molecule 2: PC4 and SFRS1 interacting protein



- Molecule 2: PC4 and SFRS1 interacting protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.42Å 60.59Å 71.13Å 90.00° 109.06° 90.00°	Depositor
Resolution (Å)	20.00 – 2.02 19.90 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.2 (20.00-2.02) 99.2 (19.90-2.02)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.87 (at 2.02Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.181 , 0.226 0.180 , 0.224	Depositor DCC
$R_{free}$ test set	1627 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.2	Xtrriage
Anisotropy	0.092	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3757	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/1067	0.77	1/1444 (0.1%)
1	B	0.76	0/1150	0.72	1/1559 (0.1%)
2	C	0.86	0/668	0.85	2/889 (0.2%)
2	D	0.73	0/668	0.87	3/889 (0.3%)
All	All	0.78	0/3553	0.79	7/4781 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	404	ARG	NE-CZ-NH2	7.65	124.12	120.30
2	C	404	ARG	NE-CZ-NH1	-7.16	116.72	120.30
2	C	404	ARG	NE-CZ-NH2	6.94	123.77	120.30
2	D	404	ARG	NE-CZ-NH1	-6.75	116.92	120.30
1	A	102	LEU	CB-CG-CD1	-6.13	100.57	111.00
2	D	351	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	B	167	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1048	0	1059	15	0
1	B	1127	0	1135	24	0
2	C	664	0	699	6	0
2	D	664	0	699	8	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	6	0	8	2	0
4	B	6	0	8	4	0
4	C	6	0	8	2	0
4	D	6	0	8	1	0
5	A	55	0	0	1	0
5	B	58	0	0	0	0
5	C	58	0	0	1	0
5	D	49	0	0	1	0
All	All	3757	0	3624	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:171:HIS:HD2	4:B:403:GOL:H32	1.30	0.96
2:D:401:LYS:HA	2:D:404:ARG:HD2	1.56	0.86
1:B:117:ASN:HD21	1:B:141:ILE:HD11	1.40	0.86
1:B:171:HIS:CD2	4:B:403:GOL:H32	2.09	0.85
1:A:107:ARG:HD2	1:B:107:ARG:HD2	1.62	0.82
2:C:401:LYS:HA	2:C:404:ARG:HD2	1.64	0.79
1:B:171:HIS:HD2	4:B:403:GOL:C3	2.02	0.72
1:B:117:ASN:ND2	1:B:141:ILE:HD11	2.07	0.70
2:C:353:GLN:HG3	4:C:443:GOL:H32	1.74	0.69
1:B:114:HIS:CE1	1:B:138:GLU:OE2	2.50	0.65
2:D:401:LYS:O	2:D:404:ARG:HD3	1.95	0.65
1:B:114:HIS:HE1	1:B:138:GLU:OE2	1.80	0.64
2:C:356:HIS:ND1	4:C:443:GOL:H2	2.14	0.63
1:B:202:ASP:O	1:B:206:THR:CG2	2.49	0.60
2:C:401:LYS:O	2:C:404:ARG:HD3	2.03	0.59
1:B:62:GLN:HG3	1:B:114:HIS:HB2	1.87	0.57
2:D:353:GLN:NE2	5:D:122:HOH:O	2.35	0.57
1:A:178:MET:HE3	2:C:365:ILE:HG12	1.88	0.56
1:A:107:ARG:CD	1:B:107:ARG:HD2	2.33	0.55
1:A:125:THR:HG22	5:A:418:HOH:O	2.07	0.55

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:GLY:O	1:B:183:HIS:HE1	1.89	0.54
1:A:82:GLY:O	1:A:183:HIS:HE1	1.90	0.54
1:B:165:VAL:HG11	1:B:182:ILE:CD1	2.40	0.51
1:A:171:HIS:CD2	4:A:404:GOL:H31	2.46	0.51
1:B:202:ASP:O	1:B:206:THR:HG22	2.10	0.51
2:D:410:GLN:O	2:D:414:GLU:HG3	2.11	0.51
1:B:101:LEU:HD21	1:B:113:VAL:HG11	1.95	0.49
1:B:107:ARG:HG2	1:B:108:TRP:CD1	2.48	0.49
1:B:202:ASP:O	1:B:206:THR:HG23	2.13	0.48
1:B:125:THR:HB	5:C:6:HOH:O	2.12	0.48
1:A:62:GLN:HE22	1:A:151:ILE:H	1.60	0.48
1:A:195:SER:OG	1:A:198:GLU:HG3	2.14	0.48
1:B:57:SER:HA	1:B:60:ILE:HG12	1.97	0.47
1:A:71:LYS:HE2	4:A:404:GOL:H11	1.96	0.47
2:D:353:GLN:HG2	4:D:443:GOL:H32	1.97	0.46
1:A:77:VAL:HG22	1:A:84:ILE:HG22	1.97	0.46
2:D:411:VAL:O	2:D:415:LYS:HG2	2.17	0.45
2:D:405:ARG:H	2:D:405:ARG:HG2	1.59	0.45
2:C:362:SER:OG	2:C:373:CYS:HB2	2.17	0.44
1:B:71:LYS:HZ1	4:B:403:GOL:H2	1.83	0.44
1:B:57:SER:N	1:B:58:PRO:CD	2.81	0.43
1:A:101:LEU:HD21	1:A:113:VAL:HG11	2.00	0.43
1:A:165:VAL:HG11	1:A:182:ILE:CD1	2.48	0.42
1:B:64:ASP:CG	1:B:151:ILE:HG12	2.40	0.42
1:A:165:VAL:HG11	1:A:182:ILE:HD12	2.02	0.42
1:A:181:PHE:O	1:A:185:LYS:HG2	2.20	0.42
1:B:85:GLU:OE1	1:B:107:ARG:NH2	2.53	0.41
2:D:401:LYS:O	2:D:404:ARG:CD	2.66	0.41
1:B:149:GLY:O	1:B:153:SER:HB2	2.21	0.40
1:A:74:LEU:O	1:A:86:ALA:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/166 (78%)	128 (99%)	1 (1%)	0	100	100
1	B	141/166 (85%)	140 (99%)	1 (1%)	0	100	100
2	C	80/98 (82%)	77 (96%)	3 (4%)	0	100	100
2	D	80/98 (82%)	78 (98%)	2 (2%)	0	100	100
All	All	430/528 (81%)	423 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/135 (83%)	109 (97%)	3 (3%)	44	44
1	B	120/135 (89%)	114 (95%)	6 (5%)	24	19
2	C	77/91 (85%)	74 (96%)	3 (4%)	32	29
2	D	77/91 (85%)	74 (96%)	3 (4%)	32	29
All	All	386/452 (85%)	371 (96%)	15 (4%)	32	29

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	ASP
1	A	72	VAL
1	A	125	THR
1	B	64	ASP
1	B	68	LEU
1	B	111	LYS
1	B	125	THR
1	B	151	ILE
1	B	206	THR
2	C	353	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	C	368	LEU
2	C	419	LEU
2	D	368	LEU
2	D	405	ARG
2	D	419	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	114	HIS
1	A	117	ASN
1	A	171	HIS
1	A	183	HIS
1	B	62	GLN
1	B	114	HIS
1	B	164	GLN
1	B	171	HIS
1	B	183	HIS
2	C	353	GLN
2	C	367	ASN
2	C	371	ASN
2	C	410	GLN
2	D	353	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	404	-	5,5,5	0.41	0	5,5,5	1.11	0
4	GOL	D	443	-	5,5,5	0.29	0	5,5,5	1.34	1 (20%)
3	PO4	B	302	-	4,4,4	0.86	0	6,6,6	1.15	1 (16%)
3	PO4	A	301	-	4,4,4	0.92	0	6,6,6	0.78	0
4	GOL	B	403	-	5,5,5	0.47	0	5,5,5	0.99	0
4	GOL	C	443	-	5,5,5	0.34	0	5,5,5	1.66	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	443	-	-	4/4/4/4	-
4	GOL	B	403	-	-	2/4/4/4	-
4	GOL	C	443	-	-	2/4/4/4	-
4	GOL	A	404	-	-	0/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	443	GOL	C3-C2-C1	-2.82	100.72	111.70
4	D	443	GOL	O2-C2-C3	2.49	120.07	109.12
3	B	302	PO4	O4-P-O2	2.13	114.82	107.97

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	403	GOL	C1-C2-C3-O3
4	C	443	GOL	C1-C2-C3-O3
4	D	443	GOL	O1-C1-C2-C3
4	D	443	GOL	C1-C2-C3-O3
4	D	443	GOL	O2-C2-C3-O3
4	B	403	GOL	O2-C2-C3-O3
4	C	443	GOL	O2-C2-C3-O3
4	D	443	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	404	GOL	2	0
4	D	443	GOL	1	0
4	B	403	GOL	4	0
4	C	443	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	135/166 (81%)	0.15	8 (5%) 22 21	26, 35, 49, 56	5 (3%)
1	B	145/166 (87%)	0.07	5 (3%) 45 45	27, 34, 47, 55	3 (2%)
2	C	82/98 (83%)	0.07	3 (3%) 41 41	30, 35, 47, 58	1 (1%)
2	D	82/98 (83%)	0.34	5 (6%) 21 20	30, 37, 48, 57	6 (7%)
All	All	444/528 (84%)	0.14	21 (4%) 31 31	26, 35, 49, 58	15 (3%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	56	CYS	4.8
2	D	425	ASN	4.0
1	B	208	ILE	3.7
2	D	426	MET	3.4
1	A	206	THR	3.3
1	A	150	VAL	3.2
1	A	208	ILE	3.0
2	C	425	ASN	2.9
1	B	207	ASP	2.8
1	B	187	ARG	2.8
1	A	151	ILE	2.7
2	D	377	LEU	2.7
2	D	400	LEU	2.6
2	C	400	LEU	2.6
1	B	119	SER	2.5
1	A	117	ASN	2.5
1	A	58	PRO	2.5
1	A	153	SER	2.5
2	C	424	LYS	2.4
1	B	141	ILE	2.3
2	D	376	ALA	2.1



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	D	443	6/6	0.72	0.24	47,49,51,55	0
4	GOL	B	403	6/6	0.82	0.17	49,56,59,61	0
4	GOL	C	443	6/6	0.85	0.21	39,45,51,55	0
4	GOL	A	404	6/6	0.90	0.19	46,56,58,62	0
3	PO4	B	302	5/5	0.94	0.18	50,50,53,56	0
3	PO4	A	301	5/5	0.95	0.26	63,65,67,67	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.