



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2024 – 06:45 PM EDT

PDB ID : 2BB2  
Title : X-RAY ANALYSIS OF BETA B2-CRYSTALLIN AND EVOLUTION OF OLIGOMERIC LENS PROTEINS  
Authors : Bax, B.; Lapatto, R.; Nalini, V.; Driessen, H.; Lindley, P.F.; Mahadevan, D.; Blundell, T.L.; Slingsby, C.  
Deposited on : 1992-09-21  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	181	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BME	A	176[A]	-	X	-	-
2	BME	A	176[B]	-	X	-	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 1570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA B2-CRYSTALLIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	181	1466	925	256	280	5	0	1	0

- Molecule 2 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
2	A	1	8	4	2	2	0	1
2	A	1	4	2	1	1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	92	Total	O	0	0
			92	92		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: BETA B2-CRYSTALLIN



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.80Å 83.60Å 109.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.10)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.186 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1570	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	0.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.35	4/1511 (0.3%)	2.09	56/2046 (2.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	6

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	70	SER	CB-OG	5.43	1.49	1.42
1	A	27	GLU	CD-OE2	-5.33	1.19	1.25
1	A	174	GLN	CD-OE1	5.26	1.35	1.24
1	A	161	GLN	CD-OE1	5.22	1.35	1.24

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	ARG	NE-CZ-NH1	11.82	126.21	120.30
1	A	107	ASP	CB-CG-OD2	-10.73	108.64	118.30
1	A	93	TYR	CG-CD2-CE2	-9.81	113.45	121.30
1	A	71(A)	ARG	C-N-CA	9.73	146.03	121.70
1	A	169	ARG	NE-CZ-NH2	-8.47	116.06	120.30
1	A	27	GLU	CB-CG-CD	-8.37	91.59	114.20
1	A	85	SER	C-N-CA	8.10	141.95	121.70
1	A	27	GLU	OE1-CD-OE2	7.69	132.52	123.30
1	A	71	ARG	NE-CZ-NH1	7.68	124.14	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	134	TYR	CG-CD2-CE2	-7.28	115.48	121.30
1	A	43	VAL	CA-CB-CG1	7.24	121.76	110.90
1	A	134	TYR	CB-CG-CD1	-7.18	116.69	121.00
1	A	62	TYR	CB-CG-CD2	-7.13	116.72	121.00
1	A	126	VAL	CA-CB-CG2	7.12	121.57	110.90
1	A	125	ARG	O-C-N	-6.98	111.53	122.70
1	A	56	VAL	CA-CB-CG2	-6.92	100.53	110.90
1	A	107	ASP	CB-CG-OD1	6.81	124.43	118.30
1	A	27	GLU	CA-CB-CG	-6.70	98.66	113.40
1	A	79	ARG	CD-NE-CZ	-6.45	114.57	123.60
1	A	103	MET	CG-SD-CE	6.44	110.51	100.20
1	A	64	ARG	NE-CZ-NH1	6.43	123.52	120.30
1	A	62	TYR	CD1-CE1-CZ	-6.34	114.09	119.80
1	A	161	GLN	O-C-N	6.29	133.06	121.10
1	A	168	ARG	NE-CZ-NH2	-6.14	117.23	120.30
1	A	85	SER	CA-C-O	-6.11	107.28	120.10
1	A	-1	ASN	CB-CA-C	-6.08	98.24	110.40
1	A	132	VAL	CA-CB-CG2	6.03	119.94	110.90
1	A	25	LEU	O-C-N	-5.99	113.12	122.70
1	A	93	TYR	CZ-CE2-CD2	5.97	125.17	119.80
1	A	30	GLU	OE1-CD-OE2	5.96	130.45	123.30
1	A	12	GLN	N-CA-CB	-5.82	100.12	110.60
1	A	27	GLU	CG-CD-OE1	-5.81	106.67	118.30
1	A	64	ARG	NE-CZ-NH2	-5.77	117.42	120.30
1	A	171	ARG	NE-CZ-NH2	5.70	123.15	120.30
1	A	97	ASN	O-C-N	-5.61	113.72	122.70
1	A	22[A]	CYS	CA-CB-SG	-5.57	103.97	114.00
1	A	22[B]	CYS	CA-CB-SG	-5.57	103.97	114.00
1	A	51	LYS	CB-CA-C	-5.57	99.26	110.40
1	A	93	TYR	CB-CG-CD1	-5.44	117.74	121.00
1	A	146	LEU	CB-CG-CD2	5.42	120.21	111.00
1	A	59	LYS	C-N-CA	-5.41	110.93	122.30
1	A	71	ARG	C-N-CA	5.41	135.22	121.70
1	A	172	ASP	CB-CG-OD1	5.36	123.12	118.30
1	A	86	GLN	CB-CA-C	-5.35	99.70	110.40
1	A	148	LYS	O-C-N	-5.31	114.18	123.20
1	A	89	LYS	N-CA-CB	-5.30	101.05	110.60
1	A	75	LEU	CB-CG-CD2	5.30	120.01	111.00
1	A	144	TYR	CB-CG-CD2	-5.27	117.84	121.00
1	A	151	TYR	O-C-N	5.21	131.04	122.70
1	A	88	HIS	CA-CB-CG	-5.20	104.75	113.60
1	A	28(A)	GLY	O-C-N	-5.16	114.44	122.70

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	62	TYR	CG-CD2-CE2	-5.06	117.25	121.30
1	A	127	GLN	CA-CB-CG	-5.05	102.29	113.40
1	A	167	VAL	CA-CB-CG2	5.04	118.46	110.90
1	A	116	HIS	CA-CB-CG	-5.00	105.10	113.60
1	A	118(A)	GLN	CG-CD-OE1	-5.00	111.60	121.60

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	69	THR	CB

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	125	ARG	Sidechain
1	A	140	ARG	Sidechain
1	A	171	ARG	Sidechain
1	A	64	ARG	Sidechain
1	A	71	ARG	Sidechain
1	A	71(A)	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1466	0	1389	81	1
2	A	12	0	13	3	0
3	A	92	0	0	5	1
All	All	1570	0	1402	81	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:TRP:NE1	3:A:267:HOH:O	1.84	1.02
1:A:8:GLN:HB2	1:A:12:GLN:HB3	1.46	0.98
1:A:70(A):SER:OG	1:A:71(A):ARG:HB2	1.68	0.93
1:A:8:GLN:CB	1:A:12:GLN:HB3	2.00	0.92
1:A:12:GLN:HA	1:A:12:GLN:OE1	1.77	0.84
1:A:175:TRP:CZ2	3:A:267:HOH:O	2.29	0.84
1:A:175:TRP:CE2	3:A:267:HOH:O	2.27	0.76
1:A:70(A):SER:OG	1:A:71(A):ARG:CB	2.33	0.76
1:A:125:ARG:HG3	1:A:150:ASP:OD1	1.87	0.75
1:A:46:GLU:HG3	1:A:72:THR:HG21	1.69	0.74
1:A:89:LYS:HD2	1:A:106:ILE:HD11	1.68	0.74
1:A:12:GLN:OE1	1:A:12:GLN:CA	2.37	0.72
1:A:8:GLN:HB2	1:A:12:GLN:O	1.90	0.72
1:A:63:PRO:HG2	1:A:64:ARG:HH11	1.55	0.71
1:A:9:GLU:O	1:A:12:GLN:HB2	1.91	0.71
1:A:26:LYS:HG3	1:A:27:GLU:N	2.09	0.68
1:A:49:ASN:O	1:A:50:CYS:HB2	1.93	0.68
1:A:161:GLN:HB3	1:A:162:PRO:HD2	1.74	0.68
1:A:118:TYR:C	1:A:118(A):GLN:HG3	2.13	0.68
1:A:24:ASN:O	1:A:27:GLU:CG	2.44	0.65
1:A:65:TRP:CE3	1:A:69:THR:OG1	2.51	0.63
1:A:24:ASN:O	1:A:27:GLU:HG3	1.98	0.63
1:A:24:ASN:OD1	1:A:26:LYS:HG2	1.99	0.63
1:A:27:GLU:HB2	2:A:176[B]:BME:S2	2.40	0.62
1:A:24:ASN:ND2	3:A:199:HOH:O	2.32	0.61
1:A:69:THR:HG21	1:A:72:THR:O	2.01	0.61
1:A:103:MET:CE	1:A:116:HIS:CB	2.79	0.60
1:A:29:VAL:HG12	1:A:29:VAL:O	2.01	0.60
1:A:8:GLN:HB3	1:A:12:GLN:HB3	1.82	0.60
1:A:127:GLN:HG2	1:A:127:GLN:O	2.04	0.58
1:A:63:PRO:CG	1:A:64:ARG:HH11	2.16	0.58
1:A:103:MET:HE1	1:A:116:HIS:HB2	1.87	0.57
1:A:103:MET:HE3	1:A:116:HIS:HB3	1.87	0.57
1:A:65:TRP:HE3	1:A:69:THR:HG1	1.48	0.56
1:A:37:VAL:HG21	1:A:57:PHE:HB3	1.87	0.55
1:A:120:LYS:HB2	1:A:162:PRO:O	2.06	0.55
1:A:89:LYS:HB2	1:A:106:ILE:HD13	1.88	0.55
1:A:9:GLU:HG3	1:A:10:ASN:ND2	2.22	0.55
1:A:103:MET:HE3	1:A:116:HIS:CB	2.38	0.54
1:A:66:ASP:HB2	1:A:71:ARG:HD2	1.90	0.53
1:A:70(A):SER:O	1:A:71:ARG:HB2	2.08	0.53
1:A:70(A):SER:OG	1:A:71(A):ARG:HG3	2.07	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:SER:HA	1:A:157:PHE:CE2	2.44	0.53
1:A:136:TYR:HB3	1:A:137:PRO:HD2	1.92	0.52
1:A:125:ARG:CG	1:A:150:ASP:OD1	2.57	0.52
1:A:70(A):SER:OG	1:A:71(A):ARG:CG	2.58	0.52
1:A:98:PHE:HZ	1:A:150:ASP:HB3	1.75	0.52
1:A:65:TRP:CZ3	1:A:69:THR:OG1	2.62	0.51
1:A:53:GLU:OE2	1:A:70(A):SER:HB3	2.10	0.51
1:A:103:MET:HE1	1:A:116:HIS:CB	2.39	0.50
1:A:2:LYS:HD3	1:A:17:GLU:OE2	2.12	0.50
1:A:119:GLU:N	1:A:119:GLU:CD	2.65	0.50
1:A:45:TYR:CD1	1:A:50:CYS:HA	2.47	0.49
1:A:103:MET:CE	1:A:116:HIS:CG	2.97	0.48
1:A:136:TYR:HB3	1:A:137:PRO:CD	2.44	0.47
1:A:175:TRP:HZ2	3:A:267:HOH:O	1.82	0.47
1:A:118:TYR:CZ	1:A:119:GLU:HG2	2.50	0.47
1:A:16:HIS:CG	1:A:28:THR:HB	2.49	0.46
1:A:23:PRO:HD2	2:A:176[B]:BME:H11	1.97	0.46
1:A:98:PHE:CZ	1:A:150:ASP:HB3	2.51	0.45
1:A:118:TYR:CE2	1:A:119:GLU:HG2	2.51	0.45
1:A:8:GLN:HB2	1:A:12:GLN:CB	2.31	0.45
1:A:22[B]:CYS:SG	1:A:25:LEU:HD13	2.58	0.44
1:A:89:LYS:HD2	1:A:106:ILE:CD1	2.42	0.44
1:A:22[B]:CYS:SG	1:A:25:LEU:CD1	3.06	0.44
1:A:26:LYS:HG3	1:A:27:GLU:HG2	2.00	0.44
1:A:8:GLN:HB2	1:A:12:GLN:C	2.37	0.44
1:A:65:TRP:CZ3	1:A:71:ARG:HA	2.53	0.43
1:A:27:GLU:HB2	2:A:176[B]:BME:C2	2.49	0.43
1:A:1:PRO:CG	1:A:80:PRO:HG3	2.48	0.43
1:A:9:GLU:CG	1:A:10:ASN:N	2.83	0.42
1:A:83:VAL:HG12	1:A:85:SER:HB2	2.01	0.42
1:A:106:ILE:HG23	1:A:106:ILE:HD12	1.74	0.42
1:A:66:ASP:CB	1:A:71:ARG:HD2	2.49	0.41
1:A:103:MET:CE	1:A:116:HIS:CD2	3.03	0.41
1:A:24:ASN:HB3	1:A:27:GLU:HG3	2.02	0.41
1:A:47:GLN:HB2	1:A:51:LYS:CB	2.50	0.41
1:A:55:PHE:CE1	1:A:69:THR:HG23	2.56	0.41
1:A:161:GLN:HB3	1:A:162:PRO:CD	2.45	0.41
1:A:89:LYS:HB2	1:A:106:ILE:CD1	2.51	0.40
1:A:25:LEU:HD12	1:A:25:LEU:HA	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:ARG:NH1	3:A:189:HOH:O[2_665]	2.13	0.07

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	180/181 (99%)	175 (97%)	5 (3%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	160/159 (101%)	143 (89%)	17 (11%)	<b>6</b> <b>4</b>

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLU
1	A	12	GLN
1	A	15	SER
1	A	25	LEU
1	A	30	GLU
1	A	51	LYS
1	A	64	ARG
1	A	70(A)	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	71(A)	ARG
1	A	74	SER
1	A	86	GLN
1	A	118(A)	GLN
1	A	119	GLU
1	A	120	LYS
1	A	148	LYS
1	A	172	ASP
1	A	173	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	116	HIS
1	A	163	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BME	A	176[B]	1	3,3,3	4.79	1 (33%)	1,2,2	2.00	1 (100%)
2	BME	A	177	1	3,3,3	0.77	0	1,2,2	1.36	0
2	BME	A	176[A]	1	3,3,3	4.58	1 (33%)	1,2,2	2.29	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BME	A	176[B]	1	-	1/1/1/1	-
2	BME	A	177	1	-	1/1/1/1	-
2	BME	A	176[A]	1	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	176[B]	BME	C2-S2	-8.21	1.52	1.80
2	A	176[A]	BME	C2-S2	-7.85	1.53	1.80

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	176[A]	BME	O1-C1-C2	2.29	119.86	110.83
2	A	176[B]	BME	O1-C1-C2	2.00	118.74	110.83

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	176[A]	BME	O1-C1-C2-S2
2	A	176[B]	BME	O1-C1-C2-S2
2	A	177	BME	O1-C1-C2-S2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	176[B]	BME	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.